

CLIMATE SENSITIVITY ANALYSIS OF ALLERGENIC TAXA IN CENTRAL EUROPE WITH NEW ECOLOGICAL INDICATORS

JÁ DEÁK¹, L MAKRA², I MATYASOVSKY³, Z CSÉPE² and B MULADI¹

¹Department of Physical Geography and Geoinformatics, University of Szeged, P.O.Box 653, 6701 Szeged, Hungary
E-mail: aron@geo.u-szeged.hu

²Department of Climatology and Landscape Ecology, University of Szeged, P.O.Box 653, 6701 Szeged, Hungary

³Department of Meteorology, Eötvös Loránd University, Pázmány Péter Street 1/A, 1117 Budapest, Hungary

Summary: The aim of the study is to analyse trends of the pollination season with its duration, start and end dates, as well as trends of the annual total pollen count and annual peak pollen concentration for the Szeged agglomeration in Southern Hungary. The data set covers an 11-year period (1997-2007) including 19 taxa and seven daily climate variables. Trend analysis is performed on both annual and daily bases. Trend analysis on daily bases is a new approach providing information on the annual cycles of trends. For quantifying the strength of the relationship between the annual cycle of the slope of daily trends of the pollen concentrations and the annual cycles of slopes of daily trends of the climate variables an association measure and a multiple association measure are introduced. Individual taxa are sorted into three categories according to their climate sensitivities. These are compared with two novel climate change related indicators, namely risk potential and expansion potential due to the climate change. A novel procedure was applied to separate the effect of the past and current weather conditions in influencing current *Ambrosia* pollen concentration. The potential effect of land use changes on the pollen release of the taxa considered is also discussed using CORINE Land Cover Database.

Key words: pollen, pollen season, trend, ecological indicator, climate change, respiratory allergy

1. INTRODUCTION

Recently, the Earth's ecosystem is experiencing a global warming. Climate change is responsible for the observed northward and uphill distribution shifts of many European plant species. By the late 21st century, distributions of European plant species are projected to have shifted several hundred kilometres to the north (Emanuel et al. 1985, Pearson 2006, Parry et al. 2007, Lindner et al. 2010); forests are likely to have contracted in the south (Penuelas and Boada 2003) and expanded in the north (Leemans et al. 1996, Pearson 2006, Lindner et al. 2010). The rate of change will exceed the ability of many species to adapt. Concerning plant phenology, the timing of seasonal events in plants is changing across Europe due to changes in climate conditions. Between 1971 and 2000, the average advance of spring and summer was 2.5 days per decade. The pollen season starts on average 10 days earlier and is longer than 50 years ago (Feehan et al. 2009).

A recent warming is associated with an earlier onset (Frei 2008, Rodríguez-Rajo et al. 2011), an earlier end date (Stach et al. 2007, Recio et al. 2010), a longer pollen season (Stach et al. 2007, Ariano et al. 2010), an increase in the total annual pollen load (Ariano et

al. 2010, Cristofori et al. 2010), furthermore an increase of patient number sensitized to pollen throughout the year (Ariano et al. 2010).

The scope of the studies is generally limited to only one taxon (Peternel et al. 2006, Alcázar et al. 2011), or a very small number of taxa (García-Mozo et al. 2010, Kaminski and Glod 2011, Rodríguez-Rajo et al. 2011). A comprehensive spectrum of the regional pollen flora was only analysed in three studies, namely in Clot (2003, 25 plant taxa), Damialis et al. (2007, 16 plant taxa) and Cristofori et al. (2010, 63 plant taxa). An overall analysis of the pollen season characteristics for a given source area provides a more reliable picture of the climate sensitivity for each taxon studied based on their different optimum environmental conditions.

Precognition of pollen season characteristics is important for those people suffering from pollen-induced respiratory diseases, who can prepare in due time for days of extreme high pollen load. At the same time, climate change can affect pollen characteristics of different taxa diversely. The object of this paper is to study an extended spectrum of airborne pollen characteristics (19 plant taxa) for the Szeged region in Southern Hungary. Trends of both quantity-related and phenological pollen season characteristics are calculated for each taxon. A multiple association measure (MAM) is introduced that describes how well the annual cycle of daily pollen concentration trends can be represented by a linear combination of annual cycles of climate variable trends. Two novel climate change related indicators, namely risk potential (RP) and expansion potential (EP) due to the climate change are also introduced and these indicators are evaluated for each taxon. Additionally, a novel procedure was applied to separate the effect of the past and current weather conditions in influencing current *Ambrosia* pollen concentration. The potential effect of land use change on *Ambrosia* pollen concentration is also discussed using CORINE Land Cover Database.

2. MATERIALS AND METHODS

2.1. Location and data

Szeged (46.25°N; 20.10°E), the largest settlement in South-eastern Hungary is located at the confluence of the Rivers Tisza and Maros. The area is characterised by an extensive flat landscape of the Great Hungarian Plain with an elevation of 79 m above sea level. The city is the centre of the Szeged region with 203,000 inhabitants. The climate of Szeged belongs to Köppen's *Ca* type (warm temperate climate) with relatively mild and short winters and hot summers (Köppen 1931).

The pollen content of the air was measured using a 7-day recording "Hirst-type" volumetric trap (Hirst 1952). The air sampler is located on top of the building of the Faculty of Arts at the University of Szeged approximately 20 m above the ground surface (Makra et al. 2010). Meteorological variables include daily values of minimum (T_{\min} , °C), maximum (T_{\max} , °C) and mean temperature (T , °C), total solar radiation (TR, $W \cdot m^{-2}$), relative humidity (RH, %), wind speed (WS, $m \cdot s^{-1}$) and rainfall (R, mm). They were recorded in a meteorological station located in the inner city area of Szeged. The data set consists of daily pollen counts (per m^3 of air) of 19 taxa taken over the period 1997-2007. With their Latin (English) names they are: *Alnus* (alder), *Ambrosia* (ragweed), *Artemisia* (mugwort), *Betula*

(birch), *Cannabis* (hemp), Chenopodiaceae (goosefoots), *Juglans* (walnut), *Morus* (mulberry), *Pinus* (pine), *Plantago* (plantain), *Platanus* (plane), Poaceae (grasses), *Populus* (poplar), *Quercus* (oak), *Rumex* (dock), *Taxus* (yew), *Tilia* (linden), *Ulmus* (elm) and *Urtica* (nettle). The 19 taxa studied produce 93.2% of the total pollen amount for the given period. Taxa with the highest pollen levels include *Ambrosia* (32.3%), Poaceae (10.5%), *Populus* (9.6%) and *Urtica* (9.1%), which together account for 61.5% of the total pollen production.

As regards the taxa with the highest pollen concentrations, the *Ambrosia* genus has only one species, namely *Ambrosia artemisiifolia* (Common Ragweed) in the Szeged region that appears both in the urban environment and in the countryside. Ragweed occurs especially frequently west of the city. The ruling north-western winds can easily transport pollen into the city. Since in the sandy region, northwest of Szeged, stubble stripping is not necessary for ground-clearance due to the mechanical properties of sandy soils, *Ambrosia* can spread unchecked. Owing to newly-built motorways around Szeged, several farmland areas have been left untouched for a long time that also favour the expansion of *Ambrosia*. Several species of the Poaceae family occur in the Szeged area, namely *Agropyron repens* (Common Couch), *Poa trivialis* (Rough Meadow-grass), as well as *Poa bulbosa* (Bulbous Meadow-grass) over untouched areas, furthermore *Poa angustifolia* (Narrow-leaved Meadow-grass) and *Alopecurus pratensis* (Meadow Foxtail) in the floodplain, and along the dyke surrounding Szeged. Along the urban lakesides *Phragmites australis* (Common Reed) is the most frequent Poaceae. Furthermore, on short grass steppes of sand, loess and saline areas *Festuca pseudovina* and *Festuca rupicola* also occur. For *Populus* genus, natural species of *Populus alba* (White Poplar) and *Populus canescens* (Grey Poplar) are the most frequent in the city and are characteristic in the floodplain forests along the Tisza and Maros Rivers. In addition, cultivated poplars such as I-273 Poplar and *Populus x euroamericana* (Canadian Poplar) and its variants are frequently planted in urban parklands, public places, as well as along roads in peripheries. At the same time, the *Urtica* genus with its only species of *Urtica dioica* (Common Nettle) has a high frequency in the floodplain forest underwood of the Tisza and Maros Rivers, road-, and channel-sides and in locust-tree plantations around the city. *Urtica* also occurs in the neglected grassy areas of the city.

The remaining species are rare. *Alnus* species are only found in the Botanical Garden of Szeged. The pollen of *Artemisia*, *Cannabis*, Chenopodiaceae and *Rumex* can come from neglected areas of both the city and its surroundings, as well as from stubble pastures. *Betula*, *Juglans*, *Pinus*, *Platanus*, *Taxus* and *Tilia* species have been planted exclusively in public places and gardens; they have no natural habitats in the Szeged region. However, since the 1960s *Pinus* (*Pinus sylvestris* and *Pinus nigra*) species have been extensively planted in the sandy regions north-west of Szeged within the framework of an afforestation programme. Their pollen can easily reach Szeged via the north-western winds. *Morus* is planted along avenues and in public places. *Plantago* species occur in natural grassy areas of both the city and its surroundings. *Quercus* species are planted along the embankment surrounding the city, as well as north of the city. *Ulmus* is planted in the city too; however it is not very common. At the same time, *Ulmus minor* is quite frequent in all landscape types around Szeged on boundaries, road-sides and channel-sides. Its scattered monodominant plantations can appear in loess landscapes and beyond the dykes (saved-side) floodplains, as well as rarely in sand landscapes. In the above-mentioned places the formerly planted *Ulmus pumila*, as an adventive species, also occurs, but its spread is not

characteristic. Large natural stands of *Ulmus minor* together with *Ulmus laevis* live in the oak-elm-ash alluvial forests alongside the River Maros, where both planted and spontaneous stands appear. These species can be found spontaneously even in the willow-poplar alluvial forests, thanks to the mature stands of the Pécska forest on the Romanian side (Deák 2010).

The pollen season is defined by its start and end dates. For the start (end) of the season we used the first (last) date on which 1 pollen grain m^{-3} of air is recorded and at least 5 consecutive (preceding) days also show 1 or more pollen grains m^{-3} (Galán et al. 2001). For a given pollen type, the longest pollen season during the 11-year period was considered for each year.

2.2. Methods

2.2.1. Trend analysis

A common way of estimating trends in data is linear trend analysis. The existence of trends is examined generally by the t -test based on the estimated slopes and their variances. This test, however, may only be used for normally distributed data. Data having probability distributions far from the normal one can be tested against monotone trends by the Mann-Kendall (MK) test (Önöz and Bayazit 2003). Therefore, this method is used here, although the slopes have also been calculated.

It may happen that some trends might have overly complex forms to be well approximated by global linear fits, so nonparametric methods are preferable. Nonparametric methods assume some smoothness of the trends to be estimated. Each version of these techniques results in linear combinations of observations lying within an interval around the points where the trends are estimated. The size of this interval is controlled by a parameter called the bandwidth. There are several versions of such estimators, but local linear fittings have useful properties (Fan 1993). When estimating the trends, the choice of the bandwidths has a crucial role in the overall accuracy. A large bandwidth provides small variances with large biases of the estimates, while a small bandwidth results in large variances with small biases. Thus, an optimal bandwidth producing relatively small variances and small biases has to be found. A technique proposed by Francisco-Fernández and Vilar-Fernández (2004) is used here for the purpose. Note that the local linear fits become globally linear with infinite bandwidths.

2.2.2. Taxon-specific ecological indicators as a basis for introducing new climate change related indicators

In order to evaluate the response of plants to climate change, two indicators were introduced: risk potential due to the climate change (RP) and expansion potential due to the climate change (EP). RP describes the endangerment of the species of different taxa in their present habitats indicating survival potential on their present places. EP shows the capability of the species to move in the landscape, so this term measures the landscape-scale rescue effect. Both indicators were determined for a specific taxonomic group (genus, family) of the most allergenic plants. The species-pool of the Hungarian vegetation was collected according to the Flora Database of Hungary (Horváth et al. 1995). The above-mentioned two new terms were determined on the basis of four main ecological indicators

of the taxa examined. The selected indicators were: temperature requirement due to Zólyomi (TZ-value) (Zólyomi and Précsényi 1964), temperature requirement due to Soó (TS-value) (Soó 1964-1980), heat supply of species interpreted with the climate of the vegetation belts due to Borhidi (TB-value) (Borhidi 1995), as well as the degree of continentality and climate extremity tolerance according to the distribution of species due to Borhidi (CB-value) (Borhidi 1995). Substitution of species within a taxon and habitat shifts were also considered using the Hungarian National Flora Database (Horváth et al. 1995) and field knowledge (Deák 2010) detecting the effect of different climate extremities on the vegetation (Deák 2011a) (Table 1). In order to calculate the new climate change related indicators, besides the above-mentioned ecological indicators, local effects of the prospective changes predicted by recent climate models were also considered. These models count with a 2-4°C increase in annual mean temperature, lower rainfall total in the summer season, annual precipitation decreasing by 150-200 mm and a more extreme rainfall distribution throughout the year for a 100-year time scale until the end of the century (Láng et al. 2007, Czúcz 2009, Faragó et al. 2010).

Table 1 The values of the ecological indicators and the climate change related indicators for the most common allergenic taxa for Hungary

Taxa	² TZ-value	³ TS-value	⁴ TB-value	⁵ CB-value	Risk potential due to climate change (RP)	Expansion potential due to climate change (EP)
<i>Alnus</i>	2, 3, 6	1, 2	3, 4, 5	4	***	-2
¹ <i>Ambrosia</i>	0	0	8	6	*	2
<i>Artemisia</i>	5, 6, 7,	2, 3, 4, 5	6, 7, 8, 9	5, 6, 7, 8, 9	*	2
<i>Betula</i>	3	1, 2	3, 4	3, 4	***	-2
<i>Cannabis</i>	5	0	7	7	*	0
Chenopodiaceae	0, 5, 6, 7	0, 2, 3, 4, 5	5, 6, 7, 8, 9	0, 2, 3, 4, 5, 6, 7, 8, 9	*(few taxa)	1
<i>Juglans</i>	5	4	8	2, 5	*	2
<i>Morus</i>	-	-	7	5	**	-1
<i>Pinus</i>	3	1, 2, 4	4, 8	4, 7	**	-1
<i>Plantago</i>	5, 6, 7	0, 2, 3, 4	5, 6, 7, 8	0, 1, 3, 6, 7, 8	*(few taxa)	1
<i>Platanus</i>	5, 6, 7	4	6, 7, 8, 9	6	*	2
¹ Poaceae	0, 3, 4, 5, 6, 7	0, 2, 3, 4, 5	3, 4, 5, 6, 7, 8, 9	2, 3, 4, 5, 6, 7, 8, 9	*(few taxa) **(few taxa) *** (few taxa)	1
¹ <i>Populus</i>	3, 5	3, 4	5, 7, 8	5, 6, 7	*(few taxa)	1
<i>Quercus</i>	5, 6	3, 4	6, 7, 8	4, 5, 6, 7	*(few taxa)	1
<i>Rumex</i>	0, 5	0, 2, 3, 4	4, 5, 6, 7, 8	2, 3, 5, 6, 7, 8	*(few taxa)	1
<i>Taxus</i>	5	2	5	2	***	-2
<i>Tilia</i>	5, 6	3, 4	5, 7	2, 4, 7	*(few taxa)	1
<i>Ulmus</i>	5	2, 3, 4	5, 6, 7	3, 5, 6	*(few taxa)	1
¹ <i>Urtica</i>	5, 6	0, 4	6, 7	4, 6	*(few taxa)	1

¹**Bold:** taxa with the highest pollen levels;

²**TZ-value: temperature requirement due to Zólyomi (Zólyomi and Précényi 1964, Horváth et al. 1995):**

–: no value is determined; 0: not characteristic; 2: in accordance with woody tundra belt; 3: in accordance with taiga belt; 4: in accordance with broad-leaved/needle-leaved mixed forest belt; 5: in accordance with broad-leaved forest belt; 6: in accordance with sub-Mediterranean forest belt; 7: in accordance with Mediterranean, Atlantic evergreen belt; (Taxa with TZ-value of 1 does not occur.)

³**TS-value: temperature requirement due to Soó (Soó 1964-1980, Horváth et al. 1995):**

–: no value is determined; 0: indifferent species to temperature; 1: highly cold-tolerant, arctic or alpine species; 2: cold-tolerant species; 3: slightly cold-tolerant species; 4: cold-sensitive, warm-needed species; 5: highly warm-needed species

⁴**TB-value: heat supply of species interpreted with the climate of the vegetation belts due to Borhidi (Borhidi 1995, Horváth et al. 1995):**

3: in accordance with sub-alpine or sub-boreal belt; 4: in accordance with montane needle-leaved forests or tajga belt; 5: in accordance with montane broad-leaved forest belt; 6: in accordance with sub-montane broad-leaved forest belt; 7: in accordance with thermophilous forest belt; 8: in accordance with the belts of sub-Mediterranean woodlands and continental steppes; 9: in accordance with Mediterranean evergreen belt; (Taxa with TB-values of 0, 1 and 2 do not occur.)

⁵**CB-value: degree of continentality and climate extremity tolerance in association to the distribution of species due to Borhidi (Borhidi 1995, Horváth et al. 1995):**

0: indifferent species; 1: eu-oceanic species (Atlantic species occurring just exceptionally on continental climate); 2: oceanic species (living mainly in Western-Europe and Western-Central Europe); 3: oceanic-sub-oceanic species (distribution focus on Central Europe); 4: sub-oceanic species (distribution focus on Central Europe, but rarely expanding to East); 5: sub-oceanic - sub-continental intermediate species; 6: sub-continental species (distribution focus on Eastern Central Europe); 7: continental-sub-continental species (distribution focus on Eastern Europe, but occurring in Central Europe, as well); 8: continental species (distribution focus on Eastern Europe reaching only Eastern Central Europe); 9: eu-continental species (Eastern European and Asian steppe species occurring just exceptionally in Central Europe)

2.2.3. Risk potential (RP) and expansion potential (EP) as new indicators due to climate change

RP describes the endangerment of the species of different taxa in their present habitats indicating survival potential of the species with 3 categories. Non-endangered taxa (*) can survive climate change since they contain species for warmer and drier conditions, whereas the climatically endangered taxa (***) have no species in the present flora for future changed conditions. In the first case, change of species within a taxon in a certain landscape could help the adaptation of the taxon to global warming, whereas in the latter case the lack of heat tolerant species can lead to the disappearance of a given taxon. The wider tolerance range (the more ecological indicator values, i.e. TZ-, TS-, TB-, or CB-values) (Table 1) and the more species (especially heat and drought tolerant species) a taxon has, the less exposure to climate change it has. Moderately endangered taxa (**) could survive partly in their places, but populations of some species may decrease regionally.

Three variables must be considered for a given taxon: (1) the number of species within a taxon, (2) the value range of the ecological indicators of the species within a taxon and (3) the number of heat and drought tolerant species within a taxon (this is the most important factor). For example, grasses (Poaceae) have a lot of species [see: (1)] with a wide value range of the ecological indicators of their species [see: (2)], and many of them are heat tolerant [see: (3)], so they will have enough species to adapt to the changes. However, the answers of species can be different to the climate change (all three categories of RP can occur for Poaceae) (Table 1). If a taxon contains mainly warm- and drought tolerant species (e.g. *Ambrosia*, *Juglans* and *Platanus*) less species need adaptation, so they

are less endangered taxa. If a taxon has only a few species and neither of them favours warmer and drier conditions the chance for its extinction is significantly high (***) (e.g. *Betula* and *Alnus*) (Table 1).

EP shows the capability of the species to move in the landscape thus characterizing the rescue effect. If a taxon belongs to several categories of RP, these categories can be grouped into different classes on the basis of field experiments (Horváth et al. 1995, Deák 2010, 2011b, Bölöni et al. 2011). This feature is described with 5 classes as a wide range of responses are expected due to the different climate-tolerance of the species-pool of taxa. These classes are as follows. (0): Taxa not influenced by global warming. They could survive and their distribution area will remain about the same. (+1): Taxa not influenced by global warming, but for some species area-increase and for some others area-decrease is possible. They can survive the changes through moving in the landscape, but their expansion is limited. (+2): Taxa significantly influenced by global warming. For some species area-increase is expected. They show the best adaptation to the climate change as they will not just survive but can spread well in the landscape. For example, potential spread of species with (*) (e.g. *Ambrosia*, *Juglans* and *Platanus*) is expected in the landscape, so +2 value is given for them showing that their significant area increase is expected as a response for the awaited climate change. (-1): For some species regional area-decrease is possible. They can survive in a few places, their spread in the landscape is limited and area-decrease is expected. (-2): Taxa significantly influenced by global warming. For the majority of species area-decrease is expected. They have the smallest adaptation capability; they will gradually disappear and even the rescue effect in some refuges is doubtful (Table 1). For instance, *Betula* and *Alnus* are endangered taxa indicated with (***) in the RP system, so their EP value is the lowest meaning that area-decrease is expected as a result of climate change. These five categories can be useful not only to evaluate the future distribution of allergenic species groups, but they can also help to reveal the possible changes in endangered plant groups and invasive species (nature conservation aspect), as well as the spread of weeds (agricultural aspect).

On the whole, RP concentrates on the local survival specifying what will happen with the species in their present habitats, while EP concentrates on the landscape-level response. The scale of the two indicators is thus different. Note that this work is the first attempt to reveal the climate sensitivity of the main species groups of vegetation since the database applied has not been available before and sufficient field-knowledge has now been collected by several botanists.

2.2.4. Multiple association measure (MAM) and its association with RP and EP

We examined whether there were any associations between the annual cycles of daily slopes of pollen concentration trends and the annual cycles of daily slopes of climate variables trends. Here an association measure (AM) is used to characterize these relationships by calculating the correlations between the annual cycles of slopes obtained by the nonparametric trend estimation procedure of Section 2.2.1. This quantity will not be referred to as a correlation because correlation is defined for random variables, but in this case similarities between deterministic functions (annual cycles) have to be quantified. In addition, an overall measure called multiple association measure (MAM) characterizing how well the annual cycle of the slope of a pollen concentration trend can be represented by a linear combination of annual cycles of slopes of climate variable trends was also introduced. MAM varies between zero and unit approaching the unit under increasing

accuracy of this above mentioned representation. Technically, MAM is calculated as a multiple correlation between a random variable and a number of other random variables, but again it should not be considered as correlation. AM and MAM are based on elementary considerations of linear algebra (see e.g. chapter 5.15 in Meyer 2001) because an annual cycle of slopes of daily trends covering an n -day period can be considered an n -dimensional vector. AM and MAM relate to angle between the vector corresponding to the annual cycle of the slope of a pollen concentration trend and the hyperplane defined by vectors corresponding to the annual cycles of slopes of climate variable trends.

MAMs do not necessarily harmonize with RP and EP for the given taxa. EPs were determined using RPs based on the ecological indicators of the species-pool; hence, these indicators are in strong association. These latter two plant associated indicators count (a) with the intra-taxonic species change, (b) with a higher range of species (species in and around the Carpathian basin), (c) with the transformation of the abiotic features of habitats, (d) with the moving capability of the species and (e) with the rescue effect of habitats due to special microclimates. At the same time, MAMs consider only the climate sensitivity of the taxa for a given time period, area and their species-pool. Furthermore, RP and EP are suitable to detect regional changes for longer (centuries, millennia) time periods, whereas MAMs can be used for observing local changes for shorter (decades) periods (Tables 1, 3, 4).

2.2.5. Factor analysis and special transformation

Factor analysis identifies any linear relationships among subsets of examined variables and this helps to reduce the dimensionality of the initial database without substantial loss of information. First, a factor analysis was applied to the initial datasets consisting of 9 variables (8 explanatory variables and 1 resultant variable defined by the daily *Ambrosia* pollen concentration) in order to transform the original variables to fewer variables. These new variables (called factors) can be viewed as latent variables explaining the joint behaviour of past and current meteorological elements – current *Ambrosia* pollen concentration variables. The optimum number of retained factors can be determined by different statistical criteria (Jolliffe 1993). The most common and widely accepted one is to specify a least percentage (80%) of the total variance in the original variables that has to be achieved (Liu 2009). After performing the factor analysis, a special transformation of the retained factors was made to discover to what degree the above-mentioned explanatory variables (4 climatic variables in the past and the same 4 climatic variables on the actual day) affect the resultant variable (daily *Ambrosia* pollen concentration), and to give a rank of their influence (Jahn and Vahle 1968). When performing factor analysis on the standardized variables the factor loadings received are correlation coefficients between the original variables and, after rotation, the coordinate values belonging to the turned axes (namely, factor values). If the resultant variable is strongly correlated with the factor and an influencing variable is highly correlated with the same factor, then the influencing variable is also highly correlated with the resultant variable. Accordingly, it is advisable to combine all the weights of the factors, together with the resultant variable, into one factor. Namely, it is effective to rotate so that only one factor has great load with the resultant variable. The remaining factors are uncorrelated with the resultant variable; that is to say, are of 0 weight (Jahn and Vahle 1968). This latter procedure is called special transformation.

3. RESULTS

3.1. Trend analysis

Only a few trends have been clearly identified compared to the total number of annual MK tests performed (Table 2). It is not surprising as the inter-annual variability of the characteristics studied is quite high, while the size of the data set is quite small. Therefore, MK tests are performed and linear trends are estimated for each particular day of each pollen season of all 19 taxa considered using 11-element pollen concentration data sets corresponding to the 11-year study period. This kind of trend analysis provides information on the annual cycles of trends. In the absence of a trend for each day of the pollen season, the MK test values are distributed normally with zero expectation and unit variance. Therefore, deciding on the existence of a trend is identical with the problem of deciding whether the annual mean of daily MK test values corresponds to the expectation zero. The classical *t*-test has been simplified for the purpose as the variance is known (unit), but modified based on the autocorrelations among the consecutive MK test values. First order autoregressive (AR(1)) models are used to describe these autocorrelations. Averaging values of daily slopes of linear trends over the pollen seasons gives rates of change of the total annual pollen counts (TAPC). Note that the trend analysis carried out on a daily basis detects much more significant trends of TAPC than the trend analysis on an annual basis (Table 2).

Table 2 Change in the total annual pollen count (TAPC) (pollen grains·m⁻³ / 10 years), annual peak pollen concentration (APP) (pollen grains·m⁻³ / 10 years), start, end and duration of the pollen season (days / 10 years) calculated by using linear trends. Significant values on annual basis are denoted by *** (1%), ** (5%) and * (10%). Significant values on daily basis are denoted by +++ (1%), ++ (5%) and + (10%).

Taxa	Mean total annual pollen counts	TAPC	APP	Pollen season		
				Start	End	Duration
<i>Alnus</i>	505	-214	-59*	18	16	-2
¹ <i>Ambrosia</i>	7826	-1170	230	14*	-9	-22
<i>Artemisia</i>	772	-61	-133	-4	15	19
<i>Betula</i>	901	-60	0	-1	2	3
<i>Cannabis</i>	432	47 +	-4	8	36**	28
Chenopodiaceae	854	-175 ++	-9	-2	3	5
<i>Juglans</i>	284	253 +++	30*	-8	-7	1
<i>Morus</i>	667	400 +++	44	-7	-4	3
<i>Pinus</i>	500	-194 +++	-20	-2	-1	0
<i>Plantago</i>	409	91 ++	3	-23**	-19	4
<i>Platanus</i>	400	271 ++	48	-7	-3	4
¹ Poaceae	2552	176	43	-10	17*	27***
¹ <i>Populus</i>	2322	2981** +++	610**	-2	3	4
<i>Quercus</i>	423	236 +	25	4	9	5
<i>Rumex</i>	462	-505 +++	-45	-11**	3	15
<i>Taxus</i>	572	697* +++	59	-4	29***	32
<i>Tilia</i>	225	-65 +	-1	-4	-1	3
<i>Ulmus</i>	260	-160 +++	-12	5	-13	-18
¹ <i>Urtica</i>	2200	1183* +++	25	-13**	18**	31***

¹**Bold:** taxa with the highest pollen levels

Needless to say, the daily MK test statistics have a big variability. Therefore, daily MK test values are smoothed with the nonparametric regression technique outlined in Section 2.2.1. In the absence of a trend for each day the estimated bandwidth is extremely large (practically infinite) producing a line close to zero because the local linear approximation to the annual cycle of the daily trends becomes globally linear. Hence, well-defined finite bandwidths obtained for every taxon indicate trends even for *Alnus*, *Ambrosia*, *Artemisia*, *Betula* and Poaceae, the 5 taxa not exhibiting overall trends on yearly basis at even 10% significance level. The nonparametric regression technique was used also to estimate annual cycles of the slopes of daily trends.

Trends of taxa (Table 2) related to trends of climatic variables (Fig. 1) can be explained as follows. *Alnus* and *Betula* occur around Szeged with very little populations (especially *Alnus*). Due to the tolerance range of these taxa increasing temperature and drying climate do not favour them, so their decreasing pollen trend would not be surprising as *Alnus glutinosa* – the only representative of this taxon appearing just in few bogs in the Great Hungarian Plain and parks in Szeged – and *Betula pendula* (planted in parks) like a wetter, more humid, balanced climate. Due to their low occurrences the decrease of their TAPC cannot be significant in the trends. However, we are now in a preliminary period of the climate change (Parry et al. 2007) to which they could have been already adapted. Furthermore, the trend back-shots (e.g. individual years with extreme high precipitation in the decreasing trend) could ensure enough water for their survival and adaptation, but it is questionable whether these back-shots will be enough if the trends continue in the future.

Pinus show a clear decreasing trend as a result of non-suitable choice of species because foresters and gardeners plant non-heat tolerant *Pinus* species in many cases. In the forest plantations of Kiskunság sand-ridge the co-plantation of *Pinus sylvestris* favouring a wetter, cooler climate and the more heat tolerant sub-Mediterranean *Pinus nigra* is common. The increasing temperature and the lack of water, especially in summer does not suit *Pinus sylvestris* and a substantial decrease of TAPC for *Pinus* can only be partially compensated by the increase of pollen counts of *Pinus nigra*.

TAPC for *Tilia* show a slight decreasing tendency. This taxon is not widely represented in the forests of the Great Hungarian Plain, but it is quite common in the parks and gardens of Szeged city. Of the *Tilia* species *Tilia platyphyllos* is often planted, which tolerates the warmer and drier climate less compared to *Tilia cordata* and *Tilia tomentosa*, which are more heat tolerant, especially the latter. *Tilia tomentosa* is represented along the Illyrian-Dacical pincer which is a route for sub-Mediterranean species approaching the Carpathian basin from the south. Note that the *Tilia* species can be found as ornamental vegetation mainly around Szeged, but they also occur within a 100-150 km range in natural forests in the surroundings of Szeged (Mecsek Mountain and South Transdanubian Hills west from Szeged, Banatian and Partiumian Mountains east of Szeged in Romania and Fruska Gora Mountains in Serbia, south of Szeged).

The lack of humidity and the increasing temperatures can be a limiting factor also for *Ulmus* as a notable decrease in its TAPC was observed. Stocks of its most common species *Ulmus minor* grow in dry places without shading trees like oak; they appear several times alone alongside roads and channels. This species together with *Ulmus laevis* can be found in the hard-wood floodplain forests as well. These taxa can also occur in parks where *Ulmus glabra* is sometimes planted. *Ulmus* is a spring pollinating species. A warming may involve higher spring temperatures and maximum temperatures. Its pollination season therefore can begin earlier and can last longer during this season mainly because in

springtime no limitation of available groundwater can be detected. From the melted snow, from the springtime rainfall and even from the floods enough water is available for this genus. However, in dry places the lack of shading trees, while in floodplains the absence of floods and the lower level of the summer groundwater can limit their pollination.




Fig. 1 Annual cycles of the slopes of daily linear trends for maximum temperature (T_{max}), mean temperature (T), rainfall total (R), relative humidity (RH) and wind speed (WS) for *Ambrosia* (solid), *Poaceae* (dash), *Populus* (dot) and *Urtica* (dash dot)

Change of TAPC for *Artemisia*, Chenopodiaceae and *Rumex* calculated on yearly basis is not significant. However, Chenopodiaceae and *Rumex* show decreasing trends when calculating on daily basis. These plants are typical species of young fallows, which appeared after the change of political system at the beginning of 1990s, especially in sand landscapes. However, due to spontaneous regeneration facilitated by grazing and mowing these new stocks began to disappear as these sand fallows turned into sand steppe grasslands (Deák 2010) and only the populations associated with natural habitats or settlements remained. At the same time, the abandonment of arable lands have decreased during the last 10 years as the extension of fallows was reduced by in-buildings and forest plantations, respectively. The strong decrease of *Rumex* can be explained by the fact that *Rumex* (especially *Rumex crispus*) often appears in the summer in depressions covered with inland water in springtime, which were transformed into arable lands. The lack of rainfall can hence reduce their stocks. However, wetter years do not favour the pollination of this genus since the plants' anther is not dry enough to open, furthermore rainfall can wash out its pollen from the air.

In contrast, TAPC of *Plantago* shows a clear increase, which can also be explained by the above-mentioned regeneration processes of fallows. *Plantago* species (especially *P. media* and *P. lanceolata*) are typical plants of old fallows regenerating into sand steppe grasslands, but also occur frequently in urban grasslands. It is typical that weeds of *Artemisia*, Chenopodiaceae and *Rumex* disappear in the later stages of fallow-succession, while steppe species like *Plantago* appear especially in the treated (mown, grazed) fallows. The fallows of the sand-ridge around Szeged have apparently reached the stage when *Plantago* became more frequent.

The duration of the pollen season of Poaceae is significantly increasing due to the warming climate, but TAPC exhibits no significant trend, which can also be in association with the regenerating fallows. The older fallows are all characterized by a huge coverage of grasses, so fallow regeneration led to a slight extension of grasslands during the last 10 years. In warmer years the pollination season of Poaceae can be substantially longer (Makra et al. 2012), but without higher pollen concentrations due to decreasing summer rainfall amounts.

Populus indicates a substantial increase of TAPC. This can be the result of its wide climate tolerance as both wet- (e.g. *P. nigra* and *P. canescens*) and drought tolerant species are represented in the landscape. Especially *Populus alba* living in floodplains, sand lands and parks shows a great adaptation potential. *Populus* (both wild and cultivated types) were planted widely in the sand lands west from Szeged and in the floodplains. Plantation of these species has not yet stopped during the last 10 years. Besides the locust-tree (*Robinia pseudo-acacia*) they are the most favoured trees for the plantation of forests. The stands planted in the last decades are in mature state, so they can pollinate on a high level.

Quercus (mainly *Quercus robur*) was not planted so widely during the last 10 years, but mainly in the 1960s and partly in the 1970-1980s significant oak plantations appeared southeast from Szeged in the saved-side floodplains and in the active floodplain of river Maros. The stands became more and more mature during the last decades, so they are on the level of their full pollination potential. As *Quercus robur* is a continental species with wide climate tolerance the increasing temperature can help its generative processes limited by the available water. These forests produce the average pollen count of this habitat-type. It must also be considered that these forests are managed. In a young forest there are smaller trees producing a certain amount of pollen. Some of them are then cut off in order

to enhance the growth of the healthier individuals. They grow higher but the number of trees decreases significantly and so no major pollen count increase is expected due to their growing age. This is the reason why only a weakly significant increase in their pollen release occurs. Climate variables and pathogens can have a more important role in controlling the pollen counts from these stands and these two factors are also associated.

Surprisingly, the TAPC of *Taxus* showed a remarkable increase during the last 10 years. This species does not appear in natural communities around Szeged, only in parks and gardens. *Taxus baccata* – as a representative of this taxon in Szeged – is a Western European species favouring a more humid, balanced oceanic climate, which is warmer in the winter and springtime compared to the continental climate of Hungary. However, this trend is influenced by the city since the spring begins earlier and a longer pollination period with more pollen grains is expected due to the urban heat island. In this way, freezing temperatures can occur beyond the city, while inside above-zero temperatures are more typical.

Highly significant increase was observed for TAPC and the duration of the pollen season of *Urtica*. *Urtica dioica*, the only representative of this species around Szeged, has a wide climate tolerance. It can be found both in dry and wet habitats. The under-use of urban habitats, the frequent plantation of locust-tree (*Robinia pseudo-acacia*) and the increasing area of fallows also contributed to the expansion of their population. Their pollination is also promoted by increasing maximum temperatures facilitating an earlier start and later end of its pollen season.

Growing trends were observed for all heat tolerant, non-endemic taxa. TAPC of *Cannabis* (originated from South Asia) shows just a slight increase, but stronger increase was observed for *Juglans*, *Platanus* (originating from the Mediterranean) and *Morus* (originating from East Asia).

Ambrosia (represented only with one species) shows no significant trend as a moderate warming is favourable for this taxon, but the lack of available water during the hottest summer period can limit its pollination as the plant concentrates on preserving water and maintaining its vegetative life functions instead of the generative. *Ambrosia* appears year by year in stubble fields, especially in sand landscapes and in abandoned places around settlements. The populations of young fallows represent just a smaller part of their population in the landscape.

3.2. Analysis of the individual taxa based on MAM, RP and EP

Possible future change of taxa due to the anticipated climate change based on RP, EP (Table 3) and MAM (Table 4) can be explained as follows. *Alnus*, *Betula* and *Taxus* are endangered species in their existence according to their RP (***), as well as they are highly sensitive according to their MAM values (+++). This is because they live on the edge of their distribution area in Hungary. Hence, global warming can affect them negatively since they cannot tolerate warm and dry climate for a longer time and they can become extinct in several habitats and other competitors can be more successful. Especially *Alnus* is sensitive to the lack of water but the urban heat island can help its pollination as the pollen season can start earlier. *Betula* favours a much cooler climate according to its distribution area and climate tolerance indicators. Therefore, *Betula* adapted to cooler springs does not tolerate earlier springs with higher temperatures. The anticipated warmer climate may thus reduce its pollen release. Increased temperatures can be more severe limiting factors for *Betula* than the lack of water considering its existence.

Ambrosia is unaffected according to its MAM value. However, its higher potential increase is expected due to its ecological indicator values and high climate-tolerance. Namely, this genus can adapt well to dry and hot conditions, but is highly influenced by future land use.

Artemisia species are heat tolerant, so their EP is high; even Mediterranean and more continental species could appear in the Carpathian basin. The EP of agricultural weeds can be high on fallows that appear fast in all landscapes in unfavourable weather conditions for farming. *Artemisia* pollen is partly released by *Artemisia santonicum*, a natural dominant species of *Artemisia* short grass alkali steppes – main habitats of saline grasslands formed on loess (Deák 2010). These habitats and this species are proved to be very climate-sensitive – in contrast with *Artemisia* weeds. Its reason is the leaching out of their solonetz soils, due to the drainage of saline grasslands and the decrease of rainfall resulting in the decrease of salty groundwater-table.

Table 3 Association measure (^aAM) between the annual cycles of the daily slopes of pollen concentration trends and the annual cycles of the daily slopes of climate variables trends

Taxa	T _{min}	T _{max}	T	R	TR	RH	WS	^b MAM
<i>Alnus</i>	0.718*	0.775*	0.742*	0.313	-0.028	-0.620*	-0.455	0.992
¹ <i>Ambrosia</i>	0.100	0.207	-0.641*	0.398	0.049	0.087	0.223	0.827
<i>Artemisia</i>	-0.249	0.676*	-0.486	0.140	-0.004	-0.230	-0.049	0.998
<i>Betula</i>	-0.689*	-0.192	-0.544*	-0.663*	-0.006	0.542*	0.070	0.973
<i>Cannabis</i>	0.602*	-0.559*	0.763*	-0.531*	-0.152	0.106	-0.147	0.993
Chenopodiaceae	0.071	0.306	-0.869*	0.644*	0.047	0.112	0.307	0.965
<i>Juglans</i>	0.271	-0.392	-0.466	0.613*	-0.129	-0.726*	0.452	0.925
<i>Morus</i>	0.329	-0.668*	-0.874*	0.821*	-0.216	-0.893*	0.684*	0.978
<i>Pinus</i>	0.093	0.144	0.241	-0.269	-0.160	-0.294	-0.079	0.963
<i>Plantago</i>	0.183	-0.642*	-0.093	0.337	-0.131	0.371	0.490	0.947
<i>Platanus</i>	0.308	-0.265	-0.354	0.368	-0.020	-0.576*	0.328	0.948
¹ Poaceae	-0.088	-0.649*	-0.816*	0.826*	-0.057	0.309	0.643*	0.959
¹ <i>Populus</i>	0.361	0.358	0.395	0.407	-0.093	-0.378	-0.349	0.869
<i>Quercus</i>	-0.046	0.165	0.360	0.616*	-0.076	-0.640*	-0.062	0.911
<i>Rumex</i>	-0.093	-0.026	0.450	-0.244	-0.060	-0.365	-0.087	0.979
<i>Taxus</i>	0.618*	0.305	0.428	0.446	0.010	0.009	-0.264	0.985
<i>Tilia</i>	0.284	-0.378	-0.171	0.327	0.062	-0.106	0.428	0.973
<i>Ulmus</i>	0.381	0.565*	0.462	0.063	-0.069	-0.766*	-0.256	0.934
¹ <i>Urtica</i>	-0.467	0.612*	0.451	-0.396	0.076	-0.580*	-0.705*	0.827

¹**Bold:** taxa with the highest pollen levels;

T_{min}: minimum temperature (°C), T_{max}: maximum temperature (°C), T: mean temperature (°C),

R: rainfall (mm), TR: total radiation (W·m⁻²), RH: relative humidity (%), WS: wind speed (m·s⁻¹);

^aAM (association measure): reflects the strength of the relationship between the annual cycle of the daily slopes of pollen concentration trends and the annual cycles of the daily slopes of climate variables trends for each individual taxon;

^bMAM (multiple association measure): describes how well the annual cycle of the slope of the daily pollen concentrations can be represented by a linear combination of the annual cycles of the slopes of the daily climate variable trends. MAM varies between zero and one, approaching one with increasing accuracy of this above-mentioned representation (Meyer 2001);

*AM >|0.5| indicates a strong association

Cannabis is a heat tolerant species but too high maximum temperatures can be a limit for its pollen production due to water shortage. Future drier conditions are expected during its pollen season, though the June rainfall peak (Medard-day rainfall) overlaps it.

The MAM of Chenopodiaceae indicates high sensitivity, but its response to climate change varies according to its species as it has a wide range of species-pool. Both increase

and decrease of its species-pool are expected. These plants frequently appear in areas affected by inland water around Szeged that can disappear due to a drying climate.

Juglans being heat tolerant shows a potential increase due to warming, and according to its MAM, has medium climate sensitivity. Rainfall is a major limiting factor. Warming can help this species as it happened in interglacial times, but a certain minimum rainfall is required. This species cannot be seen over the driest areas of the Mediterranean, but still has a relatively large EP in Hungary.

Table 4 Climate change related indicators and significance of the different pollen season characteristics for each individual taxon

Taxa	RP	EP	² MAM	³ TAPC by linear trend	⁴ APP	⁵ Pollen season			⁶ TAPC via daily linear trend
						onset	end	duration	
<i>Alnus</i>	***	-2	+++		(-10)				
¹ <i>Ambrosia</i>	*	2	+			(+10)			
<i>Artemisia</i>	*	2	+++						
<i>Betula</i>	***	-2	+++						
<i>Cannabis</i>	*	0	+++				+5		(+10)
Chenopodiaceae	*(few taxa)	1	+++						-5
<i>Juglans</i>	*	2	++		(+10)				+1
<i>Morus</i>	**	-1	+++						+1
<i>Pinus</i>	**	-1	+++						-1
<i>Plantago</i>	*(few taxa)	1	++			-5			+5
<i>Platanus</i>	*	2	++						+5
¹ Poaceae	*(few taxa) *** (few taxa)	1	+++				(+10)	+1	
¹ <i>Populus</i>	*(few taxa)	1	+	+5	+5				+1
<i>Quercus</i>	*(few taxa)	1	++						(+10)
<i>Rumex</i>	*(few taxa)	1	+++			-5			-1
<i>Taxus</i>	***	-2	+++	(+10)			+1		+1
<i>Tilia</i>	*(few taxa)	1	+++						(-10)
<i>Ulmus</i>	*(few taxa)	1	++						-1
¹ <i>Urtica</i>	*(few taxa)	1	+	(+10)		-5	+5	+1	+1

¹**Bold:** taxa with the highest pollen levels;

²**MAM (multiple association measure):** + low sensitivity; ++ medium sensitivity; +++ high sensitivity;

³**TAPC by linear trend:** change in the total annual pollen count calculated by using linear trends;

⁴**APP:** change in the annual peak pollen concentration calculated by using linear trends;

⁵**Pollen season:** change of start, end and duration of the pollinations season calculated by using linear trends;

⁶**TAPC via daily linear trend:** change in the total annual pollen count calculated by using daily linear trends;

^{3,4,5,6} ±1, ±5: a significant increasing/decreasing trend at the 1%, 5% probability levels; (±10): a tendency of trend at the 10% probability level

Morus is assigned to the moderately endangered category (**). Lack of rainfall and too high temperature are a barrier for their pollen production during their pollen season

(summer). Its AM and MAM values emphasize that decreasing rainfall can decrease their pollination intensely. On its original distribution area in China, high temperatures are accompanied with spring-time or monsoon rainfalls. This high precipitation amount is missing in Central Europe. Hence, *Morus* try to preserve as much water as possible in summer.

Pinus is endangered moderately according to its RP (**) but MAM shows much higher sensitivity (+++). This is because *Pinus sylvestris* representing a big portion of this species cannot tolerate warming climate. However, change in species composition, plantation and the appearance of Mediterranean species can make this genus more adaptive for the expected changes.

Plantago, *Quercus* and *Ulmus* have medium sensitivity (++) according to their MAM values and due to their diversified species-pool they can react well to climate change with intra-taxonic species changes. RP values show that they are moderately endangered or not endangered, and for heat tolerant, sub-Mediterranean or continental species further expansion can be awaited at the expense of present species. It can be expected mostly for genera *Ulmus* and *Quercus*, where more heat tolerant species also exist in the landscape or around the Carpathian-basin.

A medium sensitivity indicated by MAM (++) is observed for *Platanus*. It stands warm climate, but a shortage of water can be a limiting factor. This species appears mainly in mountain valleys alongside streams in the Mediterranean, where a certain amount of water is available. It has a potential for expansion in Hungary but not in every landscapes.

Poaceae show high sensitivity according to MAM (+++), since the available water and high temperature can mean limits for them. However, the species-pool of this family is the widest among the studied plant groups, so there will be species to substitute the actual grasses and even species from the Mediterranean and the more continental areas can reach the Carpathian-basin in the future. This means a high risk for the present species, but intra-taxonic re-assembly could happen. Shortage of water and too high temperatures can cause lower pollen production in natural grasslands and also in crops produced in arable lands. Certain species in certain places and time periods can suffer from climate change, but the change in species composition will give good chance for the survival of this family.

Rumex is moderately endangered in the Szeged region as it can give a wide range of responses to climate change according to its species-pool. Since *Rumex* species around Szeged live rather in semi-humid conditions favouring inland water covered areas (see *Rumex crispus*, the most common *Rumex* species around Szeged) climate change could affect them intensely due to water shortage.

The *Tilia* genus can give a wide range of responses according to its species-pool. Potential increase, especially for *Tilia tomentosa*, is expected in the Great Hungarian Plain, but this heat tolerant species currently occurs only in the parks of Szeged. The existing species-pool is not favourable for the expected climate change. Especially *Tilia platyphyllos* characterized by high sensitivity according to its MAM (+++) will not stand the warming. *Tilia cordata* could survive better, but its natural stock is very small.

Urtica and *Populus* have a wide climate-tolerance, so they are not climate-sensitive according to MAM. Both genera could increase their population in the future. They are not endangered or only certain species are moderately endangered by warming. *Urtica dioica* is not endangered according to its RP and even population increase is expected. For the extremely rare *Urtica kioviensis* living in boggy wetlands major decrease is expected in the

Carpathian basin. The better climate tolerance of *Populus* can be explained by the wide adaptation of its different species.

4. DISCUSSION AND CONCLUSIONS

Climate change can modify the pollen season characteristics of different allergenic taxa in diverse ways and can exert a substantial influence on habitat regions. In our best knowledge, only three previous studies (Clot 2003, Damialis et al. 2007, Cristofori et al. 2010) analysed comprehensive spectra of the regional pollen flora. The present study analyses one of the largest spectra with 19 taxa. Our study can be considered unique in the sense that trends of pollen concentration data for each taxon and those of all seven climate variables are calculated on a daily basis. This kind of trend analysis provides information on annual cycles of daily slopes of trends.

On a yearly basis only *Populus*, *Taxus* and *Urtica* show a significant increase of the total annual pollen count. *Populus* and *Juglans* display the most important increase, while *Alnus* exhibits the biggest decrease of the annual peak pollen counts. Poaceae and *Urtica* show a significant increase in the duration of the pollen season. Based on the 5% level, 11 of the 19 taxa indicated significant trends of the total annual pollen count, and 7 of these 11 trends is increasing on a daily basis (Table 2). Phenological characteristics (onset, end and duration of the pollen season) show changes only in 8 (10) out of 57 cases (19 taxa x 3 phenological characteristics) at 5% (10%) significance level. Here, Poaceae and *Urtica* are the most important with notable changes in at least two characteristics. Our conclusions are in good agreement with those of several previous researches. For Thessaloniki (Greece), the total annual pollen counts, as well as the daily peak pollen counts show significant increasing trends for the majority of taxa, but there are no important changes for the phenological characteristics (Damialis et al. 2007). Looking at a bigger region of Central Europe, for Zurich, Switzerland (Frei 2008, *Betula*), as well as for Vienna, Austria (Jäger et al. 1996, *Alnus*, *Corylus*, *Betula*, *Pinus* and *Ulmus*) the pollen concentrations for most of the pollen types have been increasing. Furthermore, for Zurich (Frei 2008, *Betula*), Poznań, Poland (Stach et al. 2007, *Artemisia*) and Vienna (Jäger et al. 1996, *Alnus*, *Corylus*, *Betula*, *Pinus* and *Ulmus*) the pollen season starts earlier, the daily maximum pollen concentration has increased (Frei 2008, *Betula*) and the days of peak pollen counts occur earlier (Stach et al. 2007, *Artemisia*).

Note that all taxa examined in the study are families or genera involving a number of species. Accordingly, analysing pollen season and phenological characteristics of a family or genus instead of given species involves a high variability of pollen season data. An observed trend in the above characteristics incorporates the variability of a given parameter for all species belonging to a given taxon, but this variability is influenced by meteorological variables. The important role of sunshine hours is stressed here, since its high values enhance pollen production (Valencia-Barrera et al. 2001, Kasprzyk and Walanus 2010). We found increasing trends (at 5% significance level) in the total solar radiation, relative humidity and wind speed. Temperature and rainfall do not display overall significant trends, but the smoothing of daily MK test values shows stages of positive and negative trends within the year for these latter two variables as well (Fig. 1).

Based on an association measure (AM) – introduced to characterise the strength of the relationship between annual cycles of daily slopes of pollen concentration trends and

those of climate variables trends – the individual taxa were placed into three categories according to their climate sensitivity defined by a multiple AM (MAM). These are: (1) high sensitivity: $MAM > 0.950$, involving 11 taxa (*Artemisia*, *Cannabis*, *Alnus*, *Taxus*, *Rumex*, *Morus*, *Betula*, *Tilia*, *Chenopodiaceae*, *Pinus* and *Poaceae*); (2) medium sensitivity: $0.900 < MAM \leq 0.950$, including 5 taxa (*Platanus*, *Plantago*, *Ulmus*, *Juglans* and *Quercus*); (3) low sensitivity: $MAM \leq 0.900$, comprising 3 taxa (*Populus*, *Ambrosia* and *Urtica*) (Table 3).

Risk potential (RP) and expansion potential (EP) due to climate change are compared to the MAM for each taxon (Tables 3, 4). The association measure alone cannot contain or express the climate change related indicators. However, all taxa having the lowest climate sensitivity (+) are non-endangered (*) and, except for *Ambrosia*, are characterized by a moderate EP. At the same time, for all endangered taxa (***) (even if just one species is endangered within a given taxon) MAMs indicate high sensitivity (+++). Accordingly, the association measures follow well the climate change related indicators indicating that climate parameters are important elements of the environment for the taxa examined.

However, airborne pollen concentration can be influenced not only by current values of meteorological elements but their past values as well. As it is hard to distinguish between the effect of current and past values of the meteorological variables no attempt has been made to determine the relative weight of these two components in influencing the measured current pollen concentration. The procedure proposed here was performed only for *Ambrosia*, the most allergenic of all taxa considered. Current meteorological elements were characterized by actual values, while past meteorological elements by cumulative values of daily mean temperature, daily relative humidity, daily total solar radiation and daily precipitation total, respectively. These four elements were considered since ragweed pollen concentration displays a significant positive correlation with the daily mean temperature (Bartková-Ščevková 2003, Štefanič et al. 2005, Peternel et al. 2006, Puc 2006, Kasprzyk 2008), but it shows a negative correlation with the daily relative humidity (Bartková-Ščevková 2003, Puc 2006, Kasprzyk 2008). Daily mean total solar radiation is also found to be an important predictor of local ragweed pollen levels (Laaidi et al. 2003, Štefanič et al. 2005, Oh 2009). *Ambrosia* pollen grains are negatively correlated with rainfall (Barnes et al. 2001, Kasprzyk 2008), furthermore Déchamp and Penel (2001) found that heavy rainfall reduced the risk of ragweed pollen allergy. In order to assess the effect of the antecedent and current meteorological conditions on the current pollen concentration, the 1st-day, 2nd-day, ... , 93rd-day values of both the pollen concentration and the four meteorological elements of the current pollen season were taken. (The duration of the ragweed pollen season in Szeged lasts from July 15 until October 15, namely 93 days.) Association between pollen concentration and the four meteorological variables characterizes the role of current weather conditions. Values of these meteorological variables were then cumulated for 272-day, 271-day, ..., 1-day periods starting 272 days, 271 days, ..., 1 day before the actual day of the actual pollen season. This is because there are 272 days between the end of the previous-year pollination season and the beginning of the actual pollen season. Hence, altogether 272 data sets were produced and factor analysis with special transformation was performed for each of them.

The main conclusions for *Ambrosia* as an example are as follows (Fig. 2). The total weights (summarized absolute values) of the factor loadings for the past meteorological variables (dashed line) are gradually increasing from day 272 until day 123 reaching a

maximum value of 0.498. From this day until present the effect of the past climate parameters influencing daily *Ambrosia* pollen level decreases, though two local maxima (days 82 and 21) occur. The total weights of the factor loadings for the current climate parameters (solid line) are very low from the day 272 until the day 138. Then, they are increasing steeply reaching their top values between days 62 and 50. During the last 50 days until the start of the current *Ambrosia* pollen season the total weights are sharply decreasing. Note that these findings are valid only for variations of daily *Ambrosia* pollen concentrations accounted for by the above eight explanatory variables and nothing is known about the variance portion not explained by these variables. On the whole, in agreement with our preliminary expectations, past climate has a higher weight far from, while current climate has greater importance close to the time of the current pollen release. The effect of the past climate is greater on the current *Ambrosia* pollen concentration from day 272 until day 77, while from this day until present the current climate has higher weight in influencing the current *Ambrosia* pollen level. The effect of the current climate begins to increase on day 138 (i.e. 138 days before July 15, corresponding to the date February 28) (Fig. 2). This date corresponds well to the fact that the germination ability of over 80% for *Ambrosia* pollen seeds is mostly measured from the second half of February (Hartmann et al. 2003). Under the climate conditions of Hungary the germination peak of *Ambrosia* pollen seeds occurs in April (Béres et al. 2005) indicating a good agreement with the fact that the effect of the current climate becomes predominant over the past climate influencing *Ambrosia* pollen release from day 77 (i.e. 77 days preceding July 15, corresponding to the date April 30). Further climate-related significance of these days (February 28 and April 30) is as follows. By the end of February daily mean temperatures have already been in a positive range. The weight of the past climate elements culminates in January-February. This denotes the important effect of temperature and precipitation as practically the only parameters of the past climate to be considered for *Ambrosia* pollen concentration as resultant variable in winter. The effect of autumn rainfall lasts until the end of February. As soon as photosynthesis starts, autumn and winter precipitation is gradually utilised by the plants and hence leads to the decrease of the role of precipitation as an element of the past climate and to the increase of the role of current rainfall. Additionally, from the 2nd half of April mean and maximum temperatures suddenly increase that substantially contributes to the plants producing organic material for their generative processes.

Besides meteorological variables, pollen concentrations are influenced as well by agricultural and social factors (Makra et al. 2005) including urbanisation, so-called “green meadow investments” (new investments in former agricultural areas) and newly built motorways. Land eutrophication facilitating higher pollen production is not characteristic in an agricultural area consisting of small private plots for the Szeged agglomeration. A more important factor is that large industrial areas have come into use; housing estates as well as motorways were constructed in the region during the period investigated. Stripping agricultural lands for building purposes could mean an expansion of neglected areas that contributes to an increase of habitat regions of weeds and hence to an increase in pollen production. It would be important to distinguish between the changes in atmospheric pollen concentrations resulting from the effect of climate change and changes due to land use changes. The effect of climate can be characterised by meteorological variables, while land use changes can be described by changes in the ratios of agricultural areas, industrial areas, urban areas, forests, meadows, vineyards, orchards and fallows. Applying an appropriate statistical procedure (such as factor analysis with special transformation) the weight of both

climate-related and land use related components of atmospheric pollen concentrations could be estimated. However, information on the changes in land use is only available for years 1990, 2000 and 2006 in the CORINE Land Cover Database (www.eea.europa.eu) and hence such a statistical procedure cannot be performed.




Fig. 2 Total weights of the factor loadings for the current and past meteorological elements influencing current *Ambrosia* pollen concentration

Nevertheless, the role of land use changes in the pollen release of the taxa considered was also evaluated in the following simple way. CORINE Land Cover maps for the Szeged area are available with a 100 km radius around the centre of the city for the years 1990, 2000 and 2006. For the Szeged area, from short- to medium-range pollen transport involving local pollen dispersion has higher impact on daily *Ambrosia* pollen concentration compared to long-range pollen transport (Makra et al. 2010). Hence, an area of 100 km limit of medium-range transport (Makra et al. 2010) was considered. Then, in the centre of the radius a square was fit to the circle with its sides touching the circle that covers an area of 40,000 km². Changes in the individual land use cover categories were determined in m² from year 1990 to year 2000, as well as from year 2000 to year 2006. These changes were then expressed in the unit of the total area covered by the square. To sum up, changes in land use displayed $1.68 \cdot 10^{-5}$ % from year 1990 to year 2000, while $5.39 \cdot 10^{-3}$ % from year 2000 to year 2006. Therefore, land use changes did not influence pollen concentration of any taxa considered over the Szeged area in the period examined (www.eea.europa.eu/publications/COR0-landcover).

Acknowledgements: The authors would like to thank Gábor Motika (Environmental Conservancy Inspectorate, Szeged, Hungary) for providing meteorological data of Szeged and Miklós Juhász for providing pollen data of Szeged. The European Union and the European Social Fund provided financial support for the project under the grant agreement nos. TAMOP 4.2.1/B-09/1/KMR-2010-0003, TAMOP-4.2.1/B-09/1/KONV-2010-0005 and TAMOP-4.2.2/B-10/1-2010-0012).

REFERENCES

- Alcázar P, García-Mozo H, Trigo MD, Ruiz L, González-Minero FJ, Hidalgo P, de la Guardia CD, Galán C (2011) *J Environ Monitor* 13:2502-2510
- Ariano R, Canonica GW, Passalacqua G (2010) Possible role of climate changes in variations in pollen seasons and allergic sensitizations during 27 years. *Ann Allerg Asthma Im* 104:215-222
- Barnes C, Pacheco F, Landuyt J, Hu F, Portnoy J (2001) The effect of temperature, relative humidity and rainfall on airborne ragweed pollen concentrations. *Aerobiologia* 17:61-68
- Bartková-Ščeková J (2003) The influence of temperature, relative humidity and rainfall on the occurrence of pollen allergens (*Betula*, *Poaceae*, *Ambrosia artemisiifolia*) in the atmosphere of Bratislava (Slovakia). *Int J Biometeorol* 48:1-5
- Béres I, Novák R, Hoffmanné Pathy Z, Kazinczi G (2005) Az ürömlevelű parlagfű (*Ambrosia artemisiifolia* L.) elterjedése, morfológiája, biológiája, jelentősége és a védekezés lehetőségei. [Distribution, morphology, biology, importance and weed control of common ragweed (*Ambrosia artemisiifolia* L.) (in Hungarian)] *Gyomnövények, gyomirtás* 6:1-48
- Böloni J, Molnár Z, Kun A (eds) (2011) Magyarország élőhelyei. [Habitats of Hungary. (in Hungarian)] MTA Ökológiai és Botanikai Kutatóintézet, Vácrátót
- Borhidi A (1995) Social behaviour types, the naturalness and relative ecological indicator values of the higher plants in the Hungarian Flora. *Acta Bot Hung* 39:97-181
- Clot B (2003) Trends in airborne pollen: an overview of 21 years of data in Neuchâtel (Switzerland). *Aerobiologia* 19:227-234
- Cristofori A, Cristofolini F, Gottardini E (2010) Twenty years of aerobiological monitoring in Trentino (Italy): assessment and evaluation of airborne pollen variability. *Aerobiologia* 26:253-261
- Czúcz B (2009) Élővilág és éghajlatváltozás. Tudomány az élő természetért I. [Biosphere and climate change. Science for the biosphere 1. (in Hungarian)] MTA-ÖBKI, Vácrátót
- Damialis A, Halley JM, Gioulekas D, Vokou D (2007) Long-term trends in atmospheric pollen levels in the city of Thessaloniki, Greece. *Atmos Environ* 41:7011-7021
- Deák JÁ (2010) Csongrád megye kistájainak élőhelymintázata és tájökölógiai szempontú értékelése. [Habitat-pattern and landscape ecological evaluation of the micro-regions of Csongrád County. (in Hungarian)] PhD thesis. University of Szeged, Hungary
- Deák JÁ (2011a) Vegetáció-monitoring vizsgálatok Kunpeszérén a „Conservation of Hungarian meadow viper (*Vipera ursinii rakosiensis*) in the Carpathian-basin” című LIFE+ pályázathoz kapcsolódóan. Kutatási jelentés. [Vegetation monitoring researches in Kunpeszér, in the frame of LIFE+ programme titled “Conservation of Hungarian meadow viper (*Vipera ursinii rakosiensis*) in the Carpathian-basin”. Research Report. (in Hungarian)] Szeged
- Deák JÁ (2011b) A növényzet tájléptékű változásai a Kiskunság délkeleti részén. Környezeti változások és az Alföld. [Landscape-level changes of the vegetation of south-east Kiskunság. Environmental changes and the Great Hungarian Plain. (in Hungarian)] In: Rakonczai J (ed) *A Nagyalföld Alapítvány kötetei* 7:327-338
- Déchamp C, Penel V (2002) Results of pollen counts for 2001 from the Rhone-Alpes ragweed pollen-monitoring network (SARA). *Rev Fr Allergol* 42:539-542
- Emanuel WR, Shugart HH, Stevenson MP (1985) Climatic change and the broad-scale distribution of terrestrial ecosystem complexes. *Climatic Change* 7:29-43
- Fan J (1993) Local linear regression smoothers and their ninimax efficiency. *Annal Stat* 21:196-216
- Faragó T, Láng I, Cséte L (eds) (2010) Climate change and Hungary: mitigating the hazard and preparing for the impacts. The “VAHAVA” Report. Hungarian Academy of Sciences, Budapest. 124
- Feehan J, Harley M, van Minnen J (2009) Climate change in Europe. I. Impact on terrestrial ecosystems and biodiversity. A review (Reprinted). *Agron Sustain Dev* 29:409-421
- Francisco-Fernández M, Vilar-Fernández JM (2004) Weighted local nonparametric regression with dependent data: study of real private residential fixed investment in the USA. *Stat Infer Stoch Process* 7:69-93
- Frei T (2008) Climate change and its impact on airborne pollen in Basel, Switzerland 1969-2007. *Allergologie* 31:165-169
- Galán C, Cariñanos P, García-Mozo H, Alcázar P, Domínguez-Vilches E (2001) Model for forecasting *Olea europaea* L. airborne pollen in South-West Andalusia, Spain. *Int J Biometeorol* 45:59-63
- García-Mozo H, Mestre A, Galán C (2010) Phenological trends in southern Spain: A response to climate change. *Agr Forest Meteorol* 150:575-580

- Hartmann F, Hoffmann PZ, Tóth Csantavéri S (2003) A parlagfű (*Ambrosia artemisiifolia* L.) atrazinrezisztens biotípusának országos elterjedése. [Country-side distribution of an atrazine-resistant biotype of ragweed (*Ambrosia artemisiifolia* L.) (in Hungarian)] *Növényvédelem* 39:313-318
- Hirst JM (1952) An automatic volumetric spore trap. *Ann Appl Biol* 39:257-265
- Horváth F, Dobolyi ZK, Morschhauser T, Lökös L, Karas L, Szerdahelyi T (1995) Flóra adatbázis 1.2. [Flora Database 1.2. (in Hungarian)] MTA-ÖBKI, Vácrátót
- Jäger S, Nilsson S, Berggren B, Pessi AM, Helander M, Ramfjord H (1996) Trends of some airborne tree pollen in the Nordic countries and Austria, 1980-1993 – A comparison between Stockholm, Trondheim, Turku and Vienna. *Grana* 35:171-178
- Jahn W, Vahle H (1968) Die Faktoranalyse und ihre Anwendung. Verlag die Wirtschaft, Berlin
- Jolliffe IT (1993) Principal component analysis: A beginner's guide – II. Pitfalls, myths and extensions. *Weather* 48:246-253
- Kaminski U, Glod T (2011) Are there changes in Germany regarding the start of the pollen season, the season length and the pollen concentration of the most important allergenic pollens? *Meteorol Z* 20:497-507
- Kasprzyk I (2008) Non-native *Ambrosia* pollen in the atmosphere of Rzeszow (SE Poland): Evaluation of the effect of weather conditions on daily concentrations and starting dates of the pollen season, *Int J Biometeorol* 52:341-351
- Kasprzyk I, Walanus A (2010) Description of the main Poaceae pollen season using bi-Gaussian curves, and forecasting methods for the start and peak dates for this type of season in Rzeszow and Ostrowiec Sw. (SE Poland). *J Environ Monitor* 12:906-916
- Köppen W (1931) *Grundriss Der Klimakunde*. Walter De Gruyter & Co, Berlin
- Laaidi M, Thibaudon M, Besancenot JP (2003) Two statistical approaches to forecasting the start and duration of the pollen season of *Ambrosia* in the area of Lyon (France). *Int J Biometeorol* 48:65-73
- Láng I, Jolánkai M, Csete L (eds) (2007) Globális klímaváltozás: hazai hatások és válaszok – a VAHAVA jelentés. [Global climate change: effects and responses for Hungary – VAHAVA report. (in Hungarian)] Szaktudás Kiadó Ház, Budapest
- Leemans R, Cramer W, van Minnen JG (1996) Prediction of global biome distribution using bioclimatic equilibrium models. In: Melillo, JM, Breymer A (eds) *Effects of Global Change on Coniferous Forests and Grasslands*. Wiley, New York
- Lindner M, Maroschek M, Netherer S, Kremer A, Barbati A, Garcia-Gonzalo J, Seidl R, Delzon S, Corona P, Kolström M, Lexer MJ, Marchetti M (2010) Climate change impacts, adaptive capacity, and vulnerability of European forest ecosystems. *Forest Ecol Manag* 259:698-709
- Liu PWG (2009) Simulation of the daily average PM10 concentrations at Ta-Liao with Box-Jenkins time series models and multivariate analysis. *Atmos Environ* 43:2104-2113
- Makra L, Juhász M, Béczi R, Borsos E (2005) The history and impacts of airborne *Ambrosia* (Asteraceae) pollen in Hungary. *Grana* 44:57-64
- Makra L, Sánta T, Matyasovszky I, Damialis A, Karatzas K, Bergmann KC, Vokou D (2010) Airborne pollen in three European cities: Detection of atmospheric circulation pathways by applying three-dimensional clustering of backward trajectories. *J Geophys Res-Atmos* 115:D24220
- Makra L, Matyasovszky I, Páldy A, Deák JÁ (2012) The influence of extreme high and low temperatures and precipitation totals on pollen seasons of *Ambrosia*, *Poaceae* and *Populus* in Szeged, southern Hungary. *Grana* doi:10.1080/00173134.2012.661764
- Meyer CD (2001) *Matrix Analysis and Applied Linear Algebra*. Society for Industrial and Applied Mathematics, Philadelphia
- Oh JW (2009) Development of pollen concentration prediction models. *J Korean Med Assoc* 52:579-591
- Önöz B, Bayazit M (2003) The Power of Statistical Tests for Trend Detection. *Turk J Eng Environ Sci* 27:247-251
- Parry ML, Canziani OF, Palutikof JP, van der Linden PJ, Hanson CE (eds) (2007) IPCC. *Climate Change 2007. Impacts, Adaptation and Vulnerability*. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, UK
- Pearson RG (2006) Climate change and the migration capacity of species. *Trends Ecol Evol* 21:111-113
- Penuelas J, Boada M (2003) A global change-induced biome shift in the Montseny mountains (NE Spain). *Global Change Biol* 9:131-140
- Peternel R, Čulig J, Hrga I, Hercog P (2006) Airborne ragweed (*Ambrosia artemisiifolia* L.) pollen concentrations in Croatia, 2002-2004. *Aerobiologia* 22:161-168
- Puc M (2006) Ragweed and mugwort pollen in Szczecin, Poland. *Aerobiologia* 22:67-78
- Recio M, Docampo S, García-Sánchez J, Trigo MM, Melgar M, Cabezudo B (2010) Influence of temperature, rainfall and wind trends on grass pollination in Malaga (western Mediterranean coast). *Agr Forest Meteorol* 150:931-940

- Rodríguez-Rajo FJ, Aira MJ, Fernández-González M, Seijo C, Jato V (2011) Recent trends in airborne pollen for tree species in Galicia, NW Spain. *Climate Res* 48:281-291
- Soó R (1964-1980) A magyar flóra és vegetáció rendszertani-növényföldrajzi kézikönyve, I-VI. [Taxonomical and plant geographical handbook of the Hungarian flora and vegetation. (in Hungarian)] Akadémiai Kiadó, Budapest
- Stach A, García-Mozo H, Prieto-Baena JC, Czarnecka-Operacz M, Jenerowicz D, Silny W, Galán C (2007) Prevalence of *Artemisia* species pollinosis in western Poland: Impact of climate change on aerobiological trends, 1995-2004. *J Invest Allerg Clin* 17:39-47
- Štefanič E., Kovačević V, Lazanin Ž (2005) Airborne ragweed pollen concentration in north - eastern Croatia and its relationship with meteorological parameters. *Ann Agric Environ Med* 12:75-79
- Valencia-Barrera RM, Comtois P, Fernández-González D (2001) Biogeography and bioclimatology in pollen forecasting – An example of grass in Leon (Spain) and Montreal (Canada). *Grana* 40:223-229
- Zólyomi B, Précsényi I (1964) Methode zur ökologischen Charakterisierung der Vegetationseinheiten und zum Vergleich der Standorte. *Acta Bot Hung* 10:377-416
- www.eea.europa.eu/publications/COR0-landcover