

## SILICATE-CARBONATE-CO<sub>2</sub> IMMISCIBILITY IN ALKALINE ROCKS: EXAMPLE FROM INCLUSIONS IN PLAGIOCLASE OF TRACHYTIC LATITES AT CASERIO FUENTE DE MECA, MAZARRON, SE SPAIN

SHARYGIN, V. V. (Institute of Mineralogy and Petrography, Koptyuga prospect 3, 630090 Novosibirsk, Russia)  
sharygin@uiggm.nsc.ru

### Introduction

The Miocene magmatic province of SE Spain is characterized by calc-alkaline/shoshonitic and lamproitic rocks. In general, lamproites are mantle-derived rocks, whereas calc-alkaline and shoshonitic rocks have a more complicated origin involving both mantle and crust components (Venturelli et al., 1984; 1991). The origin of some rocks from the southern group of the Spanish lamproite occurrences (Zeneta, Vera) is suggested to be as a result of mixing between lamproitic and shoshonitic components (Fuster, 1956; Venturelli et al., 1984). To other side, the studies of trachytic latites from Caserio Fuente de Meca, Mazarron showed that these rocks possibly also derived due to mixing (Venturelli et al., 1991). This work is concerning to study of unusual silicate melt inclusions in the rocks of Caserio Fuente de Meca.

### Petrography of studied rocks

The trachytic latites contain phenocrysts of olivine altered to serpentine, resorbed phlogopite (TiO<sub>2</sub> – 5.2, Al<sub>2</sub>O<sub>3</sub> – 14.5 wt. %, Mg# - 83), biotite (TiO<sub>2</sub> – 3.6-4.2, Al<sub>2</sub>O<sub>3</sub> – 15.3-16.5 wt. %, Mg# - 52-57), plagioclase (An<sub>80-49</sub>Ab<sub>19-48</sub>Or<sub>1-3</sub>) and apatite (sizes – up to 0.5 mm). Groundmass consists of rhyolitic colorless glass, mica and apatite. In addition, large carbonate globules (up to 0.5 mm), xenocrysts of cordierite and lobated quartz occur in the groundmass. The minute crystals of zircon, graphite and spinel sometimes are observed both as single inclusions in phenocrysts and as accessory phases in the groundmass.

### Silicate melt inclusions

Silicate melt inclusions are found in all phenocrystal minerals, but they are more common for plagioclase. In this mineral the inclusions are mainly situated in the central parts or outline the growth zones. Their phase composition is diverse: some inclusions contain rhyolitic glass, fluid and daughter/trapped mica, but most inclusions consist of glass and heterogeneous fluid bubble (CO<sub>2</sub>vapor, CO<sub>2</sub>liquid, and carbonate) and rarely individual carbonate globule (Fig. 1, Table 1). It should be noted that individual carbonate inclusions coexisting with melt ones were not found. The heterogeneity in fluid bubble is expressed in variable amount of constituent phases. Majority of fluid bubbles contain mainly CO<sub>2</sub>vapor and CO<sub>2</sub>liquid, carbonate is minor. In inclusions, in which carbonate dominates; the vapor represents the CO<sub>2</sub> phase.




Figure 1. Silicate melt inclusions in plagioclase phenocrysts from trachytic latite, Caserio Fuente de Meca, SE Spain. Gl – rhyolitic glass; Phl – mica; Carb – carbonate; CO<sub>2</sub> v and CO<sub>2</sub> lq – carbon dioxide vapor and liquid, respectively.

**Table 1.** Chemistry of silicate melt inclusion (glass + carbonate + fluid) in plagioclase.

Phase	SiO <sub>2</sub>	TiO <sub>2</sub>	Al <sub>2</sub> O <sub>3</sub>	FeO	MnO	MgO	CaO	BaO	SrO	Na <sub>2</sub> O	K <sub>2</sub> O	P <sub>2</sub> O <sub>5</sub>	Cl	Sum
<b>Rhyolitic glass n=2</b>	72,37	0,07	13,24	1,34	0,09	0,19	0,85	0,04	0,00	2,08	4,34	0,11	0,38	95,09
<b>Carbonate globule</b>	0,00	0,00	0,00	1,01	0,81	1,67	55,49	0,00	0,12	0,00	0,07	0,00	0,00	59,18

### Discussion

Silicate melt inclusions in latites of Caserio Fuente de Meca demonstrate a unique case when CO<sub>2</sub> and carbonate may coexist in silicate glass. This case shows an evidence of silicate-carbonate-CO<sub>2</sub> immiscibility in siliceous volcanic rocks. It is unknown what is the main reason for this feature in the studied rocks. Possibly, it was contamination of carbonate rocks by siliceous magma within the upper-middle crust. The silicate-carbonate liquid immiscibility is more common for evolution of silica-undersaturated igneous rocks. For example, this phenomenon was observed in mineral-hosted melt inclusions of the Umbrian kamafugitic rocks (Italy), the Leucite Hill lamproites (USA) and mantle-derived xenoliths of the Canary Islands (Sharygin, 2001a, b; Frezzotti et al., 2002).

This work is supported by the Russian Foundation for Basic Research (grant no. 03-05-64030).

### Reference

- FREZZOTTI, M. L., ANDERSEN, T., NEUMANN, E.-R., SIMONSEN, S. L. (2002): Carbonatite melt-CO<sub>2</sub> fluid inclusions in mantle xenoliths from Tenerife, Canary Islands: a story of trapping, immiscibility and fluid-rock interaction in the upper mantle. *Lithos*, 64, 77-96.
- FUSTER, J. M. (1956): Las erupciones delleniticas del Terciario superior de la fosa de Vera (provincial de Almeria). *Bol. R. Soc. Esp. Hist. Nat. (Madrid)*, 54, 53-88.
- SHARYGIN, V.V. (2001a): Silicate-carbonate liquid immiscibility in melt inclusions from melilitolite minerals: the Pian di Celle volcano (Umbria, Italy). *Memórias*, no. 7, 399-402 (Abstracts of XVI ECROFI, Porto).
- SHARYGIN, V. V. (2001b): Lamproites: a review of magmatic inclusions in minerals. N.V. Vladykin (ed.): *Alkaline magmatism and the problems of mantle sources*. Irkutsk, 85-117.
- VENTURELLI, G., CAPEDE, S., DI BATTISTINI, G., CRAWFORD, A., KOGARKO, L. N., CELESTINI, S. (1984): The ultrapotassic rocks from southeastern Spain. *Lithos*, 17, 37-54.
- VENTURELLI, G., TOSCANI, L., SALVIOLI-MARIANI, E., CAPEDE, S. (1991): Mixing between lamproitic and dacitic components in Miocene volcanic rocks of S.E. Spain. *Mineralogical Magazine*, 55, 282-285.