



# Existence and stability of periodic solutions for a delayed prey–predator model with diffusion effects

Hongwei Liang, Jia-Fang Zhang<sup>✉</sup> and Zhiping Zhang

School of Mathematics and Statistics, Henan University, Kaifeng, 475001, China

Received 11 September 2015, appeared 8 January 2016

Communicated by Eduardo Liz

**Abstract.** Existence and stability of spatially periodic solutions for a delay prey–predator diffusion system are concerned in this work. We obtain that the system can generate the spatially nonhomogeneous periodic solutions when the diffusive rates are suitably small. This result demonstrates that the diffusion plays an important role in deriving the complex spatiotemporal dynamics. Meanwhile, the stability of the spatially periodic solutions is also studied. Finally, in order to verify our theoretical results, some numerical simulations are also included.

**Keywords:** existence, stability, spatially periodic solutions.

**2010 Mathematics Subject Classification:** 35K57, 35B10.

## 1 Introduction

In recent years, the interactions between two species have attracted much attention due to their theoretical and practical significance since the pioneering theoretical works by Lotka [22] and Volterra [28], see [4, 8, 10, 19, 30, 32]. It is well known that the interactions between two species have mainly three kinds of fundamental forms such as competition, cooperation and prey–predation in population biology. Among these interactions, extreme attention has been paid to the prey–predation mechanism because it possesses a very significant function as a kind of restriction factor in the process of evolvement of biology [6, 9, 17, 23, 25]. Understanding the dynamics of predator–prey models will be very helpful for investigating multiple species interactions. In [1], Beretta and Kuang have explored the dynamics of the following delayed Leslie–Gower model.

$$\begin{cases} \frac{du(t)}{dt} = r_1 u(t) \left[ 1 - \frac{u(t)}{K} \right] - mu(t)v(t), & t > 0, \\ \frac{dv(t)}{dt} = r_2 v(t) \left[ 1 - \frac{v(t-\tau)}{ru(t-\tau)} \right], & t > 0, \end{cases} \quad (1.1)$$

where  $u(t), v(t)$  are the population densities of the prey and the predator, respectively;  $r_1 > 0$ ,  $r_2 > 0$  denote the intrinsic growth rates of the prey and the predator, respectively.  $K > 0$  is the

<sup>✉</sup> Corresponding author. Email: jfzhang@henu.edu.cn

carrying capacity of the prey and  $ru$  takes on the role of a prey-dependent carrying capacity for the predator. The parameter  $r > 0$  is a measure of the quality of the prey as food for the predator. They presented some results on the boundedness of solutions permanence, global stability of the boundary equilibrium and local stability results of the positive equilibrium. Following this work, Song et al. [27] considered the properties of the local Hopf bifurcation and the global continuation of the local Hopf bifurcation for model (1.1).

In fact, the distribution of species is generally spatially inhomogeneous and therefore the species always tend to migrate toward regions of lower population density to improve the possibility of survival [29]. Therefore, spatial diffusion should be considered in modelling biological interactions, see [2, 3, 12, 13, 16, 20, 21, 26, 31]. Thus, the dynamics behavior of two species to model (1.1) should be described by the following model

$$\begin{cases} \frac{\partial u(t,x)}{\partial t} = d_1 \Delta u(t,x) + r_1 u(t,x) \left[1 - \frac{u(t,x)}{K}\right] - mu(t,x)v(t,x), & t > 0, x \in \Omega \\ \frac{\partial v(t,x)}{\partial t} = d_2 \Delta v(t,x) + r_2 v(t,x) \left[1 - \frac{v(t-\tau,x)}{ru(t-\tau,x)}\right], & t > 0, x \in \Omega, \\ u(t,x) = \phi(t,x) \geq 0, v(t,x) = \psi(t,x) \geq 0, & (t,x) \in [-\tau, 0] \times \Omega, \end{cases} \quad (1.2)$$

with Neumann boundary conditions

$$\frac{\partial u(t,x)}{\partial \nu} = \frac{\partial v(t,x)}{\partial \nu} = 0, \quad x \in \partial\Omega, t \geq 0. \quad (1.3)$$

$\Omega \subseteq \mathbb{R}^N$  is a bounded domain with smooth boundary  $\partial\Omega$ ;  $\nu$  is the unit outward normal vector on the boundary of  $\Omega$  and the Neumann boundary conditions in (1.3) imply that two species have zero flux across the domain boundary  $\partial\Omega$ ;  $d_1 > 0, d_2 > 0$  denote the diffusion coefficients of two species;  $(\phi, \psi) \in C = C([-\tau, 0], X)$ ,  $X$  is defined by

$$X = \left\{ (u, v) : u, v \in W^{2,2}(\Omega) : \frac{\partial u}{\partial \nu} = \frac{\partial v}{\partial \nu} = 0, x \in \partial\Omega \right\},$$

with the inner product  $\langle \cdot, \cdot \rangle$ .

The main goals of the present paper are to consider the existence and stability of spatially periodic solutions of system (1.2). By regarding the time delay  $\tau$  as the bifurcation parameter and analyzing the associated characteristic equation, we find that an increase of  $\tau$  can lead to the occurrence of spatially nonhomogeneous periodic solutions at  $(u^*, v^*)$ . Moreover, the stability of the spatially nonhomogeneous periodic solutions is studied.

The remaining parts of this paper are organized as follows. In Section 2, the existence of spatially nonhomogeneous periodic solutions is investigated. In Section 3, we derive conditions for determining the stability of the spatially nonhomogeneous periodic solutions on the center manifold. Finally, some conclusions and numerical simulations are presented in Section 4. Throughout the paper, we denote by  $\mathbb{N}$  the set of all positive integers, and  $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ .

## 2 Existence of spatially periodic solutions

In this section, we focus on investigating the local stability and the existence of spatially periodic solutions of the positive constant steady-state of system (1.2). It is easy to see that system (1.2) has two feasible boundary equilibria  $(0, 0)$ ,  $(K, 0)$  and a unique positive constant steady-state  $E^*(u^*, v^*)$ , where

$$u^* = \frac{1}{r}v^*, \quad v^* = \frac{Kr_1r}{r_1 + Kmr}.$$

Let  $\bar{u}(t, x) = u(t, x) - u^*$ ,  $\bar{v}(t, x) = v(t, x) - v^*$ , for convenience, we still use  $u$  and  $v$  to denote  $\bar{u}$  and  $\bar{v}$ . Then system (1.2) can be transformed into the following reaction-diffusion system when  $\Omega$  is restricted to the one-dimensional spatial domain  $(0, \pi)$ :

$$\begin{cases} \frac{\partial u(t, x)}{\partial t} = d_1 \frac{\partial^2 u(t, x)}{\partial x^2} + r_1(u(t, x) + u^*) \left[ 1 - \frac{u(t, x) + u^*}{K} \right] - m(u(t, x) + u^*)(v(t, x) + v^*), \\ \frac{\partial v(t, x)}{\partial t} = d_2 \frac{\partial^2 v(t, x)}{\partial x^2} + r_2(v(t, x) + v^*) \left[ 1 - \frac{v(t-\tau, x) + v^*}{r(u(t-\tau, x) + u^*)} \right], & t > 0, x \in \Omega, \\ \frac{\partial u(t, x)}{\partial x} = \frac{\partial v(t, x)}{\partial x} = 0, & t \geq 0, x \in \partial\Omega, \\ u(t, x) = \phi(t, x) - u^*, v(t, x) = \psi(t, x) - v^*, & (t, x) \in [-\tau, 0] \times \Omega. \end{cases} \quad (2.1)$$

Thus, the positive constant steady state  $E^*(u^*, v^*)$  of system (1.2) is transformed into the zero steady state of system (2.1).

By virtue of the Taylor expansions, system (2.1) can be rewritten as the following system

$$\begin{cases} \frac{\partial u(t, x)}{\partial t} = d_1 \Delta u(t, x) + \beta_{11}u(t, x) + \beta_{12}v(t, x) + \beta_{13}u^2(t, x) + \beta_{14}u(t, x)v(t, x), \\ \frac{\partial v(t, x)}{\partial t} = d_2 \Delta v(t, x) + \beta_{21}u(t - \tau, x) + \beta_{22}v(t - \tau, x) \\ \quad + \sum_{i+j+k \geq 2} \frac{1}{i!j!k!} f_{ijk} u^i(t - \tau, x) v^j(t - \tau, x) v^k(t, x), \end{cases} \quad (2.2)$$

where

$$\begin{aligned} \beta_{11} &= \frac{-r_1}{K} u^* < 0, & \beta_{12} &= -m u^* < 0, & \beta_{13} &= \frac{-r_1}{K} < 0, & \beta_{14} &= -m < 0, \\ \beta_{21} &= r r_2 > 0, & \beta_{22} &= -r_2 < 0 \\ f_{ijk} &= \frac{\partial^{i+j+k} f(0, 0)}{\partial u^i \partial v^j \partial v_1^k}, & f(u, v) &= r_2 v_1(t, x) \left( 1 - \frac{v(t, x)}{r u(t, x)} \right). \end{aligned}$$

Let  $u_1(t) = u(t, \cdot)$ ,  $u_2(t) = v(t, \cdot)$ , and  $U(t) = (u_1(t), u_2(t))^T$ . According to [11, 12], then system (2.2) can be rewritten as a delay differential equation in the phase space  $C = C([- \tau, 0], X)$

$$\frac{d}{dt} U(t) = d \Delta U(t) + L(U_t) + F(U_t), \quad (2.3)$$

where

$$d = \begin{pmatrix} d_1 & 0 \\ 0 & d_2 \end{pmatrix}, \quad \Delta = \begin{pmatrix} \frac{\partial^2}{\partial x^2} & 0 \\ 0 & \frac{\partial^2}{\partial x^2} \end{pmatrix},$$

$U_t(\theta) = U(t + \theta)$ ,  $-\tau \leq \theta \leq 0$ ,  $L : C \rightarrow X$  and  $F : C \rightarrow X$  are given, respectively, by

$$\begin{aligned} L(\varphi) &= \begin{pmatrix} \beta_{11} \varphi_1(0) + \beta_{12} \varphi_2(0) \\ \beta_{21} \varphi_1(-\tau) + \beta_{22} \varphi_2(-\tau) \end{pmatrix}, \\ F(\varphi) &= \begin{pmatrix} \beta_{13} \varphi_1^2(0) + \beta_{14} \varphi_1(0) \varphi_2(0) \\ \sum_{i+j+k \geq 2} \frac{1}{i!j!k!} f_{ijk} \varphi_1^i(-\tau) \varphi_2^j(-\tau) \varphi_2^k(0) \end{pmatrix}, \end{aligned}$$

where  $\varphi(\theta) = U_t(\theta)$ ,  $-\tau \leq \theta \leq 0$ ,  $\varphi = (\varphi_1, \varphi_2)^T \in C$ .

Linearizing (2.3) at  $(0, 0)$  gives the linear equation

$$\frac{d}{dt} U(t) = d \Delta U(t) + L(U_t). \quad (2.4)$$

The characteristic equation for the linearized equation (2.4) is

$$\lambda y - d\Delta y - L(e^{\lambda \cdot} y) = 0, \quad (2.5)$$

where  $y \in \text{dom}(\Delta) \setminus \{0\}$  and  $\text{dom}(\Delta) \subset X$ .

It is well known that the linear operator  $\Delta$  on  $(0, \pi)$  with homogeneous Neumann boundary conditions has the eigenvalues  $-k^2$  ( $k \in \mathbb{N}_0$ ) and the corresponding eigenfunctions are

$$\beta_k^1 = \begin{pmatrix} \gamma_k \\ 0 \end{pmatrix}, \quad \beta_k^2 = \begin{pmatrix} 0 \\ \gamma_k \end{pmatrix}, \quad \gamma_k = \frac{\cos(kx)}{\|\cos(kx)\|_{2,2}}, \quad k \in \mathbb{N}_0.$$

Notice that  $(\beta_k^1, \beta_k^2)_{k=0}^\infty$  construct an orthogonal basis of the Banach space  $X$  (see [12]). Therefore  $L(\beta_k^1, \beta_k^2) \subset \text{span}\{\beta_k^1, \beta_k^2\}$  and thus any element  $y$  in  $X$  can be expanded a Fourier series in the form

$$\begin{aligned} y &= \sum_{k=0}^{\infty} \left( \langle y, \beta_k^1 \rangle \beta_k^1 + \langle y, \beta_k^2 \rangle \beta_k^2 \right) \\ &= \sum_{k=0}^{\infty} \left( \langle y, \beta_k^1 \rangle, \langle y, \beta_k^2 \rangle \right) \begin{pmatrix} \beta_k^1 \\ \beta_k^2 \end{pmatrix}. \end{aligned} \quad (2.6)$$

In addition, some easy computations can show that

$$L \left( \varphi^T \begin{pmatrix} \beta_k^1 \\ \beta_k^2 \end{pmatrix} \right) = [L(\varphi)]^T \begin{pmatrix} \beta_k^1 \\ \beta_k^2 \end{pmatrix}, \quad (2.7)$$

where  $\varphi = (\varphi_1, \varphi_2)^T \in \mathbb{C}$ .

From (2.6) and (2.7), (2.5) is equivalent to

$$\sum_{k=0}^{\infty} \left( \langle y, \beta_k^1 \rangle, \langle y, \beta_k^2 \rangle \right) \left[ \begin{pmatrix} \lambda + d_1 k^2 & 0 \\ 0 & \lambda + d_2 k^2 \end{pmatrix} - \begin{pmatrix} \beta_{11} & \beta_{12} \\ \beta_{21} e^{-\lambda \tau} & \beta_{22} e^{-\lambda \tau} \end{pmatrix} \right] \begin{pmatrix} \beta_k^1 \\ \beta_k^2 \end{pmatrix} = 0.$$

Hence, we conclude that the characteristic equation (2.4) is equivalent to the sequence of the characteristic equations

$$\begin{aligned} \lambda^2 + [(d_1 + d_2)k^2 - \beta_{11}] \lambda + [d_1 d_2 k^4 - d_2 \beta_{11} k^2] \\ + [-\beta_{22} \lambda - d_1 \beta_{22} k^2 + \beta_{11} \beta_{22} - \beta_{12} \beta_{21}] e^{-\lambda \tau} = 0, \quad k \in \mathbb{N}_0. \end{aligned} \quad (2.8)$$

It is obvious that equation (2.8) has no zero roots since  $\beta_{11} < 0$ ,  $\beta_{12} < 0$ ,  $\beta_{21} > 0$ ,  $\beta_{22} < 0$ ,  $d_1 > 0$ ,  $d_2 > 0$ .

When  $\tau = 0$ , (2.8) reduces to the following quadratic equation with respect to  $\lambda$

$$\begin{aligned} \lambda^2 + [(d_1 + d_2)k^2 - \beta_{11} - \beta_{22}] \lambda \\ + [d_1 d_2 k^4 - d_2 \beta_{11} k^2 - d_1 \beta_{22} k^2 + \beta_{11} \beta_{22} - \beta_{12} \beta_{21}] = 0, \quad k \in \mathbb{N}_0, \end{aligned} \quad (2.9)$$

where

$$\begin{aligned} [(d_1 + d_2)k^2 - \beta_{11} - \beta_{22}] &> 0, \\ [d_1 d_2 k^4 - d_2 \beta_{11} k^2 - d_1 \beta_{22} k^2 + \beta_{11} \beta_{22} - \beta_{12} \beta_{21}] &> 0. \end{aligned}$$

Consequently, all roots of equations (2.9) have negative real parts. Therefore, the positive steady state  $E(u^*, v^*)$  of system (1.2) is locally asymptotically stable in the absence of delay.

When  $d_1 = d_2 = 0$  and  $\tau = 0$ , system (1.2) becomes an ordinary differential equation, we know that roots of the characteristic equation of ordinary differential equations have negative real parts. This indicates that the diffusion coefficients  $d_1, d_2$  have no effect on the stability of the positive steady state  $E(u^*, v^*)$  in the absence of delay.

Denote

$$(H) \quad d_1\beta_{22} - d_2\beta_{11} > 0, \text{ and } d_1d_2 + (d_1\beta_{22} - d_2\beta_{11}) + (\beta_{12}\beta_{21} - \beta_{11}\beta_{22}) > 0.$$

**Lemma 2.1.** *Assume that the condition (H) holds. If*

$$(d_1^2 + d_2^2) - 2\beta_{11}d_1 < \beta_{22}^2 - \beta_{11}^2 < 16(d_1^2 + d_2^2) - 8\beta_{11}d_1, \quad (2.10)$$

$$[A_1^2 - 2B_1^2 - \beta_{22}^2]^2 - 4(B_1^2 - C_1^2) > 0, \quad (2.11)$$

then (2.8) with  $k = 1$  has purely imaginary roots  $\pm i\omega_1$ , where

$$\omega_1 = \sqrt{\frac{-(A_1^2 - 2B_1 - \beta_{22}^2) \pm \sqrt{(A_1^2 - 2B_1 - \beta_{22}^2)^2 - 4(B_1^2 - C_1^2)}}{2}}.$$

*Proof.* Assuming  $i\omega$  ( $\omega > 0$ ) is a solution of (2.8) with  $k \geq 1$ , then substituting  $i\omega$  into equation (2.8) and separating the real and imaginary parts, one can get that

$$-\omega^2 + B_k - \beta_{22}\omega \sin(\omega\tau) + C_k \cos(\omega\tau) = 0, \quad (2.12)$$

$$A_k\omega - \beta_{22}\omega \cos(\omega\tau) - C_k \sin(\omega\tau) = 0, \quad (2.13)$$

where

$$A_k = (d_1 + d_2)k^2 - \beta_{11} > 0, \quad B_k = d_1d_2k^4 - d_2\beta_{11}k^2 \geq 0,$$

$$C_k = -d_1\beta_{22}k^2 + \beta_{11}\beta_{22} - \beta_{12}\beta_{21} > 0, \quad k \in \mathbb{N}_0.$$

From (2.12) and (2.13), it is easy to see that

$$\omega^4 + (A_k^2 - 2B_k - \beta_{22}^2)\omega^2 + B_k^2 - C_k^2 = 0, \quad k \in \mathbb{N}_0. \quad (2.14)$$

By computing, we have  $B_k - C_k = d_1d_2k^4 + (d_1\beta_{22} - d_2\beta_{11})k^2 + \beta_{12}\beta_{21} - \beta_{11}\beta_{22}$ . It is clear that  $d_1d_2k^4 + (d_1\beta_{22} - d_2\beta_{11})k^2 + \beta_{12}\beta_{21} - \beta_{11}\beta_{22} \geq d_1d_2 + (d_1\beta_{22} - d_2\beta_{11}) + \beta_{12}\beta_{21} - \beta_{11}\beta_{22}$  when  $d_1\beta_{22} - d_2\beta_{11} > 0$  ( $k \geq 1$ ). According to  $B_k \geq 0, C_k > 0$ , if the condition (H) holds, we can get  $B_k^2 > C_k^2$  when  $k \geq 1$ . Obviously,  $A_k^2 - 2B_k - \beta_{22}^2 = (d_1^2 + d_2^2)k^4 - 2d_1\beta_{11}k^2 + \beta_{11}^2 - \beta_{22}^2$ ; if  $16(d_1^2 + d_2^2) - 8d_1\beta_{11} + \beta_{11}^2 - \beta_{22}^2 > 0$ , that is,  $\beta_{22}^2 - \beta_{11}^2 < 16(d_1^2 + d_2^2) - 8d_1\beta_{11}$ , then (2.14) with  $k \geq 2$  has no positive roots.

Clearly, if  $d_1^2 + d_2^2 - 2d_1\beta_{11} + \beta_{11}^2 - \beta_{22}^2 < 0$ , that is,  $\beta_{22}^2 - \beta_{11}^2 > d_1^2 + d_2^2 - 2d_1\beta_{11}$ , and  $[A_1^2 - 2B_1 - \beta_{22}^2]^2 - 4(B_1^2 - C_1^2) \geq 0$ , then (2.14) with  $k = 1$  has at least one positive root  $\omega_1$ . From (2.10), (2.11), (H) and (2.14), we have

$$\omega_1 = \sqrt{\frac{-(A_1^2 - 2B_1 - \beta_{22}^2) \pm \sqrt{(A_1^2 - 2B_1 - \beta_{22}^2)^2 - 4(B_1^2 - C_1^2)}}{2}}. \quad (2.15)$$

That is, it has  $\omega_1$  such that (2.8) with  $k = 1$  has purely imaginary eigenvalues  $\pm i\omega_1$ . Thus the proof is complete.  $\square$

According to (2.12),(2.13) and Lemma 2.1, we get

$$\tau_j = \frac{1}{\omega_1} \left( \arccos \frac{(A_1\beta_{22} + C_1)\omega_1^2 - B_1C_1}{C_1^2 + \beta_{22}^2\omega_1^2} + 2j\pi \right), \quad j = 0, 1, \dots \quad (2.16)$$

**Lemma 2.2.** *Let  $\lambda(\tau) = \mu(\tau) \pm i\omega(\tau)$  be the root of (2.8) with  $k = 1$  near  $\tau = \tau_j$  satisfying  $\mu(\tau_j) = 0$ ,  $\omega(\tau_j) = \omega_1$ ,  $j = 0, 1, \dots$ . Then, the following transversality condition holds*

$$\text{sign Re} \left[ \frac{d\lambda}{d\tau} \right]_{\tau=\tau_j} \neq 0.$$

*Proof.* Taking the derivative for equation (2.8) with respect to  $\tau$  at  $\tau_j$ , we have

$$\left[ \frac{d\lambda}{d\tau} \right]^{-1} = \frac{(2\lambda + A_1)e^{\lambda\tau} - \beta_{11}}{(C_1 - \beta_{11}\lambda)\lambda} - \frac{\tau}{\lambda}. \quad (2.17)$$

From (2.17), we get

$$\begin{aligned} \text{sign Re} \left[ \frac{d\lambda}{d\tau} \right]_{\tau=\tau_j}^{-1} &= \text{sign Re} \left[ \frac{(2\lambda + A_1)e^{\lambda\tau} - \beta_{22}}{(C_1 - \beta_{22}\lambda)\lambda} - \frac{\tau}{\lambda} \right]_{\tau=\tau_j} \\ &= \frac{(A_1 \cos \omega_1 \tau_j - 2\omega_1 \sin \omega_1 \tau_j - \beta_{22})\beta_{11}\omega_1^2}{(\beta_{22}^2\omega_1^2 + C_1^2)\omega_1^2} \\ &\quad + \frac{(A_1 \sin \omega_1 \tau_j + 2\omega_1 \cos \omega_1 \tau_j)C_1\omega_1}{(\beta_{22}^2\omega_1^2 + C_1^2)\omega_1^2} \\ &= \frac{1}{\beta_{22}^2\omega_1^2 + C_1^2} [A_1^2 - 2B_1 - \beta_{22}^2 + 2\omega_1^2] \\ &= \frac{1}{\beta_{22}^2\omega_1^2 + C_1^2} \left[ \sqrt{(A_1^2 - 2B_1 - \beta_{22}^2)^2 - 4[B_1^2 - C_1^2]} \right] \neq 0, \end{aligned}$$

then

$$\text{sign Re} \left[ \frac{d\lambda}{d\tau} \right]_{\tau=\tau_j} \neq 0.$$

Thus the proof is complete.  $\square$

Therefore, we have the following conclusions,

**Theorem 2.3.** *Suppose that the conditions in Lemma 2.1 are satisfied. Let  $\tau_j$  be defined as in (2.16).*

- (i) *If  $\tau \in [0, \tau_0)$ , then the positive constant steady-state solution  $E^* = (u^*, v^*)$  of system (1.2) is stable and unstable when  $\tau > \tau_0$ .*
- (ii) *System (1.2) can have spatially nonhomogeneous periodic solutions at the positive constant steady-state solution  $E^* = (u^*, v^*)$  when  $\tau = \tau_j$ .*

### 3 Stability of spatially periodic solutions

In the previous section, we have obtained the existence of spatially periodic solutions of system (1.2) when the parameter  $\tau$  crosses through the critical value  $\tau_j$  ( $j = 0, 1, 2, \dots$ ). In this

section, we shall study the stability of periodic solutions by applying the normal form theory of partial functional differential equations developed by [15,29].

Normalizing the delay  $\tau$  in system (2.2) by the time-scaling  $t \rightarrow \frac{t}{\tau}$ , (2.2) is transformed into

$$\begin{cases} \frac{\partial u(t,x)}{\partial t} = \tau \{d_1 \Delta u + \beta_{11}u(t) + \beta_{12}v(t) + \beta_{13}u^2(t) + \beta_{14}u(t)v(t), \\ \frac{\partial v(t,x)}{\partial t} = \tau \left\{ d_2 \Delta v + \beta_{21}u(t-1) + \beta_{22}v(t-1) + \sum_{i+j+k \geq 2} \frac{1}{i!j!k!} f_{ijk} u^i(t-1)v^j(t-1)v^k \right\}, \end{cases} \quad (3.1)$$

where  $f_{ijk}$  is defined by (2.2). Let  $\tau = \tau_j + \alpha$ , then, (3.1) can be written in abstract form in  $\mathcal{C} = C([-1,0] : X)$  as

$$\frac{d}{dt}U(t) = (\tau_j + \alpha)d\Delta U(t) + L(\tau_j)(U_t) + F(U_t, \alpha), \quad (3.2)$$

where  $d = \begin{pmatrix} d_1 & 0 \\ 0 & d_2 \end{pmatrix}$ ,  $L(\alpha)(\cdot) : \mathcal{C} \rightarrow X$ ,  $F(\cdot, \alpha) : \mathcal{C} \rightarrow X$  are given by

$$\begin{aligned} L(\alpha)(\varphi) &= (\tau_j + \alpha) \begin{pmatrix} \beta_{11}\varphi_1(0) + \beta_{12}\varphi_2(0) \\ \beta_{21}\varphi_1(-1) + \beta_{22}\varphi_2(-1) \end{pmatrix}, \\ F(\varphi, \alpha) &= \alpha \Delta \varphi(0) + L(\alpha)\varphi + f(\varphi, \alpha), \end{aligned}$$

and

$$f(\varphi, \alpha) = (\tau_j + \alpha) \begin{pmatrix} \beta_{13}\varphi_1^2(0) + \beta_{14}\varphi_1(0)\varphi_2(0), \\ \sum_{i+j+k \geq 2} \frac{1}{i!j!k!} f_{ijk} \varphi_1^i(-1)\varphi_2^j(-1)\varphi_2^k(0) \end{pmatrix},$$

for  $\varphi = (\varphi_1, \varphi_2)^T \in \mathcal{C}$ .

Linearizing (3.2) at  $(0,0)$  leads to the following linear equation

$$\frac{d}{dt}U(t) = \tau_j d\Delta U(t) + L(\tau_j)(U_t). \quad (3.3)$$

It is easy to see from the discussions in the previous section that (2.8) has two purely imaginary eigenvalues  $\pm i\omega_1$  ( $\omega_1$  is defined by (2.15)).

Let  $\Lambda_1 = \{-i\omega_1, i\omega_1\}$ , consider the following FDE on  $C([-1,0], \mathbb{R}^2)$

$$\dot{z}(t) = L(\tau_j)(z_t), \quad (3.4)$$

that is,

$$\begin{pmatrix} \dot{z}_1(t) \\ \dot{z}_2(t) \end{pmatrix} = (\tau_j + \alpha) \left\{ \begin{pmatrix} \beta_{11} & \beta_{12} \\ 0 & 0 \end{pmatrix} \begin{pmatrix} z_1(t) \\ z_2(t) \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ \beta_{21} & \beta_{22} \end{pmatrix} \begin{pmatrix} z_1(t-1) \\ z_2(t-1) \end{pmatrix} \right\}.$$

As is well known,  $L(\tau_j)$  is a continuous linear function mapping  $C([-1,0], \mathbb{R}^2)$  into  $\mathbb{R}^2$ . According to the Riesz representation theorem, there exists a  $2 \times 2$  matrix function  $\eta(\theta, \tau)$  ( $-1 \leq \theta \leq 0$ ), whose elements are of bounded variation such that

$$L(\tau_j)(\phi) = \int_{-1}^0 d\eta(\theta, \tau_j) \phi(\theta) \quad \text{for } \phi \in \mathcal{C}. \quad (3.5)$$

Thus, we can choose

$$\eta(\theta, \tau_j) = (\tau_j + \alpha) \begin{pmatrix} \beta_{11} & \beta_{12} \\ 0 & 0 \end{pmatrix} \delta(\theta) - (\tau_j + \alpha) \begin{pmatrix} 0 & 0 \\ \beta_{21} & \beta_{22} \end{pmatrix} \delta(\theta + 1), \quad (3.6)$$

where  $\delta(0) = 1$ ,  $\delta(\theta) = 0$ ,  $-1 \leq \theta < 0$ , then (3.5) is satisfied.

If  $\phi$  is any given function in  $C([-1, 0], \mathbb{R}^2)$  and  $u(\phi)$  is the unique solution of the linear equation (3.3) with the initial function  $\phi$  at zero, then the solution operator

$$T(t) : C([-1, 0], \mathbb{R}^2) \rightarrow C([-1, 0], \mathbb{R}^2)$$

is defined by

$$T(t)\phi = u_t(\phi), \quad t \geq 0.$$

Let  $A(\tau_j)$  denote the infinitesimal generator of the strongly continuous semigroup, according to [14], then,

$$A(\tau_j)\phi(\theta) = \begin{cases} d\phi(\theta)/d\theta, & \theta \in [-1, 0), \\ L(\tau_j)(\phi) \stackrel{\text{def}}{=} \int_{-1}^0 d\eta(t, \tau_j)\phi(t), & \theta = 0, \end{cases} \quad (3.7)$$

where  $\phi \in C^1([-1, 0], \mathbb{R}^2)$ .

For  $\psi \in C^1([0, 1], (\mathbb{R}^2)^*)$ , define

$$A^*\psi(s) = \begin{cases} -d\psi(s)/ds, & s \in (0, 1], \\ \int_{-1}^0 \psi(-t)d\eta(t, \tau_j), & s = 0, \end{cases} \quad (3.8)$$

and a bilinear inner product of the Sobolev space  $W^{2,2}(0, \pi)$ .

$$\begin{aligned} (\psi(s), \phi(\theta)) &= \psi(0)\phi(0) - \int_{-1}^0 \int_{\xi=0}^{\theta} \psi(\xi - \theta)d\eta(\theta)d\xi \\ &= \psi(0)\phi(0) - \tau_j \int_{-1}^0 \psi(s+1) \begin{pmatrix} 0 & 0 \\ \beta_{21} & \beta_{22} \end{pmatrix} \phi(s) ds, \end{aligned}$$

where  $\eta(\theta) = \eta(\theta, \tau_j)$  and  $A^*$  is the formal adjoint of  $A(\tau_j)$ .

Obviously, the characteristic equation of the linear operator  $A(\tau_j)$  is (2.8) with  $k = 1$ . So, it is easy to see from Section 2 that  $A(\tau_j)$  has a pair of simple purely imaginary eigenvalues  $\pm i\omega_1$  and they are also eigenvalues of  $A^*$  since  $A(\tau_j)$  and  $A^*$  are adjoint operators. Let  $\mathcal{P}$  and  $\mathcal{P}^*$  be the center spaces, that is, the generalized eigenspaces, of  $A(\tau_j)$  and  $A^*$  associated with  $\Lambda_1$ , respectively, then  $\mathcal{P}^*$  is the adjoint space of  $\mathcal{P}$  and  $\dim \mathcal{P} = \dim \mathcal{P}^* = 2$ .

In addition, according to [11, 27], a few simple calculations, we can choose  $\Phi$  and  $\Psi$  be the bases for  $\mathcal{P}$  and  $\mathcal{P}^*$ , respectively. It is known that  $\dot{\Phi} = \Phi B$ , where  $B$  is the  $2 \times 2$  diagonal matrix  $B = \begin{pmatrix} i\omega_1 & 0 \\ 0 & i\omega_1 \end{pmatrix}$ .

Let  $\Phi = (\Phi_1, \Phi_2)$  and  $\Psi = (\Psi_1, \Psi_2)^T$ , where

$$\begin{aligned} \Phi_1(\theta) &= e^{i\omega_1\theta} \left( 1, \frac{i\omega_1 + d_1 - \beta_{11}}{\beta_{12}} \right)^T, & \Phi_2(\theta) &= \overline{\Phi_1(\theta)}, & -1 \leq \theta \leq 0, \\ \Psi_1(s) &= \frac{1}{\rho} \left( 1, -\frac{i\omega_1 - d_1 + \beta_{11}}{\beta_{21}e^{i\omega_1\tau_j}} \right)^T e^{-i\omega_1 s}, & \Psi_2(s) &= \overline{\Psi_1(s)}, & 0 \leq s \leq 1, \end{aligned}$$

$$\bar{\rho} = \frac{1}{1 + \rho v - \tau_j(-d_1 + \beta_{11} + \beta_{21}\rho + \beta_{12}v + (\beta_{22} - d_2)\rho v'}$$



where  $q = \frac{i\omega_1 + d_1 - \beta_{11}}{\beta_{12}}$ ,  $v = -\frac{i\omega_1 - d_1 + \beta_{11}}{\beta_{21}e^{i\omega_1\tau_j}}$ . Let  $f_1 = (\beta_1^1, \beta_1^2)$ ,  $c \cdot f_1$  be defined by  $c \cdot f_1 = c_1\beta_1^1 + c_2\beta_1^2$  for  $c = (c_1, c_2)^T \in \mathbb{R}^2$  and  $(\psi \cdot f_1)(\theta) = \psi(\theta) \cdot f_1$  for  $\theta \in [-1, 0]$ . Then the center space of linear equation (3.3) is given by  $\mathcal{P}_{CN}\mathcal{C}$ , where

$$P_{CN}\varphi = \Phi(\Psi, \langle \varphi, f_1 \rangle) \cdot f_1, \quad \varphi \in \mathcal{C}, \quad (3.9)$$

and  $\mathcal{C} = \mathcal{P}_{CN}\mathcal{C} \oplus \mathcal{P}_Q\mathcal{C}$ , here  $\mathcal{P}_Q\mathcal{C}$  denotes the complementary subspace of  $\mathcal{P}_{CN}\mathcal{C}$  in  $\mathcal{C}$ .

Let  $A_{\tau_j}$  be defined by:

$$A_{\tau_j}\varphi(\theta) = \dot{\varphi}(\theta) + X_1(\theta) [\tau_j\Delta\varphi(0) + L_*(\tau_j)(\varphi(\theta)) - \dot{\varphi}(0)], \quad \varphi \in \mathcal{C},$$

where  $X_1: [-1, 0] \rightarrow B(X, X)$  is given by

$$X_1 = \begin{cases} 0, & \theta \in [-1, 0), \\ I, & \theta = 0. \end{cases}$$

Then the infinitesimal generator  $A_{\tau_j}$  induced by the solution of (3.3) and (3.2) can be rewritten as the following operator differential equation

$$\dot{U}_t = A_{\tau_j}U_t + X_1F(U_t, \alpha). \quad (3.10)$$

Using the decomposition  $\mathcal{C} = \mathcal{P}_{CN} \oplus \mathcal{P}_Q\mathcal{C}$  and (3.9), the solution of (3.1) can be written as

$$U_t = \Phi \begin{pmatrix} x_1(t) \\ x_2(t) \end{pmatrix} \cdot f_1 + h(x_1, x_2, \alpha),$$

where  $(x_1, x_2)^T = (\Psi, \langle U_t, f_1 \rangle)$ , and  $h(x_1, x_2, \alpha) \in \mathcal{P}_Q\mathcal{C}$  with  $h(0, 0, 0) = Dh(0, 0, 0) = 0$ .

Thus, the flow on the center manifold for (3.2) can be described as

$$\begin{pmatrix} \dot{x}_1(t) \\ \dot{x}_2(t) \end{pmatrix} = \begin{pmatrix} 0 & \omega_1 \\ -\omega_1 & 0 \end{pmatrix} \begin{pmatrix} x_1(t) \\ x_2(t) \end{pmatrix} + \Psi(0)F(0, x_1(t), x_2(t)),$$

where

$$F(0, x_1(t), x_2(t)) = \left\langle f \left( \alpha, \Phi \begin{pmatrix} x_1(t) \\ x_2(t) \end{pmatrix} \cdot f_1 + h(\alpha, x_1(t), x_2(t)) \right), f_1 \right\rangle.$$

Let  $z = x_1 - ix_2$ , and  $\Psi(0) = (\Psi_1(0), \Psi_2(0))^T$ , when  $\alpha = 0$ , then  $z$  satisfies

$$\dot{z} = i\omega_1 z + g(z, \bar{z}), \quad (3.11)$$

where

$$g(z, \bar{z}) = (\Psi_1(0) - i\Psi_2(0)) \left\langle f \left( 0, \frac{1}{2} \Phi \begin{pmatrix} z + \bar{z} \\ (z - \bar{z})i \end{pmatrix} \cdot f_1 + w(z, \bar{z}) \right), f_1 \right\rangle,$$

$$w(z, \bar{z}) = h \left( 0, \frac{z + \bar{z}}{2}, \frac{(z - \bar{z})i}{2} \right), \quad (3.12)$$

$$w(z, \bar{z}) = w_{20} \frac{z^2}{2} + w_{11} z\bar{z} + w_{02} \frac{\bar{z}^2}{2} + w_{21} \frac{z^2\bar{z}}{2} + \dots \quad (3.13)$$

Noticing that  $p_1 = \Phi_1 + i\Phi_2$ ,  $p_2 = \bar{p}_1$ , therefore, solutions of (3.10) can be rewritten as

$$U_t = \frac{1}{2} \Phi \begin{pmatrix} z + \bar{z} \\ i(z - \bar{z}) \end{pmatrix} \cdot f_1 + w(z, \bar{z}) = \frac{1}{2} (p_1 z + p_2 \bar{z}) \cdot f_1 + w(z, \bar{z}). \quad (3.14)$$

In addition, (3.11) can be rewritten as the following form

$$\dot{z} = i\omega_1 z + g_{20} \frac{z^2}{2} + g_{11} z\bar{z} + g_{02} \frac{\bar{z}^2}{2} + g_{21} \frac{z^2\bar{z}}{2} + \cdots. \quad (3.15)$$

Let

$$g(z, \bar{z}) = g_{20} \frac{z^2}{2} + g_{11} z\bar{z} + g_{02} \frac{\bar{z}^2}{2} + g_{21} \frac{z^2\bar{z}}{2} + \cdots.$$

From (3.12), we have

$$\begin{aligned} & \langle F(U_t, 0), f_1 \rangle \\ &= \frac{\tau_j}{4} \left( \begin{array}{c} (\varrho\beta_{14} + \frac{1}{2}\beta_{13}) z^2 \\ e^{-2i\omega_1\tau_j}(\varrho f_{110} + e^{i\omega_1\tau_j}\varrho f_{101} + e^{i\omega_1\tau_j}\varrho^2 f_{011}^{(2)} + \frac{1}{2}f_{200} + \frac{1}{2}\varrho^2 f_{020})z^2 \end{array} \right) \\ &+ \frac{\tau_j}{4} \left( \begin{array}{c} [(\bar{\varrho} + \varrho)\beta_{14} + \beta_{13}] z\bar{z} \\ [(\bar{\varrho} + \varrho)f_{110} + e^{-i\omega_1\tau_j}\bar{\varrho}(f_{101} + \varrho f_{011}) + e^{i\omega_1\tau_j}\varrho(f_{101} + \bar{\varrho}f_{011}) + f_{200} + \varrho\bar{\varrho}f_{020}] z\bar{z} \end{array} \right) \\ &+ \frac{\tau_j}{4} \left( \begin{array}{c} (\bar{\varrho}\beta_{14} + \beta_{13}) \bar{z}^2 \\ e^{2i\omega_1\tau_j}(\bar{\varrho}f_{110} + e^{-i\omega_1\tau_j}\bar{\varrho}f_{101} + e^{-i\omega_1\tau_j}\bar{\varrho}^2 f_{011} + \frac{1}{2}f_{200} + \frac{1}{2}\bar{\varrho}^2 f_{020})\bar{z}^2 \end{array} \right) \\ &\left( \begin{array}{c} \langle \beta_{14} \left( w_{11}^2(0) + \frac{w_{20}^2(0)}{2} + w_{11}^1(0)\varrho + \frac{w_{20}^1(0)}{2}\bar{\varrho} \right) + \beta_{13} \left( w_{11}^1(0) + \frac{w_{20}^1(0)}{2} \right), 1 \rangle \\ \langle f_{110}e^{-i\omega_1\tau_j} \left( w_{11}^2(-1) + e^{2i\omega_1\tau_j} \frac{w_{20}^2(-1)}{2} + w_{11}^1(-1)\varrho + e^{2i\omega_1\tau_j} \frac{w_{20}^1(-1)}{2}\bar{\varrho} \right) \\ + f_{101} \left( e^{-i\omega_1\tau_j} w_{11}^2(0) + e^{i\omega_1\tau_j} \frac{w_{20}^2(0)}{2} + w_{11}^1(-1)\varrho + \frac{w_{20}^1(-1)}{2}\bar{\varrho} \right) \\ + f_{011} \left( e^{-i\omega_1\tau_j} w_{11}^2(0)\varrho + e^{i\omega_1\tau_j} \frac{w_{20}^2(0)}{2}\bar{\varrho} + w_{11}^1(-1)\varrho + \frac{w_{20}^1(-1)}{2}\bar{\varrho} \right) \\ + \frac{1}{2}f_{200} \left( 2e^{-i\omega_1\tau_j} w_{11}^1(-1) + e^{i\omega_1\tau_j} w_{20}^1(-1) \right) \\ + \frac{1}{2}f_{020} \left( 2e^{-i\omega_1\tau_j} w_{11}^1(-1)\varrho + e^{i\omega_1\tau_j} w_{20}^2(-1)\varrho \right), 1 \rangle \end{array} \right) z\bar{z} \\ &+ \cdots, \end{aligned}$$

where

$$\langle w_{ij}^n(\theta), 1 \rangle = \frac{1}{\pi} \int_0^\pi w_{ij}^n(\theta)(x) dx, \quad i+j=2, n=1,2.$$

Noting that  $\Psi_1(0) - i\Psi_2(0) = \frac{2(1-i\omega_1)}{(1+\omega_1^2)(1+\varrho v)}(1, v)$ . Therefore,

$$\begin{aligned} g_{20} &= \frac{\tau_j(1-i\omega_1)}{(1+\omega_1^2)(1+\varrho v)} \\ &\times \left[ \left( \varrho\beta_{14} + \frac{1}{2}\beta_{13} \right) + e^{-2i\omega_1\tau_j} \left( \varrho f_{110} + e^{i\omega_1\tau_j}\varrho f_{101} + e^{i\omega_1\tau_j}\varrho^2 f_{011} + \frac{1}{2}f_{200} + \frac{1}{2}\varrho^2 f_{020} \right) v \right], \end{aligned}$$

$$\begin{aligned} g_{11} &= \frac{\tau_j(1-i\omega_1)}{(1+\omega_1^2)(1+\varrho v)} \\ &\times \left\{ [(\bar{\varrho} + \varrho)\beta_{14} + \beta_{13}] \right. \\ &\quad \left. + \left[ (\bar{\varrho} + \varrho)f_{110} + e^{-i\omega_1\tau_j}\bar{\varrho}(f_{101} + \varrho f_{011}) + e^{i\omega_1\tau_j}\varrho(f_{101} + \bar{\varrho}f_{011}) + f_{200} + \varrho\bar{\varrho}f_{020} \right] v \right\}, \end{aligned}$$

$$g_{02} = \overline{g_{20}},$$

$$\begin{aligned}
 g_{21} = & \frac{2\tau_j(1 - i\omega_1)}{(1 + \omega_1^2)(1 + \varrho v)} \\
 & \times \left[ \left\langle \beta_{14} \left( w_{11}^2(0) + \frac{w_{20}^2(0)}{2} + w_{11}^1(0)\varrho + \frac{w_{20}^1(0)}{2}\bar{\varrho} \right) + \beta_{13} \left( w_{11}^1(0) + \frac{w_{20}^1(0)}{2} \right), 1 \right\rangle \right. \\
 & + \left\langle f_{110}e^{-i\omega_1\tau_j} \left( w_{11}^2(-1) + e^{2i\omega_1\tau_j} \frac{w_{20}^2(-1)}{2} + w_{11}^1(-1)\varrho + e^{2i\omega_1\tau_j} \frac{w_{20}^1(-1)}{2}\bar{\varrho} \right) \right. \\
 & + f_{101} \left( e^{-i\omega_1\tau_j} w_{11}^2(0) + e^{i\omega_1\tau_j} \frac{w_{20}^2(0)}{2} + w_{11}^1(-1)\varrho + \frac{w_{20}^1(-1)}{2}\bar{\varrho} \right) \\
 & + f_{011} \left( e^{-i\omega_1\tau_j} w_{11}^2(0)\varrho + e^{i\omega_1\tau_j} \frac{w_{20}^2(0)}{2}\bar{\varrho} + w_{11}^1(-1)\varrho + \frac{w_{20}^1(-1)}{2}\bar{\varrho} \right) \\
 & + \frac{1}{2}f_{200} \left( 2e^{-i\omega_1\tau_j} w_{11}^1(-1) + e^{i\omega_1\tau_j} w_{20}^1(-1) \right) \\
 & \left. + \frac{1}{2}f_{020} \left( 2e^{-i\omega_1\tau_j} w_{11}^2(-1)\varrho + e^{i\omega_1\tau_j} w_{20}^2(-1)\varrho \right), 1 \right\rangle v \Big].
 \end{aligned}$$

To determine the properties of the Hopf bifurcation, we need to compute  $w_{ij}$ ,  $i + j = 2$ , since  $w_{20}(\theta)$  and  $w_{11}(\theta)$  for  $(\theta \in [-1, 0])$  appear in  $g_{21}$ .

In addition, we can rewrite (3.12) as

$$\dot{w}(z, \bar{z}) = w_{20}z\dot{z} + w_{11}(\dot{z}\bar{z} + z\dot{\bar{z}}) + w_{02}\bar{z}\dot{\bar{z}} + \dots \quad (3.16)$$

and

$$A_{\tau_j}w = A_{\tau_j}w_{20}\frac{z^2}{2} + A_{\tau_j}w_{11}z\bar{z} + A_{\tau_j}w_{02}\frac{\bar{z}^2}{2} + \dots \quad (3.17)$$

According to [29], we can know

$$\dot{w} = A_{\tau_j}w + H(z, \bar{z}), \quad (3.18)$$

where

$$H(z, \bar{z}) = H_{20}\frac{z^2}{2} + H_{11}z\bar{z} + H_{02}\frac{\bar{z}^2}{2} + \dots \quad (3.19)$$

and  $H_{ij} \in \mathcal{P}_{QC}$ ,  $i + j = 2$ .

Thus, by using the chain rule

$$\dot{w} = \frac{\partial w(z, \bar{z})}{\partial z} \dot{z} + \frac{\partial w(z, \bar{z})}{\partial \bar{z}} \dot{\bar{z}}.$$

And according to (3.14) and (3.18), we can obtain

$$\begin{cases}
 (2i\omega_1 - A_{\tau_j})w_{20} = H_{20}, \\
 -A_{\tau_j}w_{11} = H_{11}, \\
 (-2i\omega_1 - A_{\tau_j})w_{02} = H_{02}.
 \end{cases} \quad (3.20)$$

Noticing that  $A_{\tau_j}$  has only two eigenvalues  $\pm i\omega_1$ , therefore, (3.20) has the unique solution  $w_{ij}$  ( $i + j = 2$ ) in  $\mathcal{P}_{QC}$  and

$$\begin{cases}
 w_{20} = (2i\omega_1 - A_{\tau_j})^{-1}H_{20}, \\
 w_{11} = -A_{\tau_j}^{-1}H_{11}, \\
 w_{02} = (-2i\omega_1 - A_{\tau_j})^{-1}H_{02}.
 \end{cases} \quad (3.21)$$

From (3.19), then, for  $-1 \leq \theta < 0$ ,

$$\begin{aligned}
H(z, \bar{z}) &= -\Phi(\theta)\Psi(0) \langle f(U_t, 0), f_1 \rangle \cdot f_1 \\
&= -\left( \frac{p_1(\theta) + p_2(\theta)}{2}, \frac{p_1(\theta) - p_2(\theta)}{2i} \right) \begin{pmatrix} \Psi_1(0) \\ \Psi_2(0) \end{pmatrix} \langle f(U_t, 0), f_1 \rangle \cdot f_1 \\
&= -\frac{1}{2} [p_1(\theta) (\Psi_1(0) - i\Psi_2(0)) + p_2(\theta) (\Psi_1(0) + i\Psi_2(0))] \langle f(U_t, 0), f_1 \rangle \cdot f_1 \\
&= -\frac{1}{4} [g_{20}p_1(\theta) + \bar{g}_{02}p_2(\theta)] z^2 \cdot f_1 + \dots
\end{aligned}$$

So, for  $-1 \leq \theta < 0$ ,

$$H_{20}(\theta) = -\frac{1}{2} [g_{20}p_1(\theta) + \bar{g}_{02}p_2(\theta)] \cdot f_1, \quad H_{11}(\theta) = 0.$$

When  $\theta = 0$ ,

$$\begin{aligned}
H_{20}(0) &= \frac{i\tau_j}{2} \begin{pmatrix} \varrho\beta_{14} + \frac{1}{2}\beta_{13} \\ e^{-2i\omega_1\tau_j} (\varrho f_{110} + e^{i\omega_1\tau_j} \varrho f_{101} + e^{i\omega_1\tau_j} \varrho^2 f_{011} + \frac{1}{2}f_{200} + \frac{1}{2}\varrho^2 f_{020}) \end{pmatrix} \\
&\quad - \frac{1}{2} [g_{20}p_1(0) + \bar{g}_{02}p_2(0)] \cdot f_1, \\
H_{11}(0) &= 0.
\end{aligned}$$

Using the definition of  $A_{\tau_j}$ , for  $-1 \leq \theta < 0$ , we have

$$w_{20}(\theta) = 2i\omega_1 w_{20}(\theta) + \frac{1}{2} [g_{20}p_1(\theta) + \bar{g}_{02}p_2(\theta)] \cdot f_1, \quad -1 \leq \theta < 0.$$

Note that  $p_1(\theta) = p_1(0) e^{i\omega_1\theta}$ ,  $-1 \leq \theta \leq 0$ , hence

$$w_{20}(\theta) = \frac{i}{2} \left[ \frac{g_{20}}{\omega_1} p_1(\theta) + \frac{\bar{g}_{02}}{3\omega_1} p_2(\theta) \right] \cdot f_1 + e^{2i\omega_1\theta} E, \quad -1 \leq \theta < 0, \quad (3.22)$$

and

$$E = w_{20}(0) - \frac{i}{2} \left[ \frac{g_{20}}{\omega_1} p_1(0) + \frac{\bar{g}_{02}}{3\omega_1} p_2(0) \right] \cdot f_1. \quad (3.23)$$

By the definition of  $A_{\tau_j}$  again, and combining (3.17) and (3.20), we have

$$\begin{aligned}
&2i\omega_1 \left[ \frac{ig_{20}}{2\omega_1} p_1(0) \cdot f_1 + \frac{i\bar{g}_{02}}{6\omega_1} p_2(0) \cdot f_1 + E \right] \\
&\quad - \tau_j \Delta \left[ \frac{ig_{20}}{2\omega_1} p_1(0) \cdot f_1 + \frac{i\bar{g}_{02}}{6\omega_1} p_2(0) \cdot f_1 + E \right] \\
&\quad - L^*(\tau_j) \left[ \frac{ig_{20}}{2\omega_1} p_1(\theta) \cdot f_1 + \frac{i\bar{g}_{02}}{6\omega_1} p_2(\theta) \cdot f_1 + E e^{2i\omega_1\theta} \right] \\
&= \frac{i\tau_j}{2} \begin{pmatrix} \varrho\beta_{14} + \frac{1}{2}\beta_{13} \\ e^{-2i\omega_1\tau_j} (\varrho f_{110} + e^{i\omega_1\tau_j} \varrho f_{101} + e^{i\omega_1\tau_j} \varrho^2 f_{011} + \frac{1}{2}f_{200} + \frac{1}{2}\varrho^2 f_{020}) \end{pmatrix} \\
&\quad - \frac{1}{2} [g_{20}p_1(0) + \bar{g}_{02}p_2(0)] \cdot f_1.
\end{aligned}$$

For

$$\tau_j \Delta p_1(0) \cdot f_1 + L_*(\tilde{\tau})(p_1(\theta) \cdot f_1) = i\omega_1 p_1(0) \cdot f_1,$$

$$\tau_j \Delta p_2(0) \cdot f_1 + L_*(\tau_j)(p_2(\theta) \cdot f_1) = -i\omega_1 p_2(0) \cdot f_1,$$

then

$$\begin{aligned} & 2i\omega_1 E - \tau_j \Delta E - L_*(\tau_j)(E e^{2i\omega_1 \theta}) \\ &= \frac{i\tau_j}{2} \left( e^{-2i\omega_1 \tau_j} \left( \varrho f_{110} + e^{i\omega_1 \tau_j} \varrho f_{101} + e^{i\omega_1 \tau_j} \varrho^2 f_{011} + \frac{1}{2} f_{200} + \frac{1}{2} \varrho^2 f_{020} \right) \right. \\ & \quad \left. \varrho \beta_{14} + \frac{1}{2} \beta_{13} \right) \end{aligned}$$

From the above expression, we can see easily that

$$\begin{aligned} E &= \begin{pmatrix} E_1 \\ E_2 \end{pmatrix} \\ &= \frac{i}{2} \begin{pmatrix} -2i\omega_1 + \beta_{11} & \beta_{12} \\ \beta_{21} & -2i\omega_1 + \beta_{22} \end{pmatrix}^{-1} \\ & \quad \times \begin{pmatrix} \varrho \beta_{14} + \frac{1}{2} \beta_{13} \\ e^{-2i\omega_1 \tau_j} \left( \varrho f_{110} + e^{i\omega_1 \tau_j} \varrho f_{101} + e^{i\omega_1 \tau_j} \varrho^2 f_{011} + \frac{1}{2} f_{200} + \frac{1}{2} \varrho^2 f_{020} \right) \end{pmatrix}. \end{aligned}$$

Thus, from the definition of  $g_{20}$  and  $g_{02}$  we see easily that

$$w_{20}(0) = \frac{i}{2} \left( \frac{g_{20}}{\omega_1} + \frac{\bar{g}_{02}}{3\omega_1} \right) \begin{pmatrix} 1 \\ \varrho \end{pmatrix} + E$$

and

$$w_{20}(-1) = \frac{1}{2} \left( \frac{g_{20}}{\omega_1} - \frac{\bar{g}_{02}}{3\omega_1} \right) \begin{pmatrix} 1 \\ \varrho \end{pmatrix} - E.$$

Therefore,  $g_{21}$  can be determined by the parameters.

In fact, by a transformation

$$z = \vartheta + a_{20} \frac{\vartheta^2}{2} + a_{11} \vartheta \bar{\vartheta} + a_{20} \frac{\bar{\vartheta}^2}{2} + \dots,$$

where  $a_{20} = \frac{g_{20}}{i\omega_1}$ ,  $a_{11} = \frac{ig_{11}}{\omega_1}$ ,  $a_{02} = \frac{ig_{02}}{3\omega_1}$ . Then, (3.15) can be written as Poincaré normal form

$$\dot{\vartheta} = i\omega_1 \vartheta + c_1(0) \vartheta |\vartheta|^2 + O(|\vartheta|^5),$$

where

$$c_1(0) = \frac{i}{2\omega_1} \left( g_{11} g_{20} - 2|g_{11}|^2 - \frac{|g_{02}^2|}{3} \right) + \frac{g_{21}}{2}.$$

Thus, we can compute  $\epsilon_2 = 2 \operatorname{Re}(c_1(0))$ . Hence, we have the following result.

**Theorem 3.1.** *If  $\epsilon_2 < 0$ , then the spatially periodic solutions are stable; if  $\epsilon_2 > 0$ , then the spatially periodic solutions are unstable.*

## 4 Conclusions and numerical simulations

In this paper, by studying the existence and stability of spatially periodic solutions for a delay Leslie–Gower diffusion system, we obtain that the system can generate the spatially nonhomogeneous periodic solutions when the diffusive rates are suitably small. We discover

that system (1.2) have more abundant dynamic behavior than system (1.1), this indicates that diffusion plays a fundamental role in classifying the rich dynamics. In addition, we considered the stability of periodic solutions by applying the normal form theory of partial functional differential equations.

To illustrate the analytical results obtained, we give some numerical simulations and consider the following case of model (1.2)

$$\begin{cases} \frac{\partial u(t,x)}{\partial t} = 10\Delta u(t,x) + u(t,x)[1 - u(t,x)] - u(t,x)v(t,x), & t > 0, x \in (0, \pi) \\ \frac{\partial v(t,x)}{\partial t} = 0.01\Delta v(t,x) + v(t,x)\left[1 - \frac{v(t-\tau,x)}{2u(t-\tau,x)}\right], & t > 0, x \in (0, \pi), \\ u(t,x) = 0.2, v(t,x) = 0.2, & (t,x) \in [-\tau, 0] \times (0, \pi), \end{cases} \quad (4.1)$$

which has a positive equilibrium  $E^* = (0.3333, 0.6667)$ . From (2.15), (2.16) and system (4.1), we know  $\omega_1 = 1.0638$ ,  $\tau_0 = 1.4796$ . Therefore, we know from Theorem 2.3 that the positive equilibrium  $E^* = (0.3333, 0.6667)$  is asymptotically stable when  $\tau \in [0, 1.4796)$ . These properties are illustrated by the numerical simulation in Figs. 1–10

When  $\tau$  passes through the critical value  $\tau_0$ ,  $E^* = (0.3333, 0.6667)$  loss its stability, a family of periodic solution bifurcates from equilibrium  $E^* = (0.3333, 0.6667)$  which is depicted by the numerical simulation in Figs. 11–12. From Theorem 3.1 and system (4.1), we can compute  $c_1(0) = -2.0788e + 0.03 - 6.9946e + 0.02i$  by the software package *Matlab R2009b*, and  $\epsilon_2 = 2 \operatorname{Re}(c_1(0)) < 0$ . Therefore, the bifurcated periodic solutions are orbitally asymptotically stable on the center manifold. These properties are depicted by the numerical simulation in Figs. 11–12.

## Acknowledgements

The authors are very grateful to the anonymous reviewers for their careful reading, useful comments and constructive suggestions for the improvement of the manuscript of the present research work. Jia-Fang Zhang was supported by the NNSF of China (11226153, 11301147), the Fundamental Research Fund of Henan University (2012YBZR032) and the Science and Technology Research Foundation Awarded by Educational Commission of Henan Province of China (14A110021).

## References

- [1] E. BERETTA, Y. KUANG, Global analysis in some delayed ratio-dependent predator–prey systems, *Nonlinear Anal.* **32**(1998), 381–408. [MR1610586](#)
- [2] S. BUSENBERG, J. MAHAFFY, Interaction of spatial diffusion and delays in models of genetic control by repression, *J. Math. Biol.* **22**(1985), 313–333. [MR0813402](#)
- [3] S. BUSENBERG, W. HUANG, Stability and Hopf bifurcation for a population delay model with diffusion effects, *J. Differential Equations* **124**(1996), 80–107. [MR1368062](#)
- [4] X. CHANG, J. WEI, Hopf bifurcation and optimal control in a diffusive predator–prey system with time delay and prey harvesting, *Nonlinear Anal. Model. Control* **17**(2012), 379–409. [MR3009928](#)

- [5] Y. K. CHANG, J. J. NIETO, Some new existence results for fractional differential inclusions with boundary conditions, *Math. Comput. Model.* **49**(2009), 605–609. [MR2483665](#)
- [6] S. CHEN, J. SHI, J. WEI, Global stability and Hopf bifurcation in a delayed diffusive Leslie–Gower predator–prey system, *Int. J. Bifurcation Chaos Appl. Sci. Eng.* **22**(2012), No. 3, 1250061, 11 pp. [MR2916689](#)
- [7] S. N. CHOW, J. K. HALE, *Methods of bifurcation theory*, Springer-Verlag, New York, Heidelberg–Berlin, 1982.
- [8] G.H. CUI, X.P. YAN, Stability and bifurcation analysis on a three-species food chain system with two delays, *Commun. Nonlinear Sci. Numer. Simul.* **16**(2011), 3704–3720. [MR2787817](#)
- [9] J. M. CUSHING, *Integrodifferential equations and delay models in population dynamics*, Springer-Verlag, Heidelberg, 1977. [MR0496838](#)
- [10] Y. H. FAN, L. L. WANG, Periodic solutions in a delayed predator-prey model with non-monotonic functional response, *Nonlinear Anal. Real World Appl.* **10**(2009), 3275–3284. [MR2523287](#)
- [11] T. FARIA, Normal forms and Hopf bifurcation for partial differential equations with delays, *Trans. Amer. Math. Soc.* **352**(2000), 2217–2238. [MR1491862](#)
- [12] T. FARIA, Stability and bifurcation for a delayed predator-prey model and the effect of diffusion, *J. Math. Anal. Appl.* **254**(2001), 433–463. [MR1805517](#)
- [13] S. GOURLEY, S. RUAN, Convergence and travelling fronts in functional differential equations with nonlocal term: a competition model, *SIAM J. Math. Anal.* **35**(2003), 806–822. [MR2048407](#)
- [14] J. K. HALE, *Theory of functional differential equations*, Springer-Verlag, New York, 1977. [MR0508721](#)
- [15] B. D. HASSARD, N. D. KAZARINOFF, Y. H. WAN, *Theory and applications of Hopf bifurcation*, Cambridge University Press, Cambridge, 1981. [MR603442](#)
- [16] G. HU, X. LI, Y. WANG, Pattern formation and spatiotemporal chaos in a reaction-diffusion predator–prey system, *Nonlinear Dynam.* **81**(2015), 265–275. [MR3355029](#)
- [17] X. Y. MENG, H. F. HUO, X. B. ZHANG, Stability and global Hopf bifurcation in a delayed food web consisting of a prey and two predators, *Commun. Nonlinear Sci. Numer. Simul.* **16**(2011), 4335–4348. [MR2806750](#)
- [18] X. K. SUN, H. F. HUO, H. XIANG, Bifurcation and stability analysis in predator-prey model with a stage-structure for predator, *Nonlinear Dyn.* **58**(2009), 497–513. [MR2562945](#)
- [19] Y. KUANG, *Delay differential equations with application in population dynamics*, Academic Press, New York, 1993. [MR1218880](#)
- [20] W. T. LI, X. P. YAN, C. H. ZHANG, Stability and Hopf bifurcation for a delayed cooperation diffusion system with Dirichlet boundary conditions, *Chaos Solitons Fractals* **38**(2008), 227–237. [MR2417659](#)

- [21] J. LIU, J. WEI, On Hopf bifurcation of a delayed predator-prey system with diffusion, *Int. J. Bifurcation Chaos Appl. Sci. Eng.* **23**(2013), No. 2, 1350023, 13 pp. [MR3041473](#)
- [22] A. J. LOTKA, *Elements of physical biology*, Williams and Wilkins, New York, 1925. [MR0094263](#)
- [23] Z. P. MA, H. F. HUO, C. Y. LIU, Stability and Hopf bifurcation analysis on a predator-prey model with discrete and distributed delays, *Nonlinear Anal. Real World Appl.* **10**(2009), 1160–1172. [MR2474289](#)
- [24] R. H. MARTIN, H. L. SMITH, Reaction–diffusion systems with time delays: monotonicity, invariance, comparison and convergence, *J. Reine Angew. Math.* **413**(1991), 1–35. [MR1089794](#)
- [25] J. D. MURRAY, *Mathematical biology*, Springer-Verlag, New York. 1993. [MR1239892](#); [url](#)
- [26] H. B. SHI, W. T. LI, G. LIN, Positive steady states of a diffusive predator-prey system with modified Holling–Tanner functional response, *Nonlinear Anal. Real World Appl.* **11**(2010), 3711–3721. [MR2683824](#)
- [27] Y. SONG, S. YUAN, J. ZHANG, Bifurcation analysis in the delayed Leslie–Gower predator-prey system, *Appl. Math. Modelling* **33**(2009), 4049–4061. [MR2541833](#)
- [28] V. VOLTERRA, Variazioni e fluttuazioni del numero d’individui in specie animali conviventi (in Italian), *Mem. Acad. Licei.* **2**(1926), 31–113.
- [29] J. WU, *Theory and applications of partial functional differential equations*, Springer-Verlag, New York, 1996. [MR1415838](#); [url](#)
- [30] C. XU, Y. SHAO, Bifurcations in a predator–prey model with discrete and distributed time delay, *Nonlinear Dynam.* **67**(2012), 2207–2223. [MR2877478](#)
- [31] X. YAN, Stability and Hopf bifurcation for a delayed prey–predator system with diffusion effects, *Appl. Math. Comput.* **192**(2007), 552–566. [MR2385621](#)
- [32] W. ZUO, J. WEI, Stability and Hopf bifurcation in a diffusive predator–prey system with delay effect, *Nonlinear Anal. Real World Appl.* **12**(2011), 1998–2011. [MR2800995](#)



**Figures**




Figure 1: The trajectory graph( $u(t,x)$ ) of system (4.1) with  $0.3 = \tau < \tau_0 = 1.4796$ .




Figure 2: The trajectory graph( $v(t,x)$ ) of system (4.1) with  $0.3 = \tau < \tau_0 = 1.4796$ .




Figure 3: The trajectory graph( $u(t,x)$ ) of system (4.1) with  $0.6 = \tau < \tau_0 = 1.4796$ .




Figure 4: The trajectory graph( $v(t,x)$ ) of system (4.1) with  $0.6 = \tau < \tau_0 = 1.4796$ .




Figure 5: The trajectory graph( $u(t,x)$ ) of system (4.1) with  $0.8 = \tau < \tau_0 = 1.4796$ .




Figure 6: The trajectory graph( $v(t,x)$ ) of system (4.1) with  $0.8 = \tau < \tau_0 = 1.4796$ .




Figure 7: The trajectory graph( $u(t,x)$ ) of system (4.1) with  $1.0 = \tau < \tau_0 = 1.4796$ .




Figure 8: The trajectory graph( $v(t,x)$ ) of system (4.1) with  $1.0 = \tau < \tau_0 = 1.4796$ .




Figure 9: The trajectory graph( $u(t,x)$ ) of system (4.1) with  $1.3 = \tau < \tau_0 = 1.4796$ .




Figure 10: The trajectory graph( $v(t,x)$ ) of system (4.1) with  $1.3 = \tau < \tau_0 = 1.4796$ .




Figure 11: The trajectory graph( $u(t,x)$ ) of system (4.1) with  $1.5 = \tau > \tau_0 = 1.4796$ .




Figure 12: The trajectory graph( $v(t,x)$ ) of system (4.1) with  $1.5 = \tau > \tau_0 = 1.4796$ .