

Positive Solutions to an N^{th} Order Right Focal Boundary Value Problem

Mariette Maroun
Department of Mathematics and Physics
University of Louisiana at Monroe
Monroe, Louisiana 71209, USA
maroun@ulm.edu

Abstract

The existence of a positive solution is obtained for the n^{th} order right focal boundary value problem $y^{(n)} = f(x, y)$, $0 < x \leq 1$, $y^{(i)}(0) = y^{(n-2)}(p) = y^{(n-1)}(1) = 0, i = 0, \dots, n-3$, where $\frac{1}{2} < p < 1$ is fixed and where $f(x, y)$ is singular at $x = 0, y = 0$, and possibly at $y = \infty$. The method applies a fixed-point theorem for mappings that are decreasing with respect to a cone.

Key words: Fixed point theorem, boundary value problem.

AMS Subject Classification: 34B15.

1 Introduction

In this paper, we establish the existence of a positive solution for the n^{th} order right focal boundary value problem,

$$y^{(n)} = f(x, y), \text{ for } x \in (0, 1], \quad (1)$$

$$y^{(i)}(0) = y^{(n-2)}(p) = y^{(n-1)}(1) = 0, \quad i = 0, \dots, n-3, \quad (2)$$

where $\frac{1}{2} < p < 1$ is fixed and $f(x, y)$ is singular at $x = 0, y = 0$, and may be singular at $y = \infty$.

We assume the following conditions hold for f :

- (H1) $f(x, y) : (0, 1] \times (0, \infty) \longrightarrow (0, \infty)$ is continuous, and $f(x, y)$ is decreasing in y for every x .
- (H2) $\lim_{y \rightarrow 0^+} f(x, y) = +\infty$ and $\lim_{y \rightarrow \infty} f(x, y) = 0$ uniformly on compact subsets of $(0, 1]$.

We reduce the problem to a third order integro-differential problem. We establish decreasing operators for which we find fixed points that are solutions to this third order problem. Then, we use Gatica, Olikier, and Waltman methods to find a positive solution to the integro-differential third order problem. We integrate the positive solution $n - 3$ times to obtain the positive solution to the n^{th} order right focal boundary value problem. The role of $\frac{1}{2} < p < 1$ is fundamental for the positivity of the Green's function which in turn is fundamental for the positivity of desired solutions. The existence of positive solutions to a similar third order right focal boundary value problem was established in [22].

Singular boundary value problems for ordinary differential equations have arisen in numerous applications, especially when only positive solutions are useful. For example, when $n = 2$, Taliaferro [28] has given a nice treatment of the general problem, Callegari and Nachman [9] have studied existence questions of this type in boundary layer theory, and Lunning and Perry [21] have established constructive results for generalized Emden-Fowler boundary value problems. Also, Bandle, Sperb, and Stakgold [3] and Bobisud, *et al.* [6], [7], [8], have obtained results for singular boundary value problems that arise in reaction-diffusion theory, while Callegari and Nachman [10] have considered such boundary conditions in non-Newtonian fluid theory as well as in the study of pseudoplastic fluids. Nachman and Callegari point to applications in glacial advance and transport of coal slurries down conveyor belts. See [10] for references. Other applications for these boundary value problems appear in problems such as in draining flows [1], [5] and semi-positone and positone problems [2].

In addition, much attention has been devoted to theoretical questions for singular boundary value problems. In some studies on singular boundary value problems, the underlying technique has been to obtain *a priori* estimates on solutions to an associated two-parameter family of problems, and then use these bounds along with topological transversality theorems to obtain solutions of the original problem; for example, see Granas, Guenther, and Lee [15] and Dunninger and Kurtz [11]. This method has been fairly exploited in a number of recent papers by O'Regan, [24], [25], [26]. Baxley [4] also used to some degree this latter technique in his work on singular boundary value problems for membrane response of a spherical cap. Wei [30] gave necessary and sufficient conditions for the existence of positive solutions for the singular Emden-Fowler equation satisfying Sturm-Liouville boundary

conditions employing upper and lower solutions methods. Guoliang [16] also gave necessary and sufficient conditions for a higher order singular boundary value problem, using superlinear and sublinear conditions to show the existence of a positive solution.

2 Definitions and Properties of Cones

In this section, we begin by giving some definitions and some properties of cones in a Banach space.

Let $(\mathcal{B}, \|\cdot\|)$ be a real Banach space. A nonempty set $\mathcal{K} \subset \mathcal{B}$ is called a *cone* if the following conditions are satisfied:

- (a) the set \mathcal{K} is closed;
- (b) if $u, v \in \mathcal{K}$ then $\alpha u + \beta v \in \mathcal{K}$, for all $\alpha, \beta \geq 0$;
- (c) $u, -u \in \mathcal{K}$ imply $u = 0$.

Given a cone, \mathcal{K} , a *partial order*, \leq , is induced on \mathcal{B} by $x \leq y$, for $x, y \in \mathcal{B}$ iff $y - x \in \mathcal{K}$. (For clarity we sometimes write $x \leq y$ (w.r.t. \mathcal{K})). If $x, y \in \mathcal{B}$ with $x \leq y$, let $\langle x, y \rangle$ denote the *closed order interval between x and y* given by, $\langle x, y \rangle = \{z \in \mathcal{K} | x \leq z \leq y\}$. A cone \mathcal{K} is *normal* in \mathcal{B} provided there exists $\delta > 0$ such that $\|e_1 + e_2\| \geq \delta$, for all $e_1, e_2 \in \mathcal{K}$ with $\|e_1\| = \|e_2\| = 1$.

Remark: If \mathcal{K} is a normal cone in \mathcal{B} , then closed order intervals are norm bounded.

3 Gatica, Olikier, and Waltman Fixed Point Theorem

Now we state the fixed point theorem due to Gatica, Olikier, and Waltman on which most of the results of this paper depend.

Theorem 3.1 *Let \mathcal{B} be a Banach space, \mathcal{K} a normal cone in \mathcal{B} , \mathcal{C} a subset of \mathcal{K} such that if x, y are elements of \mathcal{C} , $x \leq y$, then $\langle x, y \rangle$ is contained in \mathcal{C} , and let $T: \mathcal{C} \rightarrow \mathcal{K}$ be a continuous decreasing mapping which is compact on any closed order interval contained in \mathcal{C} . Suppose there exists $x_0 \in \mathcal{C}$ such that $T^2(x_0)$ is defined (where $T^2(x_0) = T(Tx_0)$), and furthermore, Tx_0 and T^2x_0 are order comparable to x_0 . Then T has a fixed point in \mathcal{C} provided that either,*

- (I) $Tx_0 \leq x_0$ and $T^2x_0 \leq x_0$, or $Tx_0 \geq x_0$ and $T^2x_0 \geq x_0$, or
- (II) The complete sequence of iterates $\{T^n x_0\}_{n=0}^{\infty}$ is defined, and there exists $y_0 \in \mathcal{C}$ such that $y_0 \leq T^n x_0$, for every n .

We consider the following Banach space, \mathcal{B} , with associated norm, $\|\cdot\|$:

$$\mathcal{B} = \{u : [0, 1] \rightarrow R \mid u \text{ is continuous}\},$$

$$\|u\| = \sup_{x \in [0, 1]} |u(x)|.$$

We also define a cone, \mathcal{K} , in \mathcal{B} by,

$$\mathcal{K} = \{u \in \mathcal{B} \mid u(x) \geq 0, g(x)u(p) \leq u(x) \leq u(p) \text{ and } u(x) \text{ is concave on } [0, 1]\},$$

where

$$g(x) = \frac{x(2p-x)}{p^2}, \text{ for } 0 \leq x \leq 1.$$

4 The Integral Operator

In this section, we will define a decreasing operator \mathcal{T} that will allow us to use the stated fixed point theorem.

First, we define $k(x)$ by,

$$k(x) = \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} g(s) ds,$$

Given $g(x)$ and $k(x)$ above, we define $g_\theta(x)$ and $k_\theta(x)$, for $\theta > 0$, by

$$g_\theta(x) = \theta \cdot g(x),$$

and

$$k_\theta(x) = \theta \cdot k(x),$$

and we will assume hereafter

$$(H3) \int_0^1 f(x, k_\theta(x)) dx < \infty, \text{ for each } \theta > 0.$$

We note that the function $f(x, y) = \frac{1}{\sqrt[4]{xy}}$ also satisfies (H3).

In particular, for each $\theta > 0$,

$$\int_0^1 f(x, g_\theta(x)) dx = \sqrt[4]{\left(\frac{p^2}{\theta}\right)} \left[\frac{2}{\sqrt[4]{2p}} + 4 \frac{(2p-1)^{\frac{3}{4}} - (2p)^{\frac{3}{4}}}{3(\sqrt[4]{2p})} \right] < \infty.$$

If y is a solution of (1)-(2), then

$$u(x) = y^{(n-3)}(x),$$

is positive and concave. Hence, if in addition $u \in \mathcal{K}$, then $\|u\| = u(p)$.

Also, we get,

$$u''' = f\left(x, \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} u(s) ds\right), \quad (3)$$

$$u(0) = u'(p) = u''(1) = 0. \quad (4)$$

Since $g(x)$ is concave with $g(0) = 0$ and $\|g(x)\| = g(p)$, then we observe, that for each positive solution, $u(x)$, of (3)-(4), there is some $\theta > 0$, such that $g_\theta(x) \leq u(x)$, for $0 \leq x \leq 1$.

Next, we let $\mathcal{D} \subseteq \mathcal{K}$ be defined by

$$\mathcal{D} = \{u \in \mathcal{K} \mid \text{there exists } \theta(u) > 0 \text{ so that } g_\theta(x) \leq u(x), 0 \leq x \leq 1\}.$$

We note that for each $u \in \mathcal{K}$,

$$\|u\| = \sup_{x \in [0,1]} |u(x)| = u(p).$$

Next, we define an integral operator $\mathcal{T} : \mathcal{D} \rightarrow \mathcal{K}$ by

$$(\mathcal{T}u)(x) = \int_0^1 G(x,t) f\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} u(s) ds\right) dt,$$

where $G(x,t)$ is the Green's function for $y''' = 0$ satisfying (4), and given by

$$G(x,t) = \begin{cases} \frac{x(2t-x)}{2}, & x \leq t \leq p, \\ \frac{t^2}{2}, & t \leq x, t \leq p, \\ \frac{x(2p-x)}{2}, & x \leq t, t \geq p, \\ \frac{x(2p-x)}{2} + \frac{(x-t)^2}{2}, & x \geq t \geq p; \end{cases}$$

see [14].

First, we show \mathcal{T} is a decreasing operator. Let $u \in \mathcal{D}$ be given. Then there exists $\theta > 0$ such that $g_\theta(x) \leq u(x)$. Then, by condition (H1), $f(x, u(x)) \leq f(x, g_\theta(x))$. Now, let $u(x) \leq v(x)$ for $u(x), v(x) \in \mathcal{D}$. Then,

$$\int_0^x \frac{(x-s)^{n-4}}{(n-4)!} u(s) ds \leq \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} v(s) ds.$$

Then by condition (H1),

$$f\left(x, \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} v(s) ds\right) \leq f\left(x, \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} u(s) ds\right).$$

And since $G(x, t) > 0$, we have by (H1) and (H3),

$$\begin{aligned} \int_0^1 G(x, t) f\left(t, \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} v(s) ds\right) dt &< \int_0^1 G(x, t) f\left(t, \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} u(s) ds\right) dt \\ &\leq \int_0^1 G(x, t) f\left(t, \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} g_\theta(s) ds\right) dt \\ &= \int_0^1 G(x, t) f(x, k_\theta(x)) \\ &< \infty, \end{aligned}$$

where $g_\theta(x) \leq u(x)$.

Therefore, \mathcal{T} is well-defined on \mathcal{D} and \mathcal{T} is a decreasing operator.

Remark: We claim that $\mathcal{T} : \mathcal{D} \rightarrow \mathcal{D}$. To see this, suppose $u \in \mathcal{D}$ and let

$$w(x) = (\mathcal{T}u)(x) = \int_0^1 G(x, t) f\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} u(s) ds\right) dt \geq 0.$$

Thus, for $0 \leq x \leq 1$, $w(x) \geq 0$. Also by properties of G ,

$$w'''(x) = f\left(x, \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} u(s) ds\right) > 0, \text{ for } 0 < x \leq 1,$$

and $w(x)$ satisfies (5.4). As we argued previously, $\|w\| = w(p)$.

Since we have that $w''(1) = 0$ and $w'''(x) > 0$, then w is concave.

Also, with $w(p) = \|w(x)\|$, then $w(x) \geq w(p)g(x) = g_{w(p)}(x)$. Therefore, $w \in \mathcal{D}$, and $\mathcal{T} : \mathcal{D} \rightarrow \mathcal{D}$.

Remark: It is well-known that $\mathcal{T}u = u$ iff u is a solution of (3)-(4). Hence, we seek solutions of (3)-(4) that belong to \mathcal{D} .

5 *A Priori* Bounds on Norms of Solutions

In this section, we will show that solutions of (3)-(4) have positive *a priori* upper and lower bounds on their norms. The proofs will be done by contradiction.

Lemma 5.1 *If f satisfies (H1)-(H3), then there exists $S > 0$ such that $\|u\| \leq S$ for any solution u of (3)-(4) in \mathcal{D} .*

Proof: We assume that the conclusion of the lemma is false. Then there exists a sequence, $\{u_m\}_{m=1}^\infty$, of solutions of (3)-(4) in \mathcal{D} such that $u_m(x) > 0$, for $x \in (0, 1]$, and

$$\|u_m\| \leq \|u_{m+1}\| \text{ and } \lim_{m \rightarrow \infty} \|u_m\| = \infty.$$

For a solution u of (3)-(4), we have

$$u''' = f\left(x, \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} u(s) ds\right) > 0, \text{ for } 0 < x \leq 1,$$

$$\text{or } u'' < 0, \text{ for } 0 < x \leq 1.$$

This says that u is concave. In particular, the graphs of the sequence of solutions, u_m , are concave. Furthermore, for each m , the boundary conditions (4) and the concavity of u_m give us,

$$u_m(x) \geq u_m(p)g(x) = \|u_m\|g(x) = g_{\|u_m\|}(x) \text{ for all } x,$$

and so for every $0 < c < 1$,

$$\lim_{m \rightarrow \infty} u_m(x) = \infty \text{ uniformly on } [c, 1].$$

Now, let us define

$$M := \max\{G(x, t) : (x, t) \in [0, 1] \times [0, 1]\}.$$

Then, from condition (H2), there exists m_0 such that, for all $m \geq m_0$ and $x \in [p, 1]$,

$$f\left(x, \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} u_m(s) ds\right) \leq \frac{1}{M(1-p)}.$$

Let

$$\theta = \|u_{m_0}\| = u_{m_0}(p).$$

Then, for all $m \geq m_0$,

$$u_m(x) \geq g_\theta(x) = \|u_{m_0}\|g(x), \text{ for } 0 \leq x \leq 1.$$

So, for $m \geq m_0$, and for $0 \leq x \leq 1$, we have

$$\begin{aligned} u_m(x) &= (\mathcal{T}u_m)(x) \\ &= \int_0^1 G(x, t) f\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} u_m(s) ds\right) dt \\ &= \int_0^p G(x, t) f\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} u_m(s) ds\right) dt \\ &\quad + \int_p^1 G(x, t) f\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} u_m(s) ds\right) dt \\ &\leq \int_0^p G(x, t) f\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} g_\theta(s) ds\right) dt + \int_p^1 M \frac{1}{M(1-p)} dt \\ &\leq \int_0^p G(x, t) f(t, k_\theta(t)) dt + 1 \\ &\leq \int_0^1 G(x, t) f(t, k_\theta(t)) dt + 1 \\ &\leq M \int_0^1 f(t, k_\theta(t)) dt + 1. \end{aligned}$$

This is a contradiction to $\lim_{m \rightarrow \infty} \|u_m\| = \infty$. Hence, there exists an $S > 0$ such that $\|u\| < S$ for any solution $u \in \mathcal{D}$ of (3)-(4).

□

Now we deal with positive *a priori* lower bounds on the solution norms.

Lemma 5.2 *If f satisfies (H1)-(H3), then there exists $R > 0$ such that $\|u\| \geq R$ for any solution u of (5.3)-(5.4) in \mathcal{D} .*

Proof: We assume the conclusion of the lemma is false. Then, there exists a sequence $\{u_m\}_{m=1}^\infty$ of solutions of (3)-(4) in \mathcal{D} such that $u_m(x) > 0$, for $x \in (0, 1]$, and

$$\|u_m\| \geq \|u_{m+1}\|$$

and

$$\lim_{m \rightarrow \infty} \|u_m\| = 0.$$

Now we define

$$\bar{m} := \min\{G(x, t) : (x, t) \in [p, 1] \times [p, 1]\} > 0.$$

From condition (H2), $\lim_{y \rightarrow 0^+} f(x, y) = \infty$ uniformly on compact subsets of $(0, 1]$.

Thus, there exists $\delta > 0$ such that, for $x \in [p, 1]$ and $0 < y < \delta$,

$$f(x, y) > \frac{1}{\bar{m}(1-p)}.$$

In addition, there exists m_0 such that, for all $m \geq m_0$ and $x \in (0, 1]$

$$0 < u_m(x) < \frac{\delta}{2},$$

$$0 < \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} u_m(s) ds < \frac{\delta}{2}.$$

So, for $x \in [p, 1]$ and $m \geq m_0$,

$$\begin{aligned} u_m(x) &= (\mathcal{T}u_m)(x) \\ &= \int_0^1 G(x, t) f\left(t, \int_0^t \frac{(x-s)^{n-4}}{(n-4)!} u_m(s) ds\right) dt \\ &\geq \int_p^1 G(x, t) f\left(t, \int_0^t \frac{(x-s)^{n-4}}{(n-4)!} u_m(s) ds\right) dt \end{aligned}$$

$$\begin{aligned}
&\geq \bar{m} \int_p^1 f\left(t, \int_0^t \frac{(x-s)^{n-4}}{(n-4)!} u_m(s) ds\right) dt \\
&> \bar{m} \int_p^1 f\left(t, \frac{\delta}{2}\right) dt \\
&> \bar{m} \int_p^1 \frac{1}{\bar{m}(1-p)} dt \\
&= 1.
\end{aligned}$$

Which is a contradiction to $\lim_{m \rightarrow \infty} \|u_m(x)\| = 0$ uniformly on $[0, 1]$. Thus, there exists $R > 0$ such that $R \leq \|u\|$ for any solution u in \mathcal{D} of (5.3)-(5.4). \square

In summary, there exist $0 < R < S$ such that, for $u \in \mathcal{D}$, a solution of (3)-(4), Lemma 5.1 and Lemma 5.2 give us

$$R \leq \|u\| \leq S.$$

The next section gives the main result, an existence theorem, for this problem.

6 Existence Result

In this section, we will construct a sequence of operators, $\{\mathcal{T}_m\}_{m=1}^\infty$, each of which is defined on all of \mathcal{K} . We will then show, by applications of Theorem 3.1, that each \mathcal{T}_m has a fixed point, ϕ_m , for every m , in \mathcal{K} . Then, we will show that some subsequence of the $\{\phi_m\}_{m=1}^\infty$ converges to a fixed point of \mathcal{T} .

Theorem 6.1 *If f satisfies (H1)-(H3), then (3)-(4) has at least one positive solution u in \mathcal{D} , such that $y(x) = \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} u(s) ds$ is a positive solution of (1)-(2).*

Proof: For all m , let $u_m(x) := \mathcal{T}(m)$, where m is the constant function of that value on $[0, 1]$. In particular,

$$\begin{aligned}
u_m(x) &= \int_0^1 G(x, t) f\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} m ds\right) dt \\
&= \int_0^1 G(x, t) f\left(t, \frac{m(-s)^{n-3}}{(n-3)!}\right) dt, \text{ for } 0 \leq x \leq 1.
\end{aligned}$$

But f is decreasing in its second component, giving us,

$$0 < u_{m+1}(x) \leq u_m(x), \text{ for } 0 \leq x \leq 1.$$

By condition (H2), $\lim_{m \rightarrow \infty} u_m(x) = 0$, uniformly on $[0, 1]$.

Now, we define $f_m(x, y) : (0, 1] \times [0, \infty) \rightarrow (0, \infty)$ by

$$f_m(x, y) = f\left(x, \max\left\{y, \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} u_m(s) ds\right\}\right).$$

Then f_m is continuous and f_m does not possess the singularities as found in f at $y = 0$. Moreover, for $(x, y) \in (0, 1] \times (0, \infty)$ we have that,

$$f_m(x, y) \leq f(x, y)$$

and, moreover,

$$f_m(x, y) = f\left(x, \max\left\{y, \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} u_m(s) ds\right\}\right) \leq f\left(x, \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} u_m(s) ds\right).$$

Next, we define a sequence of operators, $\mathcal{T}_m : \mathcal{K} \rightarrow \mathcal{K}$, for $\phi \in \mathcal{K}$ and $x \in [0, 1]$, by

$$\mathcal{T}_m \phi(x) := \int_0^1 G(x, t) f_m\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} \phi(s) ds\right) dt.$$

It is standard that each \mathcal{T}_m is a compact mapping on \mathcal{K} . Moreover,

$$\begin{aligned} \mathcal{T}_m(0) &= \int_0^1 G(x, t) f_m(t, 0) dt \\ &= \int_0^1 G(x, t) f\left(t, \max\left\{0, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} u_m(s) ds\right\}\right) dt \\ &= \int_0^1 G(x, t) f\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} u_m(s) ds\right) dt \\ &> 0. \end{aligned}$$

Also,

$$\mathcal{T}_m^2(0) = \mathcal{T}_m\left(\int_0^1 G(x, t) f_m(t, 0) dt\right) \geq 0.$$

Then, by theorem (3.1) with $x_0 = 0$, \mathcal{T}_m has a fixed point in \mathcal{K} for every m . Thus, for every m , there exists a $\phi_m \in \mathcal{K}$ so that

$$\mathcal{T}_m \phi_m(x) = \phi_m(x), \quad 0 \leq x \leq 1.$$

Hence, for $m \geq 1$, ϕ_m satisfies the boundary conditions (4) of the problem.

Also,

$$\begin{aligned} \mathcal{T}_m \phi_m(x) &= \int_0^1 G(x, t) f_m\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} \phi_m(s) ds\right) dt \\ &= \int_0^1 G(x, t) f\left(t, \max\left\{\int_0^t \frac{(t-s)^{n-4}}{(n-4)!} \phi_m(s) ds, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} u_m(s) ds\right\}\right) dt \\ &\leq \int_0^1 G(x, t) f\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} u_m(s) ds\right) dt \\ &= \mathcal{T} u_m(x). \end{aligned}$$

That is, $\phi_m(x) = \mathcal{T}_m\phi_m(x) \leq \mathcal{T}u_m(x)$, for $0 \leq x \leq 1$, and for every m .

Proceeding as in lemmas 5.1 and 5.2, there exists $S > 0$ and $R > 0$ such that

$$R < \|\phi_m\| < S$$

for every m .

Now, let $\theta = R$. Since $\phi_m \in \mathcal{K}$, then for $x \in [0, 1]$ and every m ,

$$\phi_m(x) \geq \phi_m(p)g(x) = \|\phi_m\| \cdot g(x) > R \cdot g(x) = \theta \cdot g(x) = g_\theta(x).$$

Thus, with $\theta = R$, $g_\theta(x) \leq \phi_m(x)$ for $x \in [0, 1]$, for every m . Thus, $\{\phi_m\}_{m=1}^\infty$ is contained in the closed order interval $\langle g_\theta, S \rangle$. Therefore, the sequence $\{\phi_m\}_{m=1}^\infty$ is contained in \mathcal{D} . Since \mathcal{T} is a compact mapping, we may assume $\lim_{m \rightarrow \infty} \mathcal{T}\phi_m$ exist; say the limit is ϕ^* .

To conclude the proof of this theorem, we still need to show that

$$\lim_{m \rightarrow \infty} \left(\mathcal{T}\phi_m(x) - \phi_m(x) \right) = 0$$

uniformly on $[0, 1]$. This will give us that $\phi^* \in \langle g_\theta, S \rangle$. Still with $\theta = R$, then $k_\theta(x) = \int_0^1 \frac{(x-s)^{n-4}}{(n-4)!} g_\theta(s) ds \leq \int_0^1 \frac{(x-s)^{n-4}}{(n-4)!} \phi_m(s) ds$ for every m and $0 \leq x \leq 1$. Let $\epsilon > 0$ be given and choose δ , $0 < \delta < 1$, such that

$$\int_0^\delta f(t, k_\theta(t)) dt < \frac{\epsilon}{2M},$$

where again $M := \max\{G(x, t) : (x, t) \in [0, 1] \times [0, 1]\}$. Then, there exists m_0 such that, for $m \geq m_0$ and for $x \in [\delta, 1]$,

$$\int_0^1 \frac{(x-s)^{n-4}}{(n-4)!} u_m(x) ds \leq k_\theta(x) \leq \int_0^1 \frac{(x-s)^{n-4}}{(n-4)!} \phi_m(x) ds.$$

So, for $x \in [\delta, 1]$,

$$\begin{aligned} f_m \left(x, \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} \phi_m(s) ds \right) = \\ f \left(x, \max \left\{ \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} \phi_m(s) ds, \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} u_m(s) ds \right\} \right) = \\ f \left(x, \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} \phi_m(s) ds \right). \end{aligned}$$

Then, for $0 \leq x \leq 1$,

$$\begin{aligned}
\mathcal{T}\phi_m(x) - \phi_m(x) &= \mathcal{T}\phi_m(x) - \mathcal{T}_m\phi_m(x) \\
&= \int_0^1 G(x,t)f\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!}\phi_m(s)ds\right)dt \\
&\quad - \int_0^1 G(x,t)f_m\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!}\phi_m(s)ds\right)dt \\
&= \int_0^\delta G(x,t)f\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!}\phi_m(s)ds\right)dt \\
&\quad + \int_\delta^1 G(x,t)f\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!}\phi_m(s)ds\right)dt \\
&\quad - \int_0^\delta G(x,t)f_m\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!}\phi_m(s)ds\right)dt \\
&\quad - \int_\delta^1 G(x,t)f_m\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!}\phi_m(s)ds\right)dt \\
&= \int_0^\delta G(x,t)f\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!}\phi_m(s)ds\right)dt \\
&\quad - \int_0^\delta G(x,t)f_m\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!}\phi_m(s)ds\right)dt.
\end{aligned}$$

Thus, for $0 \leq x \leq 1$, we have,

$$\begin{aligned}
\left| \mathcal{T}\phi_m(x) - \phi_m(x) \right| &= \left| \int_0^\delta G(x,t) f\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} \phi_m(s) ds\right) dt \right. \\
&\quad \left. - \int_0^\delta G(x,t) f_m\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} \phi_m(s) ds\right) dt \right| \\
&\leq M \left[\left| \int_0^\delta f\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} \phi_m(s) ds\right) dt \right| \right. \\
&\quad \left. + \left| \int_0^\delta f_m\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} \phi_m(s) ds\right) dt \right| \right] \\
&= M \left[\int_0^\delta f\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} \phi_m(s) ds\right) dt \right. \\
&\quad \left. + \int_0^\delta f\left(t, \max\left\{ \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} u_m(s) ds, \right. \right. \right. \\
&\quad \quad \left. \left. \left. \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} \phi_m(s) ds \right\} \right) dt \right] \\
&= M \left[\int_0^\delta f\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} \phi_m(s) ds\right) dt \right. \\
&\quad \left. + \int_0^\delta f\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} \phi_m(s) ds\right) dt \right] \\
&= 2M \int_0^\delta f\left(t, \int_0^t \frac{(t-s)^{n-4}}{(n-4)!} \phi_m(s) ds\right) dt \\
&\leq 2M \int_0^\delta f\left(t, \int_0^1 \frac{(x-s)^{n-4}}{(n-4)!} g_\theta(s) ds\right) ds \\
&= 2M \int_0^\delta f(t, k_\theta(t)) dt \\
&= 2M \frac{\epsilon}{2M} = \epsilon.
\end{aligned}$$

Thus, for $m \geq m_0$,

$$\|\mathcal{T}\phi_m - \phi_m\| < \epsilon.$$

In particular, $\lim_{m \rightarrow \infty} (\mathcal{T}\phi_m(x) - \phi_m(x)) = 0$ uniformly on $[0, 1]$, and for $0 \leq x \leq 1$

$$\begin{aligned}
\mathcal{T}\phi^*(x) &= \mathcal{T}\left(\lim_{m \rightarrow \infty} \mathcal{T}\phi_m(x)\right) \\
&= \mathcal{T}\left(\lim_{m \rightarrow \infty} \phi_m(x)\right) \\
&= \lim_{m \rightarrow \infty} (\mathcal{T}\phi_m(x)) \\
&= \phi^*(x).
\end{aligned}$$

Thus,

$$\mathcal{T}\phi^* = \phi^*,$$

and ϕ^* is a desired solution of (3)-(4).

Now, if $\phi^*(x)$ is the solution of (3)-(4), let $y(x) = \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} \phi^*(s) ds$. Then we have

$$y(0) = \int_0^0 \frac{(x-s)^{n-4}}{(n-4)!} \phi^*(s) ds = 0,$$

and by the Fundamental Theorem of Calculus,

$$y^{(n-3)}(x) = \phi^*(x).$$

Thus,

$$y^{(n-3)}(0) = \phi^*(0) = 0.$$

Also,

$$y^{(n-2)}(x) = (\phi^*)'(x),$$

thus,

$$y^{(n-2)}(p) = (\phi^*)'(p) = 0.$$

And,

$$y^{(n-1)}(x) = (\phi^*)''(x)$$

$$y^{(n-1)}(1) = (\phi^*)''(1) = 0.$$

Moreover,

$$y^{(n)}(x) = (\phi^*)'''(x) = (\mathcal{T}\phi^*)'''(x) = f\left(x, \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} \phi^*(s) ds\right) = f(x, y).$$

Thus, $y(x) = \int_0^x \frac{(x-s)^{n-4}}{(n-4)!} \phi^*(s) ds > 0$, $0 \leq x \leq 1$ solves (1)-(2).

This completes the proof. □

Remark: The results of this paper extend to Boundary Value Problems for $y^{(n)} = f(x, y, y', \cdot, y^{(n-3)})$ under the same boundary conditions.

Acknowledgment: The author is indebted to the referee's suggestions. These have greatly improved this paper.

References

- [1] R. P. Agarwal and D. O'Regan, Singular problems on the infinite interval modelling phenomena in draining flows, *IMA Journal of Applied Mathematics*, **66** (2001), 621-709.

- [2] R. P. Agarwal, D. O'Regan, and P. J. Y. Wong, *Positive Solutions of Differential, Difference and Integral Equations*, Dordrecht, The Netherlands, 1999, 86-105.
- [3] C. Bandle, R. Sperb, and I. Stakgold, Diffusion and reaction with monotone kinetics, *Nonlinear Analysis* **18** (1984), 321-333.
- [4] J. V. Baxley, A singular boundary value problem: membrane response of a spherical cap, *SIAM Journal of Applied Mathematics* **48** (1988), 855-869.
- [5] F. Bernis and L. A. Peletier, Two problems from draining flows involving third order ordinary differential equations, *SIAM Journal of Applied Mathematics* **27** (1996), 515-527.
- [6] L. E. Bobisud, D. O'Regan, and W. D. Royalty, Singular boundary value problems, *Applied Analysis* **23** (1986), 233-243.
- [7] L. E. Bobisud, D. O'Regan, and W. D. Royalty, Existence and nonexistence for a singular boundary value problem, *Applied Analysis* **28** (1988), 245-256.
- [8] L. E. Bobisud, D. O'Regan, and W. D. Royalty, Solvability of some nonlinear singular boundary value problem, *Nonlinear Analysis* **12** (1988), 855-869.
- [9] A. Callegari and A. Nachman, Some singular nonlinear differential equations arising in boundary layer theory, *Journal of Mathematical Analysis and Applications* **64** (1978), 96-105.
- [10] A. Callegari and A. Nachman, A nonlinear singular boundary value problem in the theory of pseudoplastic fluids, *SIAM Journal of Applied Mathematics* **38** (1980), 275-281.
- [11] D. R. Dunninger and J. C. Kurtz, Existence of solutions for some nonlinear boundary value problem, *Journal of Mathematical Analysis and Applications* **115** (1986), 396-405.
- [12] P. Elloe and J. Henderson, Singular nonlinear boundary value problems for higher ordinary differential equations, *Nonlinear Anal.* **17** (1991), 1-10.
- [13] J.A. Gratica, V. Olikar and P. Waltman, Singular nonlinear boundary value problems for second-order differential equations, *J. Differential Equations* **79** (1989), 62-78.
- [14] J. R. Graef and B. Yang, Positive solutions of a nonlinear third order eigenvalue problem, *Dynamic Systems and Applications*, in press.

- [15] A. Granas, R. B. Guenther, and J. W. Lee, A note on the Thomas-Fermi equation, *Zeitschrift für Angewandte Mathematik und Mechanik* **61** (1981), 204-205.
- [16] S. Guoliang, On the existence of positive solution for n order singular boundary value problems, *Chinese Quart. Journal of Math*, Vol. 13, No 3, (1998).
- [17] K. S. Ha and Y. H. Lee, Existence of multiple positive solutions of singular boundary value problems, *Nonlinear Analysis, Methods and Applications* **28** (1997), 1429-1438.
- [18] J. Henderson and W. Yin, Singular $(n, n-k)$ boundary value problems between conjugate and right focal positive solutions of nonlinear problems. *J. Comput. Appl. Math.* **88** (1998), 57-69.
- [19] M. A. Krasnosel'skii, *Positive Solutions to Operator Equations*, Noordhoff, Groningen, The Netherlands, (1964).
- [20] Y. H. Lee, A multiplicity result of positive solutions for the generalized Gelfand type singular boundary value problems *Nonlinear Analysis, Theory, Meth. and Appl.* **30** (1997), 3829-3835.
- [21] C. D. Lunning and W. L. Perry, Positive solutions of negative exponent generalized Emden-Fowler boundary value problems, *SIAM Journal of Applied Mathematics* **12** (1981), 874-879.
- [22] M. Maroun, Positive solutions to a third-order right focal boundary value problem, *Communications on Applied Nonlinear Analysis* **12** (2005), Number 3, 71-82.
- [23] D. O'Regan, Existence of solutions to third order boundary value problems, *Proc. Royal Irish Acad. Sect. A* **90** (1990), 173-189.
- [24] D. O'Regan, Positive solutions to singular and non-singular second-order boundary value problems, *Journal of Mathematical Analysis and Applications* **142** (1989), 40-52.
- [25] D. O'Regan, Some new results for second order boundary value problems, *Journal of Mathematical Analysis and Applications* **148** (1990), 548-570.
- [26] D. O'Regan, Existence of positive solutions to some singular and non-singular second order boundary value problems, *Journal of Differential Equations* **84** (1990), 228-251.
- [27] P. Singh, A second-order singular three point boundary value problem, *Appl. Math. Letters* **17** (2004), 969-976.

- [28] S. Taliaferro, A nonlinear singular boundary value problem, *Nonlinear Analysis* **3** (1979), 897-904.
- [29] W. -B. Qu, Z. -X. Zhang and J. -D. Wu, Positive solutions to a singular second order three-point boundary value problem, *Appl. Math. and Mechanics* **23** (2002), 759-770.
- [30] Z. Wei, Positive solutions of singular sublinear second order boundary value problems, *Systems Science and Mathematical Sciences* **11** (1998), 80-88.
- [31] F. -H. Wong, Positive solutions for singular boundary value problems, *Computers Math. Applic.*, **32** (1996), 41-49.
- [32] X. Yang, Positive solutions for nonlinear singular boundary value problems, *Appl. Math. and Computations* **130** (2002), 225-234.

(Received June 22, 2006)