

Triple positive solutions for a boundary value problem of nonlinear fractional differential equation

Chuanzhi Bai *

Department of Mathematics, Huaiyin Teachers College, Huaian, Jiangsu 223300, P R China

Abstract

In this paper, we investigate the existence of three positive solutions for the nonlinear fractional boundary value problem

$$D_{0+}^{\alpha} u(t) + a(t)f(t, u(t), u''(t)) = 0, \quad 0 < t < 1, \quad 3 < \alpha \leq 4,$$

$$u(0) = u'(0) = u''(0) = u''(1) = 0,$$

where D_{0+}^{α} is the standard Riemann-Liouville fractional derivative. The method involves applications of a new fixed-point theorem due to Bai and Ge. The interesting point lies in the fact that the nonlinear term is allowed to depend on the second order derivative u'' .

Keywords: Fractional derivative; Boundary value problem; Positive solution; Fixed point theorem

1. Introduction

Many papers and books on fractional calculus and fractional differential equations have appeared recently, see for example [1-3, 7-12]. Very recently, El-Shahed [5] used the Krasnoselskii's fixed-point theorem on cone expansion and compression to show the existence and non-existence of positive solutions of nonlinear fractional boundary value problem :

$$D_{0+}^{\alpha} u(t) + \lambda a(t)f(u(t)) = 0, \quad 0 < t < 1, \quad 2 < \alpha \leq 3,$$

$$u(0) = u'(0) = u'(1) = 0,$$

where D_{0+}^{α} is the standard Riemann-Liouville fractional derivative. Kaufmann and Mboumi [6] studied the existence and multiplicity of positive solutions of nonlinear fractional boundary value problem :

$$D_{0+}^{\alpha} u(t) + a(t)f(u(t)) = 0, \quad 0 < t < 1, \quad 1 < \alpha \leq 2,$$

$$u(0) = u'(1) = 0.$$

Motivated by the above works, in this paper we study the existence of three positive solutions for the following nonlinear fractional boundary value problem :

*E-mail address: czbai8@sohu.com

$$D_{0+}^{\alpha} u(t) + a(t)f(t, u(t), u''(t)) = 0, \quad 0 < t < 1, \quad 3 < \alpha \leq 4, \quad (1.1)$$

$$u(0) = u'(0) = u''(0) = u''(1) = 0, \quad (1.2)$$

by using a new fixed-point theorem due to Bai and Ge [4]. Here, the interesting point lies in the fact that the nonlinear term f is allowed to depend on the second order derivative u'' . To the best of the authors knowledge, no one has studied the existence of positive solutions for nonlinear fractional boundary value problems (1.1)-(1.2).

Throughout this paper, we assume that the following conditions hold.

(H₁) $f : [0, 1] \times [0, \infty) \times (-\infty, +\infty) \rightarrow [0, \infty)$ is continuous;

(H₂) $a \in C([0, 1], [0, +\infty))$ and there exists $0 < \omega < 1$ such that $\int_{\omega}^1 [(1-s)^{\alpha-3} - (1-s)^{\alpha-1}] a(s) ds > 0$.

The rest of this paper is organized as follows: In section 2, we present some preliminaries and lemmas. Section 3 is devoted to prove the existence of three positive solutions for BVP (1.1) and (1.2).

2. Preliminaries

For the convenience of the reader, we present some definitions from the cone theory on ordered Banach spaces.

Definition 2.1. The map ψ is said to be a nonnegative continuous concave functional on a cone P of a real Banach space E provided that $\psi : P \rightarrow [0, \infty)$ is continuous and

$$\psi(tx + (1-t)y) \geq t\psi(x) + (1-t)\psi(y), \quad \forall x, y \in P, \quad 0 \leq t \leq 1.$$

Similarly, we say the map ϕ is a nonnegative continuous convex functional on a cone P of a real Banach space E provided that $\phi : P \rightarrow [0, \infty)$ is continuous and

$$\phi(tx + (1-t)y) \leq t\phi(x) + (1-t)\phi(y), \quad \forall x, y \in P, \quad 0 \leq t \leq 1.$$

Definition 2.2. Let $r > a > 0$, $L > 0$ be given and ψ be a nonnegative continuous concave functional and γ, β be nonnegative continuous convex functionals on the cone P . Define convex sets:

$$P(\gamma, r; \beta, L) = \{x \in P \mid \gamma(x) < r, \beta(x) < L\},$$

$$\overline{P}(\gamma, r; \beta, L) = \{x \in P \mid \gamma(x) \leq r, \beta(x) \leq L\},$$

$$P(\gamma, r; \beta, L; \psi, a) = \{x \in P \mid \gamma(x) < r, \beta(x) < L, \psi(x) > a\},$$

$$\overline{P}(\gamma, r; \beta, L; \psi, a) = \{x \in P \mid \gamma(x) \leq r, \beta(x) \leq L, \psi(x) \geq a\}.$$

Suppose that the nonnegative continuous convex functionals γ, β on the cone P satisfy

(A₁) there exists $M > 0$ such that $\|x\| \leq M \max\{\gamma(x), \beta(x)\}$, for $x \in P$;

(A₂) $P(\gamma, r; \beta, L) \neq \emptyset$, for any $r > 0, L > 0$.

Lemma 2.1.[4] *Let P be a cone in a real Banach space E and $r_2 \geq d > b > r_1 > 0, L_2 \geq L_1 > 0$ constants. Assume that γ, β are nonnegative continuous convex functionals on P such that (A₁) and (A₂) are satisfied. ψ is a nonnegative continuous concave functional on P such that $\psi(x) \leq \gamma(x)$ for all $x \in \overline{P}(\gamma, r_2; \beta, L_2)$ and let $T : \overline{P}(\gamma, r_2; \beta, L_2) \rightarrow \overline{P}(\gamma, r_2; \beta, L_2)$ be a completely continuous operator. Suppose*

- (i) $\{x \in \overline{P}(\gamma, d; \beta, L_2; \psi, b) \mid \psi(x) > b\} \neq \emptyset, \psi(Tx) > b$ for $x \in \overline{P}(\gamma, d; \beta, L_2; \psi, b)$,
- (ii) $\gamma(Tx) < r_1, \beta(Tx) < L_1$ for all $x \in \overline{P}(\gamma, r_1; \beta, L_1)$,
- (iii) $\psi(Tx) > b$ for all $x \in \overline{P}(\gamma, r_2; \beta, L_2; \psi, b)$ with $\gamma(Tx) > d$.

Then T has at least three fixed points $x_1, x_2, x_3 \in \overline{P}(\gamma, r_2; \beta, L_2)$. Further,

$$x_1 \in P(\gamma, r_1; \beta, L_1), \quad x_2 \in \{\overline{P}(\gamma, r_2; \beta, L_2; \psi, b) \mid \psi(x) > b\},$$

and

$$x_3 \in \overline{P}(\gamma, r_2; \beta, L_2) \setminus (\overline{P}(\gamma, r_2; \beta, L_2; \psi, b) \cup \overline{P}(\gamma, r_1; \beta, L_1)).$$

The above fixed-point theorem is fundamental in the proof of our main result.

Next, we give some definitions from the fractional calculus.

Definition 2.3. The Riemann-Liouville fractional integral of order $\alpha > 0$ of a function $u : (0, \infty) \rightarrow \mathbb{R}$ is defined as

$$I_{0+}^{\alpha} u(t) = \frac{1}{\Gamma(\alpha)} \int_0^t \frac{u(s)}{(t-s)^{1-\alpha}} ds.$$

Definition 2.4. The Riemann-Liouville fractional derivative of order $\alpha > 0$ of a continuous function $u : (0, \infty) \rightarrow \mathbb{R}$ is

$$D_{0+}^{\alpha} u(t) = \frac{d^n}{dt^n} \left[I_{0+}^{n-\alpha} u(t) \right],$$

where $n = [\alpha] + 1$, provided that the right side is pointwise defined on $(0, \infty)$.

The following lemma is crucial in finding an integral representation of the boundary value problem (1), (2).

Lemma 2.2. [3] *Suppose that $u \in C(0, 1) \cap L(0, 1)$ with a fractional derivative of order $\alpha > 0$. Then*

$$I_{0+}^{\alpha} D_{0+}^{\alpha} u(t) = u(t) + c_1 t^{\alpha-1} + c_2 t^{\alpha-2} + \cdots + c_n t^{\alpha-n}.$$

for some $c_i \in \mathbb{R}$, $i = 1, 2, \dots, n$.

From Lemmas 2.2, we now give an integral representation of the solution of the linearized problem.

Lemma 2.3. *If $y \in C[0, 1]$, then the boundary value problem*

$$D_{0+}^{\alpha} u(t) + y(t) = 0, \quad 0 < t < 1, \quad 3 < \alpha \leq 4, \quad (2.1)$$

$$u(0) = u'(0) = u''(0) = u''(1) = 0, \quad (2.2)$$

has a unique solution

$$u(t) = \int_0^1 G(t, s)y(s)ds,$$

where

$$G(t, s) = \frac{1}{\Gamma(\alpha)} \begin{cases} t^{\alpha-1}(1-s)^{\alpha-3} - (t-s)^{\alpha-1}, & 0 \leq s \leq t \leq 1, \\ t^{\alpha-1}(1-s)^{\alpha-3}, & 0 \leq t < s \leq 1. \end{cases}$$

Proof. From Lemma 2.2, we get

$$u(t) = c_1 t^{\alpha-1} + c_2 t^{\alpha-2} + c_3 t^{\alpha-3} + c_4 t^{\alpha-4} - \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} y(s)ds.$$

By (2.2), there are $c_2 = c_3 = c_4 = 0$, and $c_1 = \frac{1}{\Gamma(\alpha)} \int_0^1 (1-s)^{\alpha-3} y(s)ds$. Hence, the unique solution of BVP (2.1), (2.2) is

$$u(t) = \int_0^1 \frac{t^{\alpha-1}(1-s)^{\alpha-3}}{\Gamma(\alpha)} y(s)ds - \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} y(s)ds = \int_0^1 G(t, s)y(s)ds.$$

The proof is complete.

Lemma 2.4. *$G(t, s)$ has the following properties.*

$$(i) \quad 0 \leq G(t, s) \leq h(s), \quad t, s \in [0, 1],$$

where

$$h(s) = \frac{(1-s)^{\alpha-3} - (1-s)^{\alpha-1}}{\Gamma(\alpha)};$$

$$(ii) \quad G(t, s) \geq \frac{1}{2} t^{\alpha-1} h(s), \quad \text{for } 0 \leq t, s \leq 1.$$

Proof. It is easy to check that (i) holds. Next, we prove (ii) holds. If $t \geq s$, then

$$\begin{aligned} \frac{G(t, s)}{h(s)} &= \frac{t^{\alpha-1}(1-s)^{\alpha-3} - (t-s)^{\alpha-1}}{(1-s)^{\alpha-3} - (1-s)^{\alpha-1}} \\ &\geq \frac{t^2(t-ts)^{\alpha-3} - (t-s)^2(t-ts)^{\alpha-3}}{(1-s)^{\alpha-3} - (1-s)^{\alpha-1}} = \frac{(t-ts)^{\alpha-3}s(2t-s)}{(1-s)^{\alpha-3}s(2-s)} \\ &= \frac{t^{\alpha-3}(2t-s)}{2-s} \geq \frac{t^{\alpha-2}}{2-s} \geq \frac{t^{\alpha-1}}{2}. \end{aligned}$$

If $t \leq s$, then

$$\frac{G(t, s)}{h(s)} = \frac{t^{\alpha-1}(1-s)^{\alpha-3}}{(1-s)^{\alpha-3} - (1-s)^{\alpha-1}} \geq \frac{t^{\alpha-1}(1-s)^{\alpha-3}}{(1-s)^{\alpha-3}} \geq \frac{t^{\alpha-1}}{2}.$$

The proof is complete.

3. Main results

Let $X = \{u \in C^2[0, 1] : u(0) = u'(0) = 0\}$. Then we have the following lemma.

Lemma 3.1. For $u \in X$, $\|u\|_0 \leq \|u'\|_0 \leq \|u''\|_0$, where $\|u\|_0 = \max_{0 \leq t \leq 1} |u(t)|$.

By Lemma 3.1, X is a Banach space when it is endowed with the norm $\|u\| = \|u''\|_0$.

It is easy to know that

$$\frac{\partial}{\partial t} G(t, s) = \frac{\alpha-1}{\Gamma(\alpha)} \begin{cases} t^{\alpha-2}(1-s)^{\alpha-3} - (t-s)^{\alpha-2}, & 0 \leq s \leq t \leq 1, \\ t^{\alpha-2}(1-s)^{\alpha-3}, & 0 \leq t < s \leq 1. \end{cases} \quad (3.1)$$

$$\frac{\partial^2}{\partial t^2} G(t, s) = \frac{(\alpha-1)(\alpha-2)}{\Gamma(\alpha)} \begin{cases} (t(1-s))^{\alpha-3} - (t-s)^{\alpha-3}, & 0 \leq s \leq t \leq 1, \\ (t(1-s))^{\alpha-3}, & 0 \leq t < s \leq 1. \end{cases} \quad (3.2)$$

We define the operator T by

$$Tu(t) = \int_0^1 G(t, s)a(s)f(s, u(s), u''(s))ds, \quad 0 \leq t \leq 1. \quad (3.3)$$

From (3.1) and (3.3), we have $Tu(0) = (Tu)'(0) = 0$. Moreover, we obtain by (H_1) , (H_2) and (3.3) that

$$(Tu)''(t) = \int_0^1 \frac{\partial^2}{\partial t^2} G(t, s)a(s)f(s, u(s), u''(s))ds \in C^2[0, 1], \quad \forall u \in X.$$

Thus, $T : X \rightarrow X$. By Lemma 2.3, $u(t)$ is a solution of the fractional boundary value problem (1.1)-(1.2) if and only if $u(t)$ is a fixed point of the operator T .

Define the cone $P \subset X$ by

$$P = \left\{ u \in X : u(t) \geq 0, \forall t \in [0, 1], \min_{\omega \leq t \leq 1} u(t) \geq \frac{1}{2} \omega^{\alpha-1} \|u\|_0 \right\},$$

where $0 < \omega < 1$ as in (H_2) .

Let the nonnegative continuous convex functionals γ, β and the nonnegative continuous concave functional ψ be defined on the cone P by

$$\gamma(u) = \max_{0 \leq t \leq 1} |u(t)|, \quad \beta(u) = \max_{0 \leq t \leq 1} |u''(t)|, \quad \psi(u) = \min_{\omega \leq t \leq 1} |u(t)|.$$

Then $\gamma, \beta, \psi : P \rightarrow [0, \infty)$ are three continuous nonnegative functionals such that $\|u\| = \max\{\gamma(u), \beta(u)\}$,

and (A_1) , (A_2) hold; γ, β are convex, ψ is concave and there holds $\psi(u) \leq \gamma(u)$, for all $u \in P$.

For $3 < \alpha \leq 4$, it is rather straightforward that

$$\max_{0 \leq t \leq 1} t^{\alpha-3}(1-t^{\alpha-2}) = \frac{\alpha-2}{2\alpha-5} \left(\frac{\alpha-3}{2\alpha-5} \right)^{\frac{\alpha-3}{\alpha-2}}, \quad \max_{0 \leq t \leq 1} t^{\alpha-1}(1-t^{\alpha-2}) = \frac{\alpha-2}{2\alpha-3} \left(\frac{\alpha-1}{2\alpha-3} \right)^{\frac{\alpha-1}{\alpha-2}}. \quad (3.4)$$

For convenience, we denote

$$M = \frac{(\alpha-1)(\alpha-2)}{\Gamma(\alpha)(2\alpha-5)} \left(\frac{\alpha-3}{2\alpha-5} \right)^{\frac{\alpha-3}{\alpha-2}}, \quad N = \frac{1}{\Gamma(\alpha)(2\alpha-3)} \left(\frac{\alpha-1}{2\alpha-3} \right)^{\frac{\alpha-1}{\alpha-2}},$$

$$m = \frac{\omega^{\alpha-1}}{2\Gamma(\alpha)} \int_{\omega}^1 \left[(1-s)^{\alpha-3} - (1-s)^{\alpha-1} \right] a(s) ds > 0, \text{ (by (H}_2\text{))}$$

where $3 < \alpha \leq 4$.

We are now in a position to present and prove our main result.

Theorem 3.2. Assume that (H_1) and (H_2) hold. Suppose there exist constants $r_2 \geq \frac{2b}{\omega^{\alpha-1}} > b > r_1 > 0$, $L_2 \geq L_1 > 0$ such that $\frac{b}{m} \leq \min\{\frac{r_2}{N\|a\|_0}, \frac{L_2}{M\|a\|_0}\}$. If the following assumptions hold

(H_3) $f(t, u, v) < \min\{\frac{r_1}{N\|a\|_0}, \frac{L_1}{M\|a\|_0}\}$, for $(t, u, v) \in [0, 1] \times [0, r_1] \times [-L_1, L_1]$;

(H_4) $f(t, u, v) > \frac{b}{m}$, for $(t, u, v) \in [\omega, 1] \times [b, \frac{2b}{\omega^{\alpha-1}}] \times [-L_2, L_2]$;

(H_5) $f(t, u, v) \leq \min\{\frac{r_2}{N\|a\|_0}, \frac{L_2}{M\|a\|_0}\}$, for $(t, u, v) \in [0, 1] \times [0, r_2] \times [-L_2, L_2]$,

then BVP (1.1)-(1.2) has at least three positive solutions u_1, u_2 , and u_3 such that

$$\begin{aligned} \max_{0 \leq t \leq 1} u_1(t) &\leq r_1, \quad \max_{0 \leq t \leq 1} |u_1''(t)| \leq L_1; \\ b &< \min_{\omega \leq t \leq 1} u_2(t) \leq \max_{0 \leq t \leq 1} u_2(t) \leq r_2, \quad \max_{0 \leq t \leq 1} |u_2''(t)| \leq L_2; \\ \max_{0 \leq t \leq 1} u_3(t) &\leq \frac{2b}{\omega^{\alpha-1}}, \quad \max_{0 \leq t \leq 1} |u_3''(t)| \leq L_2. \end{aligned}$$

Proof. By (H_1) , (H_2) , Lemma 2.4 and (3.3), for $u \in P$, we have $Tu(t) \geq 0$, $\forall t \in [0, 1]$, and

$$\begin{aligned} \min_{\omega \leq t \leq 1} Tu(t) &= \min_{\omega \leq t \leq 1} \int_0^1 G(t, s) a(s) f(s, u(s), u''(s)) ds \\ &\geq \int_0^1 \min_{\omega \leq t \leq 1} G(t, s) a(s) f(s, u(s), u''(s)) ds \\ &\geq \frac{1}{2} \omega^{\alpha-1} \int_0^1 h(s) a(s) f(s, u(s), u''(s)) ds \end{aligned}$$

$$\geq \frac{1}{2} \omega^{\alpha-1} \max_{0 \leq t \leq 1} \int_0^1 G(t, s) a(s) f(s, u(s), u''(s)) ds = \frac{1}{2} \omega^{\alpha-1} \|Tu\|_0, \quad t \in [0, 1].$$

Thus, $T(P) \subset P$. Moreover, it is easy to check by the Arzela-Ascoli theorem that the operator T is completely continuous. We now show that all the conditions of Lemma 2.1 are satisfied.

If $u \in \bar{P}(\gamma, r_2; \beta, L_2)$, then $\gamma(u) = \max_{0 \leq t \leq 1} |u(t)| \leq r_2$, $\beta(u) = \max_{0 \leq t \leq 1} |u''(t)| \leq L_2$, and assumption (H₅) implies

$$f(t, u(t), u''(t)) \leq \min \left\{ \frac{r_2}{N\|a\|_0}, \frac{L_2}{M\|a\|_0} \right\}, \quad \forall t \in [0, 1]. \quad (3.5)$$

Thus, by (3.4), (3.2) and (3.3), we get

$$\begin{aligned} \beta(Tu) &= \max_{0 \leq t \leq 1} |(Tu)''(t)| = \max_{0 \leq t \leq 1} \left| \int_0^1 \frac{\partial^2}{\partial t^2} G(t, s) a(s) f(s, u(s), u''(s)) ds \right| \\ &\leq \|a\|_0 \frac{L_2}{M\|a\|_0} \max_{0 \leq t \leq 1} \int_0^1 \frac{\partial^2}{\partial t^2} G(t, s) ds \\ &= \frac{L_2}{M} \frac{(\alpha-1)(\alpha-2)}{\Gamma(\alpha)} \max_{0 \leq t \leq 1} \left[\int_0^t (t^{\alpha-3}(1-s)^{\alpha-3} - (t-s)^{\alpha-3}) ds + \int_t^1 t^{\alpha-3}(1-s)^{\alpha-3} ds \right] \\ &= \frac{L_2}{M} \frac{\alpha-1}{\Gamma(\alpha)} \max_{0 \leq t \leq 1} t^{\alpha-3}(1-t^{\alpha-2}) = \frac{L_2}{M} \frac{\alpha-1}{\Gamma(\alpha)} \frac{\alpha-2}{2\alpha-5} \left(\frac{\alpha-3}{2\alpha-5} \right)^{\frac{\alpha-3}{\alpha-2}} \\ &= \frac{L_2}{M} \cdot M = L_2. \end{aligned}$$

Moreover, we obtain by (3.5) that

$$\begin{aligned} \gamma(Tu) &= \max_{0 \leq t \leq 1} \left| \int_0^1 G(t, s) a(s) f(s, u(s), u''(s)) ds \right| \\ &\leq \|a\|_0 \frac{r_2}{N\|a\|_0} \max_{0 \leq t \leq 1} \int_0^1 G(t, s) ds \\ &= \frac{r_2}{N} \frac{1}{\Gamma(\alpha)} \max_{0 \leq t \leq 1} \left[\int_0^t (t^{\alpha-1}(1-s)^{\alpha-3} - (t-s)^{\alpha-3}) ds + \int_t^1 t^{\alpha-1}(1-s)^{\alpha-3} ds \right] \\ &= \frac{r_2}{N} \frac{1}{\Gamma(\alpha)(\alpha-2)} \max_{0 \leq t \leq 1} t^{\alpha-1}(1-t^{\alpha-2}) = \frac{r_2}{N} \frac{1}{\Gamma(\alpha)} \frac{1}{2\alpha-3} \left(\frac{\alpha-1}{2\alpha-3} \right)^{\frac{\alpha-1}{\alpha-2}} \\ &= \frac{r_2}{N} \cdot N = r_2. \end{aligned}$$

Hence, $T : \bar{P}(\gamma, r_2; \beta, L_2) \rightarrow \bar{P}(\gamma, r_2; \beta, L_2)$. Similarly, if $u \in \bar{P}(\gamma, r_1; \beta, L_1)$, then assumption (H₃) yields $f(t, u(t), u''(t)) < \min \left\{ \frac{r_1}{N\|a\|_0}, \frac{L_1}{M\|a\|_0} \right\}$ for $t \in [0, 1]$. As in the argument above, we can obtain that $T : \bar{P}(\gamma, r_1; \beta, L_1) \rightarrow P(\gamma, r_1; \beta, L_1)$. Hence, condition (ii) of Lemma 2.1 is satisfied.

To check condition (i) of Lemma 2.1, we choose $u(t) = \frac{2b}{\omega^{\alpha-1}}$, $0 \leq t \leq 1$. It is easy to see that $u(t) = \frac{2b}{\omega^{\alpha-1}} \in \bar{P}(\gamma, \frac{2b}{\omega^{\alpha-1}}; \beta, L_2; \psi, b)$, $\psi(u) = \psi(\frac{2b}{\omega^{\alpha-1}}) > b$, and so $\{u \in \bar{P}(\gamma, \frac{2b}{\omega^{\alpha-1}}; \beta, L_2; \psi, b) \mid \psi(u) > b\} \neq \emptyset$. Hence, if $u \in \bar{P}(\gamma, \frac{2b}{\omega^{\alpha-1}}; \beta, L_2; \psi, b)$, then $b \leq u(t) \leq \frac{2b}{\omega^{\alpha-1}}$, $|u''(t)| \leq L_2$ for $\omega \leq t \leq 1$. From assumption (H₄), we have $f(t, u(t), u''(t)) > \frac{b}{m}$ for $\omega \leq t \leq 1$. Thus, by Lemma 2.4 and (3.2), we have

$$\begin{aligned} \psi(Tu) &= \min_{\omega \leq t \leq 1} \left| \int_0^1 G(t, s) a(s) f(s, u(s), u''(s)) ds \right| \\ &\geq \int_0^1 \min_{\omega \leq t \leq 1} G(t, s) a(s) f(s, u(s), u''(s)) ds \geq \frac{\omega^{\alpha-1}}{2} \int_0^1 h(s) a(s) f(s, u(s), u''(s)) ds \\ &\geq \frac{\omega^{\alpha-1}}{2} \int_\omega^1 h(s) a(s) f(s, u(s), u''(s)) ds > \frac{\omega^{\alpha-1}}{2} \frac{b}{m} \int_\omega^1 h(s) a(s) ds \\ &= \frac{\omega^{\alpha-1}}{2} \frac{b}{m} \frac{1}{\Gamma(\alpha)} \int_\omega^1 [(1-s)^{\alpha-3} - (1-s)^{\alpha-1}] a(s) ds \\ &= b, \end{aligned}$$

i.e.,

$$\psi(Tu) > b, \quad \forall u \in \bar{P}(\gamma, \frac{2b}{\omega^{\alpha-1}}; \beta, L_2, \psi, b).$$

This shows that condition (i) of Lemma 2.1 is satisfied. We finally show that (iii) of Lemma 2.1 also holds. Suppose that $u \in \bar{P}(\gamma, r_2; \beta, L_2; \psi, b)$ with $\gamma(Tu) > \frac{2b}{\omega^{\alpha-1}}$. Then, by the definition of ψ and $Tu \in P$, we have

$$\begin{aligned} \psi(Tu) &= \min_{\omega \leq t \leq 1} \left| \int_0^1 G(t, s) a(s) f(s, u(s), u''(s)) ds \right| \\ &\geq \frac{\omega^{\alpha-1}}{2} \int_0^1 h(s) a(s) f(s, u(s), u''(s)) ds \geq \frac{\omega^{\alpha-1}}{2} \max_{0 \leq t \leq 1} \int_0^1 G(t, s) a(s) f(s, u(s), u''(s)) ds \\ &= \frac{\omega^{\alpha-1}}{2} \gamma(Tu) > \frac{\omega^{\alpha-1}}{2} \cdot \frac{2b}{\omega^{\alpha-1}} = b. \end{aligned}$$

So, the condition (iii) of Lemma 2.1 is satisfied. Therefore, an application of Lemma 2.1 implies that the boundary value problem (1.1)-(1.2) has at least three positive solutions u_1, u_2 , and u_3 in $\bar{P}(\alpha, r_2; \beta, L_2)$ such that

$$\begin{aligned} \max_{0 \leq t \leq 1} u_1(t) &< r_1, \quad \max_{0 \leq t \leq 1} |u_1''(t)| < L_1; \\ b &\leq \min_{\omega \leq t \leq 1} u_2(t) \leq \max_{0 \leq t \leq 1} u_2(t) \leq r_2, \quad \max_{0 \leq t \leq 1} |u_2''(t)| \leq L_2; \\ \max_{0 \leq t \leq 1} u_3(t) &\leq \frac{2b}{\omega^{\alpha-1}}, \quad \max_{0 \leq t \leq 1} |u_3''(t)| \leq L_2. \end{aligned}$$

The proof is complete.

Finally, we give an example to illustrate the effectiveness of our result.

Example 3.1. Consider the nonlinear fractional boundary value problem :

$$D_{0+}^{3.6}u(t) + a(t)f(t, u(t), u''(t)) = 0, \quad 0 < t < 1, \quad (3.6)$$

$$u(0) = u'(0) = u''(0) = u''(1) = 0, \quad (3.7)$$

where $a(t) = 100(1 - t)$, and

$$f(t, u, v) = \begin{cases} \frac{1}{10^3}|cost| + \frac{1}{380}3u^3 + \left(\frac{|v|}{800}\right)^3, & u \leq 2, \\ \frac{1}{10^3}|cost| + \frac{6561}{380} + \left(\frac{|v|}{800}\right)^3, & u \geq 2. \end{cases}$$

Set $\alpha = 3.6$, $\omega = \frac{1}{2}$, we have

$$M = \frac{(\alpha - 1)(\alpha - 2)}{\Gamma(\alpha)(2\alpha - 5)} \left(\frac{\alpha - 3}{2\alpha - 5} \right)^{\frac{\alpha-3}{\alpha-2}} = 0.3125, \quad N = \frac{1}{\Gamma(\alpha)(2\alpha - 3)} \left(\frac{\alpha - 1}{2\alpha - 3} \right)^{\frac{\alpha-1}{\alpha-2}} = 0.8397,$$

$$\|a\|_0 = 100, \quad m = \frac{\omega^{\alpha-1}}{2\Gamma(\alpha)} \int_{\omega}^1 \left[(1-s)^{\alpha-3} - (1-s)^{\alpha-1} \right] a(s) ds = 0.12.$$

Obviously, $a(t)$ satisfies condition (H₂). Choose $r_1 = 1$, $b = 2$, $r_2 = 1500$, $L_1 = 30$, and $L_2 = 600$, then

$$\frac{2b}{\omega^{\alpha-1}} = 24.2515, \quad \frac{b}{m} = 16.6667,$$

$$\min \left\{ \frac{r_1}{N\|a\|_0}, \frac{L_1}{M\|a\|_0} \right\} = 0.0119, \quad \min \left\{ \frac{r_2}{N\|a\|_0}, \frac{L_2}{M\|a\|_0} \right\} = 17.8635.$$

Consequently, $f(t, u, v)$ satisfy

$$f(t, u, v) < 0.0119, \text{ for } (t, u, v) \in [0, 1] \times [0, 1] \times [-30, 30];$$

$$f(t, u, v) > 16.6667, \text{ for } (t, u, v) \in [1/2, 1] \times [2, 24.2515] \times [-600, 600];$$

$$f(t, u, v) < 17.8635, \text{ for } (t, u, v) \in [0, 1] \times [0, 1500] \times [-600, 600],$$

Then all the assumptions of Theorem 3.2 hold. Hence, with Theorem 3.2, nonlinear fractional boundary value problem (3.6), (3.7) has at least three positive solutions u_1, u_2, u_3 , such that

$$\max_{0 \leq t \leq 1} u_1(t) \leq 1, \quad \max_{0 \leq t \leq 1} |u_1''(t)| \leq 30;$$

$$2 < \min_{1/2 \leq t \leq 1} u_2(t) \leq \max_{0 \leq t \leq 1} u_2(t) \leq 1500, \quad \max_{0 \leq t \leq 1} |u_2''(t)| \leq 600;$$

$$\max_{0 \leq t \leq 1} u_3(t) \leq 24.2515, \quad \max_{0 \leq t \leq 1} |u_3''(t)| \leq 600.$$

Acknowledgements

The author would like to thank the reviewers for their helpful comments on this paper. This work is supported by the National Natural Science Foundation of China (10771212) and the Natural Science Foundation of Jiangsu Education Office (06KJB110010).

References

- [1] C. Bai and J. Fang, The existence of a positive solution for a singular coupled system of nonlinear fractional differential equations, *Appl. Math. Comput.* 150 (2004) 611-621.
- [2] C. Bai, Positive solutions for nonlinear fractional differential equations with coefficient that changes sign, *Nonlinear Analysis*, 64 (2006) 677-685.
- [3] Z. Bai and H. Lu, Positive solutions for boundary value problem of nonlinear fractional differential equation, *J. Math. Anal. Appl.* 311 (2005) 495-505.
- [4] Z. Bai and W. Ge, Existence of three positive solutions for some second-order boundary-value problems, *Comput. Math. Appl.* 48 (2004) 699-707.
- [5] M. El-Shahed, Positive solutions for boundary value problem of nonlinear fractional differential equation, *Abstract and Applied Analysis*, vol.2007, Article ID 10368, 8 pages, 2007, doi: 10.1155/2007/10368.
- [6] E. Kaufmann and E. Mboumi, Positive solutions of a boundary value problem for a nonlinear fractional differential equation, *Electron. J. Qual. Theory Differ. Equ.* 2008 (2008), No. 3, 1-11.
- [7] A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, vol. 204 of North-Holland Mathematics Studies, Elsevier Science, Amsterdam, The Netherlands, 2006.
- [8] A. A. Kilbas and J. J. Trujillo, Differential equations of fractional order: methods, results, and problems. I, *Appl. Anal.* 78 (2001) 153-192.
- [9] K. S. Miller and B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*, Wiley, New York, 1993.
- [10] I. Podlubny, *Fractional Differential Equations*, Mathematics in Sciences and Engineering 198, Academic Press, San Diego 1999.
- [11] S. Zhang, Existence of positive solution for a nonlinear fractional differential equation, *J. Math. Anal. Appl.* 278 (2003) 136-148.
- [12] S. Zhang, Positive solution for a boundary value problem of nonlinear fractional differential equations, *Electronic Journal of Differential Equations*, 36 (2006) 1-12.

(Received April 13, 2008)