

## EXISTENCE OF $\Psi$ -BOUNDED SOLUTIONS FOR LINEAR DIFFERENCE EQUATIONS ON $\mathbb{Z}$

Aurel Diamandescu  
University of Craiova

### Abstract

In this paper<sup>1</sup>, we give a necessary and sufficient condition for the existence of  $\Psi$ -bounded solutions for the nonhomogeneous linear difference equation  $x(n+1) = A(n)x(n) + f(n)$  on  $\mathbb{Z}$ . In addition, we give a result in connection with the asymptotic behavior of the  $\Psi$ -bounded solutions of this equation.

### 1. Introduction

The problem of boundedness of the solutions for the system of ordinary differential equations  $x' = A(t)x + f(t)$  was studied by Coppel in [2]. In [3], [4], [5], the author proposes a novel concept,  $\Psi$ -boundedness of solutions ( $\Psi$  being a matrix function), which is interesting and useful in some practical cases and presents the existence condition for such solutions. Also, in [1], the author associates this problem with the concept of  $\Psi$ -dichotomy on  $\mathbb{R}$  of the system  $x' = A(t)x$ .

Naturally, one wonders whether there are any similar concepts and results on the solutions of difference equations, which can be seen as the discrete version of differential equations.

In [7], the authors extend the concept of  $\Psi$ -boundedness to the solutions of difference equation

$$x(n+1) = A(n)x(n) + f(n) \quad (1)$$

(via  $\Psi$ -bounded sequence) and establish a necessary and sufficient condition for existence of  $\Psi$ -bounded solutions for the nonhomogeneous linear difference equation (1) in case  $f$  is a  $\Psi$ -summable sequence on  $\mathbb{N}$ .

In [6], the author proved a necessary and sufficient condition for the existence of  $\Psi$ -bounded solutions of (1) in case  $f$  is a  $\Psi$ -bounded sequence on  $\mathbb{N}$ .

Similarly, we can consider solutions of (1) which are bounded not only on  $\mathbb{N}$  but on the  $\mathbb{Z}$ .

In this case, the conditions for the existence of at least one  $\Psi$ -bounded solution are rather more complicated, as we will see below.

In this paper, we give a necessary and sufficient condition so that the nonhomogeneous linear difference equation (1) have at least one  $\Psi$ -bounded solution on  $\mathbb{Z}$  for every  $\Psi$ -summable function  $f$  on  $\mathbb{Z}$ .

Here,  $\Psi$  is a matrix function. The introduction of the matrix function  $\Psi$  permits to obtain a mixed asymptotic behavior of the components of the solutions.

---

<sup>1</sup>2000 Mathematics Subject Classification: 39A11, 39A10

Keywords: difference equations,  $\Psi$ -bounded solutions on  $\mathbb{Z}$ .

## 2. Preliminaries

Let  $\mathbb{R}^d$  be the Euclidean  $d$ -space. For  $x = (x_1, x_2, \dots, x_d)^T \in \mathbb{R}^d$ , let  $\|x\| = \max\{|x_1|, |x_2|, \dots, |x_d|\}$  be the norm of  $x$ . For a  $d \times d$  real matrix  $A = (a_{ij})$ , the norm  $|A|$  is defined by  $|A| = \sup_{\|x\| \leq 1} \|Ax\|$ . It is well-known that  $|A| = \max_{1 \leq i \leq d} \sum_{j=1}^d |a_{ij}|$ .

Let  $\Psi_i : \mathbb{Z} \rightarrow (0, \infty)$ ,  $i = 1, 2, \dots, d$  and let the matrix function

$$\Psi = \text{diag} [\Psi_1, \Psi_2, \dots, \Psi_d].$$

Then,  $\Psi(n)$  is invertible for each  $n \in \mathbb{Z}$ .

**Definition 1.** A function  $\varphi : \mathbb{Z} \rightarrow \mathbb{R}^d$  is called  $\Psi$ -bounded iff the function  $\Psi\varphi$  is bounded (i.e. there exists  $M > 0$  such that  $\|\Psi(n)\varphi(n)\| \leq M$  for all  $n \in \mathbb{Z}$ ).

**Definition 2.** A function  $\varphi : \mathbb{Z} \rightarrow \mathbb{R}^d$  is called  $\Psi$ -summable on  $\mathbb{Z}$  if  $\sum_{n=-\infty}^{\infty} \|\Psi(n)\varphi(n)\|$  is convergent (i.e.  $\lim_{\substack{p \rightarrow -\infty \\ q \rightarrow +\infty}} \sum_{n=p}^q \|\Psi(n)\varphi(n)\|$  is finite).

Consider the nonautonomous difference linear equation

$$y(n+1) = A(n)y(n) \tag{2}$$

where the  $d \times d$  real matrix  $A(n)$  is invertible at  $n \in \mathbb{Z}$ . Let  $Y$  be the fundamental matrix of (2) with  $Y(0) = I_d$  (identity  $d \times d$  matrix). It is well-known that

$$\text{i). } Y(n) = \begin{cases} A(n-1)A(n-2) \cdots A(1)A(0), & n > 0 \\ I_d, & n = 0 \\ [A(-1)A(-2) \cdots A(n)]^{-1}, & n < 0 \end{cases},$$

ii).  $Y(n+1) = A(n)Y(n)$  for all  $n \in \mathbb{Z}$

iii). the solution of (2) with the initial condition  $y(0) = y_0$  is

$$y(n) = Y(n)y_0, n \in \mathbb{Z};$$

iv).  $Y$  is invertible for each  $n \in \mathbb{Z}$  and

$$Y^{-1}(n) = \begin{cases} A^{-1}(0)A^{-1}(1) \cdots A^{-1}(n-1), & n > 0 \\ I_d, & n = 0 \\ A(-1)A(-2) \cdots A(n), & n < 0 \end{cases}$$

Let the vector space  $\mathbb{R}^d$  represented as a direct sum of three subspaces  $X_-$ ,  $X_0$ ,  $X_+$  such that a solution  $y$  of (2) is  $\Psi$ -bounded on  $\mathbb{Z}$  if and only if  $y(0) \in X_0$  and  $\Psi$ -bounded on  $\mathbb{Z}_+ = \{0, 1, 2, \dots\}$  if and only if  $y(0) \in X_- \oplus X_0$ . Also let  $P_-$ ,  $P_0$ ,  $P_+$  denote the corresponding projection of  $\mathbb{R}^d$  onto  $X_-$ ,  $X_0$ ,  $X_+$  respectively.

### 3. Main result

The main result of this paper is the following.

**Theorem 1.** The equation (1) has at least one  $\Psi$ - bounded solution on  $\mathbb{Z}$  for every  $\Psi$ - summable function  $f$  on  $\mathbb{Z}$  if and only if there is a positive constant  $K$  such that

$$\left\{ \begin{array}{ll} |\Psi(n)Y(n)P_-Y^{-1}(k+1)\Psi^{-1}(k)| & \leq K, \quad k+1 \leq \min\{0,n\} \\ |\Psi(n)Y(n)(P_0 + P_+)Y^{-1}(k+1)\Psi^{-1}(k)| & \leq K, \quad n < k+1 \leq 0 \\ |\Psi(n)Y(n)(P_0 + P_-)Y^{-1}(k+1)\Psi^{-1}(k)| & \leq K, \quad 0 < k+1 \leq n \\ |\Psi(n)Y(n)P_+Y^{-1}(k+1)\Psi^{-1}(k)| & \leq K, \quad k + 1 > \max\{0,n\} \end{array} \right. \quad (3)$$

**Proof.** First, we prove the "only if" part. We define the sets:

$$B_\Psi = \{x : \mathbb{Z} \longrightarrow \mathbb{R}^d \mid x \text{ is } \Psi\text{- bounded}\},$$

$$B = \{x : \mathbb{Z} \longrightarrow \mathbb{R}^d \mid x \text{ is } \Psi\text{- summable on } \mathbb{Z}\},$$

$$D = \{x : \mathbb{Z} \longrightarrow \mathbb{R}^d \mid x \in B_\Psi, x(0) \in X_- \oplus X_+, (x(n+1) - A(n)x(n)) \in B\}$$

Obviously,  $B_\Psi$ ,  $B$  and  $D$  are vector spaces over  $\mathbb{R}$  and the functionals

$$x \longmapsto \|x\|_{B_\Psi} = \sup_{n \in \mathbb{Z}} \|\Psi(n)x(n)\|,$$

$$x \longmapsto \|x\|_B = \sum_{n=-\infty}^{\infty} \|\Psi(n)x(n)\|,$$

$$x \longmapsto \|x\|_D = \|x\|_{B_\Psi} + \|x(n+1) - A(n)x(n)\|_B$$

are norms on  $B_\Psi$ ,  $B$  and  $D$  respectively.

**Step 1.** It is a simple exercise that  $(B_\Psi, \|\cdot\|_{B_\Psi})$  and  $(B, \|\cdot\|_B)$  are Banach spaces.

**Step 2.**  $(D, \|\cdot\|_D)$  is a Banach space.

Let  $(x_p)_{p \in \mathbb{N}}$  be a fundamental sequence in  $D$ . Then,  $(x_p)_{p \in \mathbb{N}}$  is a fundamental sequence in  $B_\Psi$ . Therefore, there exists a  $\Psi$ - bounded function  $x : \mathbb{Z} \longrightarrow \mathbb{R}^d$  such that  $\lim_{p \rightarrow \infty} \Psi(n)x_p(n) = \Psi(n)x(n)$ , uniformly on  $\mathbb{Z}$ . From

$$\|x_p(n) - x(n)\| \leq \|\Psi^{-1}(n)\| \|\Psi(n)(x_p(n) - x(n))\|,$$

it follows that the sequence  $(x_p)_{p \in \mathbb{N}}$  is almost uniformly convergent to function  $x$  on  $\mathbb{Z}$ . Because  $x_p(0) \in X_- \oplus X_+$ ,  $p \in \mathbb{N}$ , it follows that  $x(0) \in X_- \oplus X_+$ .

On the other hand, the sequence  $(f_p)_{p \in \mathbb{N}}$ ,  $f_p(n) = x_p(n+1) - A(n)x_p(n)$ ,  $n \in \mathbb{Z}$ , is a fundamental sequence in  $B$ . Therefore, there exists a function  $f \in B$  such that

$$\sum_{n=-\infty}^{\infty} \|\Psi(n)f_p(n) - \Psi(n)f(n)\| \longrightarrow 0 \text{ as } p \longrightarrow \infty.$$

It follows that  $\Psi(n)f_p(n) \longrightarrow \Psi(n)f(n)$  and  $f_p(n) \longrightarrow f(n)$  for each  $n \in \mathbb{Z}$ .

For a fixed but arbitrary  $n \in \mathbb{Z}$ ,  $n > 0$ , we have

$$\begin{aligned} x(n+1) - x(0) &= \lim_{p \rightarrow \infty} [x_p(n+1) - x_p(0)] = \\ &= \lim_{p \rightarrow \infty} \sum_{i=0}^n [x_p(i+1) - x_p(i)] = \\ &= \lim_{p \rightarrow \infty} \sum_{i=0}^n [x_p(i+1) - A(i)x_p(i) + A(i)x_p(i) - x_p(i)] = \\ &= \lim_{p \rightarrow \infty} \sum_{i=0}^n [f_p(i) - f(i) + f(i) + A(i)x_p(i) - x_p(i)] = \end{aligned}$$

$$\begin{aligned}
&= \sum_{i=0}^n [f(i) + A(i)x(i) - x(i)] = \\
&= \sum_{i=0}^{n-1} [f(i) + A(i)x(i) - x(i)] + f(n) + A(n)x(n) - x(n) = \\
&= x(n) - x(0) + f(n) + A(n)x(n) - x(n) = A(n)x(n) + f(n) - x(0).
\end{aligned}$$

Similarly, we have

$$x(1) - x(0) = A(0)x(0) + f(0) - x(0)$$

and, for  $n \in \mathbb{Z}$ ,  $n < 0$ ,

$$\begin{aligned}
x(n) - x(0) &= \lim_{p \rightarrow \infty} [x_p(n) - x_p(0)] = \lim_{p \rightarrow \infty} \sum_{i=n}^{-1} [x_p(i) - x_p(i+1)] = \\
&= \lim_{p \rightarrow \infty} \sum_{i=n}^{-1} [x_p(i) - A(i)x_p(i) + A(i)x_p(i) - x_p(i+1)] = \\
&= \lim_{p \rightarrow \infty} \sum_{i=n}^{-1} [x_p(i) - A(i)x_p(i) - f_p(i)] = \\
&= \sum_{i=n}^{-1} [x(i) - A(i)x(i) - f(i)] = \\
&= \sum_{i=n+1}^{-1} [x(i) - A(i)x(i) - f(i)] + x(n) - A(n)x(n) - f(n) = \\
&= x(n+1) - x(0) + x(n) - A(n)x(n) - f(n).
\end{aligned}$$

By the above relations, we have that

$$x(n+1) - A(n)x(n) = f(n), \quad n \in \mathbb{Z}.$$

It follows that  $x \in D$ .

Now, from the relations

$$\begin{aligned}
&\sum_{n=-\infty}^{\infty} \|\Psi(n)(x_p - x)(n+1) - \Psi(n)A(n)(x_p - x)(n)\| \longrightarrow 0 \text{ as } p \longrightarrow \infty, \\
&\|x_p - x\|_{B_\Psi} \longrightarrow 0 \text{ as } p \longrightarrow \infty,
\end{aligned}$$

it follows that  $\|x_p - x\|_D \longrightarrow 0$  as  $p \longrightarrow +\infty$ .

Thus,  $(D, \|\cdot\|_D)$  is a Banach space.

**Step 3.** There exists a positive constant  $K$  such that, for every  $f \in B$  and for corresponding solution  $x \in D$  of (1), we have

$$\|x\|_{B_\Psi} \leq K \cdot \|f\|_B. \quad (4)$$

We define the operator  $T : D \longrightarrow B$ ,  $(Tx)(n) = x(n+1) - A(n)x(n)$ ,  $n \in \mathbb{Z}$ .

Clearly,  $T$  is linear and bounded, with  $\|T\| \leq 1$ . Let  $Tx = 0$  be. Then,  $x \in D$  and  $x(n+1) = A(n)x(n)$ . This shows that  $x$  is a  $\Psi$ -bounded solution of (2) with  $x(0) \in X_- \oplus X_+$ . From the Definition of  $X_0$ , we have  $x(0) \in X_0$ . Thus,  $x(0) \in X_0 \cap (X_- \oplus X_+) = \{0\}$ . It follows that  $x = 0$ . This means that the operator  $T$  is one-to-one.

Now, for  $f \in B$ , let  $x$  be a  $\Psi$ -bounded solution of the equation (1). Let  $z$  be the solution of the Cauchy problem

$$z(n+1) = A(n)z(n) + f(n), \quad z(0) = (P_- + P_+)x(0).$$

Then, the function  $u = x - z$  is a solution of the equation (2) with

$$u(0) = x(0) - z(0) = P_0x(0) \in X_0.$$

It follows that the function  $u$  is  $\Psi$ -bounded on  $\mathbb{Z}$ . Thus, the function  $z$  is  $\Psi$ -bounded on  $\mathbb{Z}$ . It follows that  $z \in D$  and  $Tz = f$ . Consequently,  $T$  is onto.

From a fundamental result of Banach "If  $T$  is a bounded one-to-one linear operator from a Banach space onto another, then the inverse operator  $T^{-1}$  is also bounded", we have that

$$\|T^{-1}f\|_D \leq \|T^{-1}\| \|f\|_B, \text{ for } f \in B.$$

Denoting  $T^{-1}f = x$ , we have  $\|x\|_D = \|x\|_{B_\Psi} + \|f\|_B \leq \|T^{-1}\| \|f\|_B$  and then

$$\|x\|_{B_\Psi} \leq (\|T^{-1}\| - 1) \|f\|_B.$$

Thus, we have (4), where  $K = \|T^{-1}\| - 1$ .

**Step 4.** The end of the proof.

For a fixed but arbitrary  $k \in \mathbb{Z}$ ,  $\xi \in \mathbb{R}^d$ , we consider the function  $f: \mathbb{Z} \rightarrow \mathbb{R}^d$  defined by

$$f(n) = \begin{cases} \Psi^{-1}(n)\xi, & \text{if } n = k \\ 0, & \text{elsewhere} \end{cases}.$$

Obviously,  $f \in B$  and  $\|f\|_B = \|\xi\|$ . The corresponding solution  $x \in D$  of (1) is  $x(n) = G(n, k+1)f(k)$ , where

$$G(n, k) = \begin{cases} Y(n)P_-Y^{-1}(k) & k \leq \min\{0, n\} \\ -Y(n)(P_0 + P_+)Y^{-1}(k) & n < k \leq 0 \\ Y(n)(P_0 + P_-)Y^{-1}(k) & 0 < k \leq n \\ -Y(n)P_+Y^{-1}(k) & k > \max\{0, n\} \end{cases}.$$

Indeed, we prove this in more cases:

Case  $k \leq -1$ . a). for  $k + 1 \leq n \leq 0$ ,

$$\begin{aligned} x(n+1) &= G(n+1, k+1)f(k) = Y(n+1)P_-Y^{-1}(k+1)f(k) = \\ &= A(n)Y(n)P_-Y^{-1}(k+1)f(k) = A(n)x(n) = A(n)x(n) + f(n) \text{ (because } f(n) = 0); \end{aligned}$$

b). for  $n = k$ ,

$$\begin{aligned} x(n+1) &= G(n+1, k+1)f(k) = Y(n+1)P_-Y^{-1}(k+1)f(k) = \\ &= Y(k+1)(I - P_0 - P_+) \cdot Y^{-1}(k+1)f(k) = f(k) - A(k)Y(k)(P_0 + P_+)Y^{-1}(k+1)f(k) = \\ &= f(k) + A(k)G(k, k+1)f(k) = A(n)x(n) + f(n); \end{aligned}$$

c). for  $n < k$ ,

$$\begin{aligned} x(n+1) &= G(n+1, k+1)f(k) = -Y(n+1)(P_0 + P_+)Y^{-1}(k+1)f(k) = \\ &= -A(n)Y(n)(P_0 + P_+)Y^{-1}(k+1)f(k) = A(n)x(n) = A(n)x(n) + f(n); \end{aligned}$$

d). for  $n > 0$ ,

$$\begin{aligned} x(n+1) &= G(n+1, k+1)f(k) = Y(n+1)P_-Y^{-1}(k+1)f(k) = \\ &= A(n)Y(n)P_-Y^{-1}(k+1)f(k) = A(n)x(n) = A(n)x(n) + f(n); \end{aligned}$$

Case  $k > -1$ .  $\alpha$ ). for  $n < 0$ ,

$$\begin{aligned} x(n+1) &= G(n+1, k+1)f(k) = -Y(n+1)P_+Y^{-1}(k+1)f(k) = \\ &= -A(n)Y(n)P_+Y^{-1}(k+1)f(k) = A(n)x(n) = A(n)x(n) + f(n); \end{aligned}$$

$\beta$ ). for  $n = 0$  and  $k = 0$ ,

$$\begin{aligned} x(1) &= G(1, 1)f(0) = Y(1)(P_0 + P_-)Y^{-1}(1)f(0) = Y(1)(I - P_+)Y^{-1}(1)f(0) = \\ &= f(0) - A(0)Y(0)P_+Y^{-1}(1)f(0) = A(0)x(0) + f(0); \end{aligned}$$

$\gamma$ ).  $n = 0$  and  $k > 0$ ,

$$\begin{aligned} x(1) &= G(1, k+1)f(k) = -Y(1)P_+Y^{-1}(k+1)f(k) = -A(0)Y(0)P_+Y^{-1}(k+1)f(k) = \\ &= A(0)G(0, k+1)f(k) = A(0)x(0) + f(0); \end{aligned}$$

$\delta$ ). for  $0 < n = k$ ,

$$\begin{aligned} x(n+1) &= G(k+1, k+1)f(k) = Y(k+1)(P_0 + P_-)Y^{-1}(k+1)f(k) = \\ &= Y(k+1)(I - P_+)Y^{-1}(k+1)f(k) = f(k) - A(k)Y(k)P_+Y^{-1}(k+1)f(k) = \end{aligned}$$

$$= A(n)x(n) + f(n);$$

$\varepsilon$ ). for  $0 < n < k$ ,

$$\begin{aligned} x(n+1) &= G(n+1, k+1)f(k) = -Y(n+1)P_+Y^{-1}(k+1)f(k) = \\ &= -A(n)Y(n)P_+Y^{-1}(k+1)f(k) = A(n)x(n) = A(n)x(n) + f(n); \end{aligned}$$

$\zeta$ ). for  $n \geq k + 1$ ,

$$\begin{aligned} x(n+1) &= G(n+1, k+1)f(k) = Y(n+1)(P_0 + P_-)Y^{-1}(k+1)f(k) = \\ &= A(n)Y(n)(P_0 + P_-)Y^{-1}(k+1)f(k) = A(n)x(n) = A(n)x(n) + f(n). \end{aligned}$$

On the other hand,  $x(0) \in X_- \oplus X_+$ , because

$$x(0) = G(0, k+1)f(k) = \begin{cases} +P_-Y^{-1}(k+1)f(k), & k+1 \leq 0 \\ -P_+Y^{-1}(k+1)f(k), & k+1 > 0 \end{cases}.$$

Finally, we have

$$x(n) = G(n, k+1)f(k) = \begin{cases} -Y(n)(P_0 + P_+)Y^{-1}(k+1)f(k), & n < k+1 \leq 0 \\ Y(n)(P_0 + P_-)Y^{-1}(k+1)f(k), & n \geq k+1 \geq 0 \end{cases}.$$

From the Definitions of  $X_-$ ,  $X_0$  and  $X_+$ , it follows that the function  $x$  is  $\Psi$ -bounded on  $\mathbb{Z}_-$  and  $\mathbb{N}$ . Thus,  $x$  is the solution of (1) in  $D$ .

Now, we have,  $\|\Psi(n)x(n)\| = \|\Psi(n)G(n, k+1)f(k)\| = \|\Psi(n)G(n, k+1)\Psi^{-1}(k)\xi\|$ .

The inequality (4) becomes

$$\|\Psi(n)G(n, k+1)\Psi^{-1}(k)\xi\| \leq K\|\xi\|, \text{ for all } k, n \in \mathbb{Z}, \xi \in \mathbb{R}^d.$$

It follows that  $\|\Psi(n)G(n, k+1)\Psi^{-1}(k)\| \leq K$ , for all  $k, n \in \mathbb{Z}$ , which is equivalent with (3).

Now, we prove the "if" part.

For a given  $\Psi$ -summable function  $f : \mathbb{Z} \rightarrow \mathbb{R}^d$ , consider  $u : \mathbb{Z} \rightarrow \mathbb{R}^d$  defined by

$$u(n) = \begin{cases} \sum_{k=-\infty}^{n-1} Y(n)P_-Y^{-1}(k+1)f(k) - \sum_{k=n}^{-1} Y(n)P_0Y^{-1}(k+1)f(k) - \\ \quad - \sum_{k=n}^{\infty} Y(n)P_+Y^{-1}(k+1)f(k), & n < 0 \\ \sum_{k=-\infty}^{-1} Y(0)P_-Y^{-1}(k+1)f(k) - \sum_{k=0}^{\infty} Y(0)P_+Y^{-1}(k+1)f(k), & n = 0 \\ \sum_{k=-\infty}^{n-1} Y(n)P_-Y^{-1}(k+1)f(k) + \sum_{k=0}^{n-1} Y(n)P_0Y^{-1}(k+1)f(k) - \\ \quad - \sum_{k=n}^{\infty} Y(n)P_+Y^{-1}(k+1)f(k), & n > 0 \end{cases}$$

**Step 5.** The function  $u$  is well-defined.

For  $p, q \in \mathbb{Z}$ ,  $q < 0 < p$ , we have

$$\begin{aligned} &\sum_{k=q}^{-1} \|Y(0)P_-Y^{-1}(k+1)f(k)\| + \sum_{k=0}^p \|Y(0)P_+Y^{-1}(k+1)f(k)\| \leq \\ &\leq |\Psi^{-1}(0)| \sum_{k=q}^{-1} \|\Psi(0)Y(0)P_-Y^{-1}(k+1)\Psi^{-1}(k)\| \|\Psi(k)f(k)\| + \\ &+ |\Psi^{-1}(0)| \sum_{k=0}^p \|\Psi(0)Y(0)P_+Y^{-1}(k+1)\Psi^{-1}(k)\| \|\Psi(k)f(k)\| \leq \\ &\leq K|\Psi^{-1}(0)| \left( \sum_{k=q}^p \|\Psi(k)f(k)\| \right), \end{aligned}$$

and then,  $\sum_{k=-\infty}^{-1} Y(0)P_-Y^{-1}(k+1)f(k)$  and  $\sum_{k=0}^{\infty} Y(0)P_+Y^{-1}(k+1)f(k)$  are absolutely convergent series. Thus,  $u(0)$  is well-defined.

For  $m, n \in \mathbb{Z}$ ,  $m \geq n > 0$ , we have

$$\begin{aligned} & \sum_{k=n}^m \|Y(n)P_+Y^{-1}(k+1)f(k)\| = \\ &= \sum_{k=n}^m \|\Psi^{-1}(n)(\Psi(n)Y(n)P_+Y^{-1}(k+1)\Psi^{-1}(k))(\Psi(k)f(k))\| \leq \\ &\leq |\Psi^{-1}(n)| \sum_{k=n}^m |\Psi(n)Y(n)P_+Y^{-1}(k+1)\Psi^{-1}(k)| \|\Psi(k)f(k)\| \leq \\ &\leq K|\Psi^{-1}(n)| \left( \sum_{k=n}^m \|\Psi(k)f(k)\| \right), \end{aligned}$$

and then,  $\sum_{k=n}^{\infty} Y(n)P_+Y^{-1}(k+1)f(k)$  is an absolutely convergent series for  $n > 0$ .

For  $m \in \mathbb{Z}$ ,  $n \in \mathbb{N}$ ,  $m < n - 1$ , we have

$$\begin{aligned} & \sum_{k=m}^{n-1} \|Y(n)P_-Y^{-1}(k+1)f(k)\| = \\ &= \sum_{k=m}^{n-1} \|\Psi^{-1}(n)(\Psi(n)Y(n)P_-Y^{-1}(k+1)\Psi^{-1}(k))(\Psi(k)f(k))\| \\ &\leq |\Psi^{-1}(n)| \sum_{k=m}^{n-1} |\Psi(n)Y(n)P_-Y^{-1}(k+1)\Psi^{-1}(k)| \|\Psi(k)f(k)\| \leq \\ &\leq K|\Psi^{-1}(n)| \sum_{k=m}^{n-1} \|\Psi(k)f(k)\|, \end{aligned}$$

and then,  $\sum_{k=-\infty}^{n-1} Y(n)P_-Y^{-1}(k+1)f(k)$  is an absolutely convergent series for  $n > 0$ .

Thus, the function  $u$  is well defined for  $n \geq 0$ .

Similarly, the function  $u$  is well defined for  $n < 0$ .

**Step 6.** The function  $u$  is a solution of the equation (1).

Indeed, using the expression of the function  $u$ , we obtain:

$$\begin{aligned} \bullet \quad & u(1) = \sum_{k=-\infty}^0 Y(1)P_-Y^{-1}(k+1)f(k) + Y(1)P_0Y^{-1}(1)f(0) - \\ & - \sum_{k=1}^{\infty} Y(1)P_+Y^{-1}(k+1)f(k) = A(0) \left[ \sum_{k=-\infty}^0 Y(0)P_-Y^{-1}(k+1)f(k) + \right. \\ & + Y(0)P_0Y^{-1}(1)f(0) - \left. \sum_{k=1}^{\infty} Y(0)P_+Y^{-1}(k+1)f(k) \right] = \\ &= A(0) \left[ \sum_{k=-\infty}^{-1} Y(0)P_-Y^{-1}(k+1)f(k) + Y(0)P_-Y^{-1}(1)f(0) + Y(0)P_0Y^{-1}(1)f(0) \right. \\ & - \left. \sum_{k=0}^{\infty} Y(0)P_+Y^{-1}(k+1)f(k) + Y(0)P_+Y^{-1}(1)f(0) \right] = \\ &= A(0)u(0) + A(0)Y(0)(P_- + P_0 + P_+)Y^{-1}(1)f(0) = A(0)u(0) + f(0); \\ \bullet \quad & \text{for } n > 0, u(n+1) = \sum_{k=-\infty}^n Y(n+1)P_-Y^{-1}(k+1)f(k) + \\ & + \sum_{k=0}^n Y(n+1)P_0Y^{-1}(k+1)f(k) - \sum_{k=n+1}^{\infty} Y(n+1)P_+Y^{-1}(k+1)f(k) = \end{aligned}$$

$$\begin{aligned}
&= A(n) \left[ \sum_{k=-\infty}^{n-1} Y(n)P_-Y^{-1}(k+1)f(k) + Y(n)P_-Y^{-1}(n+1)f(n) + \right. \\
&+ \sum_{k=0}^{n-1} Y(n)P_0Y^{-1}(k+1)f(k) + Y(n)P_0Y^{-1}(n+1)f(n) - \\
&- \sum_{k=n}^{\infty} Y(n)P_+Y^{-1}(k+1)f(k) + Y(n)P_+Y^{-1}(n+1)f(n) \left. \right] = \\
&= A(n)u(n) + Y(n+1)(P_- + P_0 + P_+)Y^{-1}(n+1)f(n) = A(n)u(n) + f(n); \\
\bullet \quad &u(0) = \sum_{k=-\infty}^{-1} Y(0)P_-Y^{-1}(k+1)f(k) - \sum_{k=0}^{\infty} Y(0)P_+Y^{-1}(k+1)f(k) = \\
&= A(-1) \left[ \sum_{k=-\infty}^{-1} Y(-1)P_-Y^{-1}(k+1)f(k) - \sum_{k=0}^{\infty} Y(-1)P_+Y^{-1}(k+1)f(k) \right] = \\
&= A(-1) \left[ \sum_{k=-\infty}^{-2} Y(-1)P_-Y^{-1}(k+1)f(k) + Y(-1)P_-Y^{-1}(0)f(-1) - \right. \\
&- \sum_{k=-1}^{-1} Y(-1)P_0Y^{-1}(k+1)f(k) + Y(-1)P_0Y^{-1}(0)f(-1) - \\
&- \sum_{k=-1}^{\infty} Y(-1)P_+Y^{-1}(k+1)f(k) + Y(-1)P_+Y^{-1}(0)f(-1) \left. \right] = \\
&= A(-1)u(-1) + Y(0)(P_- + P_0 + P_+)Y^{-1}(0)f(-1) = \\
&= A(-1)u(-1) + f(-1); \\
\bullet \quad &\text{for } n < -1, u(n+1) = \sum_{k=-\infty}^n Y(n+1)P_-Y^{-1}(k+1)f(k) - \\
&- \sum_{k=n+1}^{-1} Y(n+1)P_0Y^{-1}(k+1)f(k) - \sum_{k=n+1}^{\infty} Y(n+1)P_+Y^{-1}(k+1)f(k) = \\
&= A(n) \left[ \sum_{k=-\infty}^{n-1} Y(n)P_-Y^{-1}(k+1)f(k) + Y(n)P_-Y^{-1}(n+1)f(n) - \right. \\
&- \sum_{k=n}^{-1} Y(n)P_0Y^{-1}(k+1)f(k) + Y(n)P_0Y^{-1}(n+1)f(n) - \\
&- \sum_{k=n}^{\infty} Y(n)P_+Y^{-1}(k+1)f(k) + Y(n)P_+Y^{-1}(n+1)f(n) \left. \right] = \\
&= A(n)u(n) + Y(n+1)(P_- + P_0 + P_+)Y^{-1}(n+1)f(n) = A(n)u(n) + f(n).
\end{aligned}$$

These relations show that the function  $u$  is a solution of the equation (1).

**Step 7.** The function  $u$  is  $\Psi$ -bounded on  $\mathbb{Z}$ .

Indeed, for  $n > 0$  we have

$$\begin{aligned}
\| \Psi(n)u(n) \| &= \left\| \sum_{k=-\infty}^{n-1} \Psi(n)Y(n)P_-Y^{-1}(k+1)\Psi^{-1}(k)\Psi(k)f(k) + \right. \\
&+ \sum_{k=0}^{n-1} \Psi(n)Y(n)P_0Y^{-1}(k+1)\Psi^{-1}(k)\Psi(k)f(k) - \\
&- \sum_{k=n}^{\infty} \Psi(n)Y(n)P_+Y^{-1}(k+1)\Psi^{-1}(k)\Psi(k)f(k) \left. \right\| = \\
&= \left\| \sum_{k=-\infty}^{-1} \Psi(n)Y(n)P_-Y^{-1}(k+1)\Psi^{-1}(k)\Psi(k)f(k) + \right. \\
&+ \sum_{k=0}^{n-1} \Psi(n)Y(n)(P_0 + P_-)Y^{-1}(k+1)\Psi^{-1}(k)\Psi(k)f(k) -
\end{aligned}$$



$$\begin{aligned}
& - \sum_{k=n}^{\infty} \Psi(n)Y(n)P_+Y^{-1}(k+1)\Psi^{-1}(k)\Psi(k)f(k) \leq \\
& \leq \sum_{k=-\infty}^{-1} |\Psi(n)Y(n)P_-Y^{-1}(k+1)\Psi^{-1}(k)| \|\Psi(k)f(k)\| + \\
& + \sum_{k=0}^{n-1} |\Psi(n)Y(n)(P_0 + P_-)Y^{-1}(k+1)\Psi^{-1}(k)| \|\Psi(k)f(k)\| + \\
& + \sum_{k=n}^{\infty} |\Psi(n)Y(n)P_+Y^{-1}(k+1)\Psi^{-1}(k)| \|\Psi(k)f(k)\| \leq \\
& \leq K \left( \sum_{k=-\infty}^{-1} \|\Psi(k)f(k)\| + \sum_{k=0}^{n-1} \|\Psi(k)f(k)\| + \sum_{k=n}^{\infty} \|\Psi(k)f(k)\| \right) = \\
& = K \sum_{k=-\infty}^{+\infty} \|\Psi(k)f(k)\| = K\|f\|_B.
\end{aligned}$$

For  $n < 0$ , we have

$$\begin{aligned}
\|\Psi(n)u(n)\| &= \left\| \sum_{k=-\infty}^{n-1} \Psi(n)Y(n)P_-Y^{-1}(k+1)\Psi^{-1}(k)\Psi(k)f(k) - \right. \\
& - \sum_{k=n}^{-1} \Psi(n)Y(n)P_0Y^{-1}(k+1)\Psi^{-1}(k)\Psi(k)f(k) - \\
& - \left. \sum_{k=n}^{\infty} \Psi(n)Y(n)P_+Y^{-1}(k+1)\Psi^{-1}(k)\Psi(k)f(k) \right\| = \\
& = \left\| \sum_{k=-\infty}^{n-1} \Psi(n)Y(n)P_-Y^{-1}(k+1)\Psi^{-1}(k)\Psi(k)f(k) - \right. \\
& - \sum_{k=n}^{-1} \Psi(n)Y(n)(P_0 + P_+)Y^{-1}(k+1)\Psi^{-1}(k)\Psi(k)f(k) - \\
& - \left. \sum_{k=0}^{\infty} \Psi(n)Y(n)P_+Y^{-1}(k+1)\Psi^{-1}(k)\Psi(k)f(k) \right\| \leq \\
& \leq \sum_{k=-\infty}^{n-1} |\Psi(n)Y(n)P_-Y^{-1}(k+1)\Psi^{-1}(k)| \|\Psi(k)f(k)\| + \\
& + \sum_{k=n}^{-1} |\Psi(n)Y(n)(P_0 + P_+)Y^{-1}(k+1)\Psi^{-1}(k)| \|\Psi(k)f(k)\| + \\
& + \sum_{k=0}^{\infty} |\Psi(n)Y(n)P_+Y^{-1}(k+1)\Psi^{-1}(k)| \|\Psi(k)f(k)\| \leq \\
& \leq K \left( \sum_{k=-\infty}^{n-1} \|\Psi(k)f(k)\| + \sum_{k=n}^{-1} \|\Psi(k)f(k)\| + \sum_{k=0}^{\infty} \|\Psi(k)f(k)\| \right) = \\
& = K \sum_{k=-\infty}^{+\infty} \|\Psi(k)f(k)\| = K\|f\|_B.
\end{aligned}$$

Similarly,  $\|\Psi(0)u(0)\| \leq K\|f\|_B$ .

Therefore,  $\|\Psi(n)u(n)\| \leq K\|f\|_B$ , for all  $n \in \mathbb{Z}$ .

Thus, the solution  $u$  of the equation (1) is  $\Psi$ -bounded on  $\mathbb{Z}$ .

The proof is now complete.

**Corollary 1.** If the homogeneous equation (2) has no nontrivial  $\Psi$ -bounded solution on  $\mathbb{Z}$ , then, the equation (1) has a unique  $\Psi$ -bounded solution on  $\mathbb{Z}$  for every  $\Psi$ -summable function  $f$  on  $\mathbb{Z}$  if and only if there exists a positive constant  $K$  such that, for  $k, n \in \mathbb{Z}$ ,

$$\begin{cases} |\Psi(n)Y(n)P_-Y^{-1}(k+1)\Psi^{-1}(k)| \leq K, & \text{for } k+1 \leq n \\ |\Psi(n)Y(n)P_+Y^{-1}(k+1)\Psi^{-1}(k)| \leq K, & \text{for } n < k+1 \end{cases} \quad (5)$$

**Proof.** Indeed, in this case,  $P_0 = 0$ . Now, the Corollary follows from the above Theorem.

Finally, we give a result in which we will see that the asymptotic behavior of  $\Psi$ - bounded solutions of (1) is determined completely by the asymptotic behavior of the fundamental matrix  $Y$  of (2).

**Theorem 2.** Suppose that:

1°. the fundamental matrix  $Y$  of (2) satisfies the conditions (3) for some  $K > 0$  and the conditions

- i).  $\lim_{n \rightarrow \pm\infty} |\Psi(n)Y(n)P_0| = 0$ ;
- ii).  $\lim_{n \rightarrow +\infty} |\Psi(n)Y(n)P_-| = 0$ ;
- iii).  $\lim_{n \rightarrow -\infty} |\Psi(n)Y(n)P_+| = 0$ .

2°. the function  $f: \mathbb{Z} \rightarrow \mathbb{R}^d$  is  $\Psi$ - summable on  $\mathbb{Z}$ .

Then, every  $\Psi$ - bounded solution  $x$  of (1) satisfies the condition

$$\lim_{n \rightarrow \pm\infty} \|\Psi(n)x(n)\| = 0.$$

**Proof.** Let  $x$  be a  $\Psi$ - bounded solution of (1). Let  $u$  be the  $\Psi$ - bounded solution of (1) from the proof of Theorem 1 ("if" part).

Let the function  $y(n) = x(n) - u(n) - Y(n)P_0(x(0) - u(0))$ ,  $n \in \mathbb{Z}$ .

It is easy to see that  $y$  is a  $\Psi$ - bounded solution of (2) and then  $y(0) \in X_0$ .

On the other hand,

$$y(0) = (I - P_0)(x(0) - u(0)) = (P_- + P_+)(x(0) - u(0)) \in X_- \oplus X_+.$$

Thus,  $y(0) \in (X_- \oplus X_+) \cap X_0 = \{0\}$ . It follows that  $y = 0$  and then

$$x(n) = u(n) + Y(n)P_0(x(0) - u(0)), n \in \mathbb{Z}.$$

Now, we prove that  $\lim_{n \rightarrow \pm\infty} \|\Psi(n)x(n)\| = 0$ .

For  $n > 0$ , we have

$$\begin{aligned} x(n) &= Y(n)P_0(x(0) - u(0)) + \sum_{k=-\infty}^{n-1} Y(n)P_-Y^{-1}(k+1)f(k) + \\ &+ \sum_{k=0}^{n-1} Y(n)P_0Y^{-1}(k+1)f(k) - \sum_{k=n}^{\infty} Y(n)P_+Y^{-1}(k+1)f(k) \end{aligned}$$

and then

$$\begin{aligned} \Psi(n)x(n) &= \Psi(n)Y(n)P_0(x(0) - u(0)) + \\ &+ \sum_{k=-\infty}^{-1} \Psi(n)Y(n)P_-Y^{-1}(k+1)\Psi^{-1}(k)\Psi(k)f(k) + \\ &+ \sum_{k=0}^{n-1} \Psi(n)Y(n)(P_0 + P_-)Y^{-1}(k+1)\Psi^{-1}(k)\Psi(k)f(k) - \\ &- \sum_{k=n}^{\infty} \Psi(n)Y(n)P_+Y^{-1}(k+1)\Psi^{-1}(k)\Psi(k)f(k). \end{aligned}$$

By the hypotheses, for a given  $\varepsilon > 0$ , there exist:

- $n_1 \in \mathbb{N}$  such that, for  $n \geq n_1$ ,  

$$\sum_{k=-\infty}^{-n} \|\Psi(k)f(k)\| < \frac{\varepsilon}{5K} \text{ and } \sum_{k=n}^{\infty} \|\Psi(k)f(k)\| < \frac{\varepsilon}{5K};$$
- $n_2 \in \mathbb{N}$ ,  $n_2 > n_1$ , such that, for  $n \geq n_2$ ,  

$$|\Psi(n)Y(n)P_-| < \frac{\varepsilon}{5} \left( 1 + \sum_{k=-n_1+1}^{-1} \|Y^{-1}(k+1)f(k)\| \right)^{-1};$$
- $n_3 \in \mathbb{N}$ ,  $n_3 > n_2$ , such that, for  $n \geq n_3$ ,  

$$|\Psi(n)Y(n)P_0| < \frac{\varepsilon}{5} (1 + \|x(0) - u(0)\|)^{-1};$$
- $n_4 \in \mathbb{N}$ ,  $n_4 > n_3$ , such that, for  $n \geq n_4$ ,  

$$|\Psi(n)Y(n)(P_0+P_-)| < \frac{\varepsilon}{5} \left( 1 + \sum_{k=0}^{n_1} \|Y^{-1}(k+1)f(k)\| \right)^{-1}.$$

Then, for  $n \geq n_4$  we have

$$\begin{aligned} \|\Psi(n)x(n)\| &\leq |\Psi(n)Y(n)P_0| \|x(0) - u(0)\| + \\ &+ \sum_{k=-\infty}^{-n_1} |\Psi(n)Y(n)P_- Y^{-1}(k+1)\Psi^{-1}(k)| \|\Psi(k)f(k)\| + \\ &+ \sum_{k=-n_1+1}^{-1} |\Psi(n)Y(n)P_-| \|Y^{-1}(k+1)f(k)\| + \\ &+ \sum_{k=0}^{n_1} |\Psi(n)Y(n)(P_0 + P_-)| \|Y^{-1}(k+1)f(k)\| + \\ &+ \sum_{k=n_1+1}^{n-1} |\Psi(n)Y(n)(P_0 + P_-)Y^{-1}(k+1)\Psi^{-1}(k)| \|\Psi(k)f(k)\| + \\ &+ \sum_{k=n}^{\infty} |\Psi(n)Y(n)P_+ Y^{-1}(k+1)\Psi^{-1}(k)| \|\Psi(k)f(k)\| < \\ &< \frac{\varepsilon}{5} (1 + \|x(0) - u(0)\|)^{-1} \|x(0) - u(0)\| + \\ &+ K \sum_{k=-\infty}^{-n_1} \|\Psi(k)f(k)\| + |\Psi(n)Y(n)P_-| \sum_{k=-n_1+1}^{-1} \|Y^{-1}(k+1)f(k)\| + \\ &+ |\Psi(n)Y(n)(P_0 + P_-)| \sum_{k=0}^{n_1} \|Y^{-1}(k+1)f(k)\| + \\ &+ K \sum_{k=n_1+1}^{n-1} \|\Psi(k)f(k)\| + K \sum_{k=n}^{\infty} \|\Psi(k)f(k)\| < \\ &< \frac{\varepsilon}{5} + K \frac{\varepsilon}{5K} + \frac{\varepsilon}{5} + \frac{\varepsilon}{5} + K \frac{\varepsilon}{5K} = \varepsilon. \end{aligned}$$

This shows that  $\lim_{n \rightarrow +\infty} \|\Psi(n)x(n)\| = 0$ .

Similarly,  $\lim_{n \rightarrow -\infty} \|\Psi(n)x(n)\| = 0$ .

The proof is now complete.

**Corollary 2.** Suppose that:

1°. the homogeneous equation (2) has no nontrivial  $\Psi$ - bounded solution on  $\mathbb{Z}$ ;

2°. the fundamental matrix  $Y$  of (2) satisfies:

a). the conditions (5) for some  $K > 0$ ;

b). the conditions:

$$\text{i). } \lim_{n \rightarrow +\infty} |\Psi(n)Y(n)P_-| = 0$$

$$\text{ii). } \lim_{n \rightarrow -\infty} |\Psi(n)Y(n)P_+| = 0.$$

2°. the function  $f : \mathbb{Z} \rightarrow \mathbb{R}^d$  is  $\Psi$ - summable on  $\mathbb{Z}$ .

Then, the equation (1) has a unique solution  $x$  on  $\mathbb{Z}$  such that

$$\lim_{n \rightarrow \pm\infty} \|\Psi(n)x(n)\| = 0.$$

Proof. It results from the above Corollary and Theorem 2.

Note that the Theorem 2 (and the Corollary 2) is no longer true if we require that the function  $f$  be  $\Psi$ - bounded on  $\mathbb{Z}$ , instead of the condition 2° of the Theorem. This is shown by the next

**Example 1.** Consider the system (1) with

$$A(n) = \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & 2 \end{pmatrix}$$

and

$$f(n) = \begin{cases} \begin{pmatrix} 1 \\ 0 \end{pmatrix}, & n = 0, 1, 2, \dots \\ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, & n = -1, -2, \dots \end{cases}.$$

Then,  $Y(n) = \begin{pmatrix} 2^{-n} & 0 \\ 0 & 2^n \end{pmatrix}$  is the fundamental matrix of (2) with  $Y(0) = I_2$ .

Consider  $\Psi(n) = \begin{pmatrix} 1 & 0 \\ 0 & 3^{-n} \end{pmatrix}$ ,  $n \in \mathbb{Z}$ .

The first hypothesis of the Theorem 2 is satisfied with

$$P_0 = O_2, P_- = I_2, P_+ = O_2 \text{ and } K = 1.$$

In addition, we have  $\|\Psi(n)f(n)\| = 1$  for  $n \geq 0$  and  $\|\Psi(n)f(n)\| = 0$  for  $n < 0$ . The function  $f$  is not  $\Psi$ - summable on  $\mathbb{Z}$ , but it is  $\Psi$ - bounded on  $\mathbb{Z}$ .

On the other hand, the solutions on  $\mathbb{Z}$  of the system (1) are

$$x(n) = \begin{cases} \begin{pmatrix} 2^{-n}c_1 \\ 2^n c_2 \end{pmatrix}, & \text{for } n < 0, \\ \begin{pmatrix} 2^{-n}c_1 + 2^{-2^{1-n}} \\ 2^n c_2 \end{pmatrix}, & \text{for } n \geq 0. \end{cases}, \quad c_1, c_2 \in \mathbb{R}.$$

It results from this that there is no solution  $x$  for  $\lim_{n \rightarrow \pm\infty} \|\Psi(n)x(n)\| = 0$ .

**Acknowledgment.** The author would like to thank the anonymous referee for his/her valuable comments and suggestions.

### References.

- [1]. P. N. Boi – Existence of  $\Psi$ - bounded solutions on  $\mathbb{R}$  for nonhomogeneous linear differential equations, *Electronic Journal of Differential Equations*, Vol. 2007(2007), No. 52, pp. 1-10.
- [2]. W. A. Coppel – *Stability and Asymptotic Behavior of Differential Equations*, Heath, Boston, 1965.
- [3]. A. Diamandescu – Existence of  $\Psi$  - bounded solutions for a system of differential equations, *Electronic Journal of Differential Equations*, Vol. 2004(2004), No. 63, pp. 1 - 6,
- [4]. A. Diamandescu – Note on the  $\Psi$  - boundedness of the solutions of a system of differential equations, *Acta. Math. Univ. Comenianae*, Vol. LXXIII, 2(2004), pp. 223 - 233
- [5]. A. Diamandescu – A Note on the  $\Psi$ - boundedness for differential systems, *Bull. Math. Soc. Sc. math. Roumanie*, Tome 48(96), No. 1, 2005, 33-43.
- [6]. A. Diamandescu – Existence of  $\Psi$ - bounded solutions for nonhomogeneous linear difference equations, to appear.
- [7]. Y. Han, J. Hong – Existence of  $\Psi$ - bounded solutions for linear difference equations, *Applied Mathematics Letters*, 20(2007) 301–305.

Author's address

Aurel Diamandescu

University of Craiova,

Department of Applied Mathematics,

13, "Al. I. Cuza" st., 200585 Craiova, România.

E-mail: adiamandescu@central.ucv.ro

(Received June 5, 2008)