

On the radially symmetric solutions of a BVP for a class of nonlinear elliptic partial differential equations

JENŐ HEGEDŰS

(JATE, Bolyai Institute, Szeged, Hungary)
hegedusj@math.u-szeged.hu

Abstract. Uniqueness and comparison theorems are proved for the BVP of the form

$$\Delta u(x) + g(x, u(x), |\nabla u(x)|) = 0, \quad x \in B, u|_{\Gamma} = a \in \mathbb{R} \quad (\Gamma := \partial B),$$

where B is the unit ball in \mathbb{R}^n centered at the origin ($n \geq 2$). We investigate radially symmetric solutions, their dependence on the parameter $a \in \mathbb{R}$, and their concavity.

AMS subject classifications: 35B30, 35J60, 35J65.

Key words: radially symmetric solution, nonlinear elliptic partial differential equations, uniqueness, monotone dependence on the boundary value, concavity.

Introduction

Radially symmetric solutions to Dirichlet problem for the nonlinearly perturbed Laplace operator are investigated by many authors, see e.g. [1]-[4].

In [1] it is proven for a wide class of perturbations that the smooth positive solutions of the homogeneous Dirichlet problem in a ball are necessarily radially symmetric. The perturbation of the Laplacian in the paper [2], is $f(u)$ with a locally Lipschitz function f ; a BVP with a condition at infinity is considered, reduced to an ODE problem and sufficient conditions are given that guarantee the solvability of the original problem. In the papers [3], [4] nonlinear ODE-BVP-s (partly related to perturbed Laplacian) are considered on the intervals (a, b) and $(0, 1)$ respectively; the term y'' is perturbed with the sum

$$g(x, y') + f(x, y), \quad \frac{n-1}{x}y' + f(x, y)$$

respectively (where g is locally Lipschitz), and sufficient conditions (certain additional restrictions on f and g) of the existence and uniqueness of the positive solution y are presented. These cases do not cover the general case of perturbations y'' of the form

$$\frac{n-1}{x}y' + f(x, y, y') \quad x \in (0, R), \quad 0 < R < \infty,$$

i.e. the case of the perturbations Δu with $f(|x|, u, \pm|\nabla u|)$.

We remark that fundamental results for the investigations in [1] were already given in [5]. The contribution of the author of [5] to the theory of radially symmetric solutions of nonlinear elliptic PDE-s (mainly on the whole space \mathbb{R}^n and more generally for the m -Laplacian) can be found in [6].

This paper is in final form and no version of it will be submitted for publication elsewhere.

Recently G. Bognár [7] considered the following BVP in the unit ball $B := \{x \in \mathbb{R}^n | \rho \equiv |x| < 1\}$ ($\Gamma := \partial B$):

$$(A_1) \quad \Delta u(x) + \exp(\lambda u(x) + \kappa |\nabla u(x)|) = 0 \quad x \in B; \quad \kappa, \lambda \leq 0 \text{ are constants,}$$

$$(A_2) \quad u \in C^2(B) \cap C(\overline{B}), u(x) = v(|x|) \equiv v(\rho),$$

$$(A_3) \quad u|_{\Gamma} = a = \text{const.} \quad a \geq 0.$$

Existence and uniqueness results were established by the author and it was shown that the solution u depends monotonically on the parameter a .

The purpose of the present paper is to prove uniqueness, monotonicity, and concavity results for the solutions of the more general BVP: **Problem 1**:

$$(1.1) \quad \Delta u(x) + f(|x|, u(x), |\nabla u(x)|) = 0 \quad x \in B,$$

$$(1.2) \quad u \in C^2(B) \cap C(\overline{B}), \exists v : [0, 1] \rightarrow \mathbb{R} : v(\rho) \equiv v(|x|) = u(x) \quad \forall x \in \overline{B},$$

$$(1.3) \quad u|_{\Gamma} = a \in \mathbb{R}.$$

Here $f \in C(G_a; (0, \infty))$, $a \in \mathbb{R}$ is arbitrarily fixed; $G_a := [0, 1] \times [a, \infty) \times [0, \infty)$, B is the unit ball centered at the origin, and $\rho := |x| \quad x \in \overline{B}$.

The method used here is, partly, a modification of that of [7]. Some results are proved without using radial symmetry. These proofs are based upon the techniques communicated in [9].

To prepare our general results we formulate some of them in simplified versions:

Theorem A. If $f \in C(G_a; (0, \infty))$ and $f(\rho, t, \beta)$ is strongly decreasing in $t \in [a, \infty)$, then **Problem 1** may have no more than one solution.

Theorem B. If $f \in C(G_a; (0, \infty))$ and $f(\rho, t_1, t_2)$ is nonincreasing both in $t_1 \in [a, \infty)$, and $t_2 \in [0, \infty)$, then for the (radially symmetric) solutions u_1 and u_2 of **Problem 1** with the property:

$$u_1|_{\Gamma} \equiv a_1 > u_2|_{\Gamma} \equiv a_2 \geq a$$

inequalities

$$v_1(\rho) \equiv u_1(|x|) \geq v_2(\rho) \equiv u_2(|x|) \quad x \in \overline{B}, \quad v'_1(\rho) \geq v'_2(\rho) \quad \rho \in [0, 1)$$

hold.

Finally a concavity result:

Theorem C. Let $f \in C(G_a; (0, \infty))$, and let $f(\rho, t_1, t_2)$ be nonincreasing both in $t_1 \in [a, \infty)$, and $t_2 \in [0, \infty)$. Then there exists a constant $K(a)$ such, that $0 < f \leq K(a) < \infty$, and any of the assumptions (C_1) , (C_2)

$$(C_1) \quad f(t, a + \frac{K(a)}{n} \frac{1-t^2}{2}, \frac{K(a)}{n} t) \geq K(a) (1 - \frac{1}{n}) \quad t \in [0, 1],$$

$$(C_2) \quad f\left(t, a + \frac{K(a)}{2n}, \frac{K(a)}{n}\right) \geq K(a)\left(1 - \frac{1}{n}\right) \quad t \in [0, 1]$$

guarantees the concavity of the solution of **Problem 1**. For the case

$$f(\rho, u, |\nabla u|) \equiv \exp(\lambda u + \kappa|\nabla u|) \quad \lambda, \kappa \leq 0$$

considered in [7], the assumption (C_1) turns into (C_3) :

$$(C_3) \quad \exp\left\{\lambda\left[a + \frac{e^{\lambda a}}{n} \frac{1-t^2}{2}\right] + \kappa \frac{e^{\lambda a}}{n} t\right\} \geq e^{\lambda a}\left(1 - \frac{1}{n}\right) \quad t \in [0, 1].$$

One of the simplest sufficient conditions for the concavity of u for this special case is

$$(C_4) \quad \kappa \leq \lambda(\leq 0), \quad -1 \leq \kappa e^{\lambda a}.$$

1. Uniqueness results.

We shall prove (under the corresponding conditions) two theorems on the uniqueness of solution of **Problem 1**. The first one will be a consequence of a classical, simple uniqueness theorem related to the problem more general than **Problem 1**.

Theorem 1. Let $w(t) := f(\alpha, t, \beta) \quad t \in [a, \infty)$ for every $\alpha \in [0, 1]$, $\beta \in [0, \infty)$ fixed be strongly decreasing in t on the interval $[a, \infty)$; then **Problem 1** has no more than one solution.

Instead of a direct proof consider **Problem 2** (mentioned above) in an arbitrary bounded domain Ω of \mathbb{R}^n with the boundary $\Gamma := \partial\Omega$:

Problem 2.

$$(4) \quad u \in C^2(\Omega) \cap C(\bar{\Omega})$$

$$(5) \quad (\Delta u)(x) + g(x, u(x), u_{x_1}(x), \dots, u_{x_n}(x)) = 0 \quad x \in \Omega,$$

$$(6) \quad u|_{\Gamma} = \varphi \in C(\Gamma),$$

where

$$g \in C(\bar{\Omega} \times \mathbb{R}^{n+1}).$$

Theorem 2. If $w(t) := g(\underline{\alpha}, t, \underline{\beta}) \quad t \in \mathbb{R}$ is strongly decreasing in t for any

$$\underline{\alpha} \in \Omega, \quad \underline{\beta} \in \mathbb{R}^n$$

fixed, then **Problem 2** has no more than one solution.

This theorem is very close to the Theorem 9.3 (p.208) of the book [8].

Proof. Suppose that there exist two different solutions of **Problem 2**: u_1 and u_2 . Define $u := u_1 - u_2$ and suppose that there exists a point $y \in \Omega$ such, that $u(y) \neq 0$. Without loss of the generality it may be supposed that $u(y) < 0$. Letting

$$m := \min_{x \in \overline{B}} u(x)$$

we see, that $m < 0$, and there exists a point $x_0 \in \Omega$ of global minimum of the function $u(x) \quad x \in \overline{\Omega}$ i.e. $\exists x_0 \in \Omega$ such that

$$0 > m = u(x_0) \leq u(x) \quad \forall x \in \overline{\Omega}.$$

Consequently we have

$$(7) \quad (\Delta u)(x_0) \geq 0, \quad u_{x_i}(x_0) = 0 \quad i = \overline{1, n}.$$

On the other hand we know, that

$$(8) \quad (\Delta u_1)(x_0) + g(x_0, u_1(x_0), (\text{grad } u_1)(x_0)) = 0,$$

$$(9) \quad (\Delta u_2)(x_0) + g(x_0, u_2(x_0), (\text{grad } u_2)(x_0)) = 0,$$

therefore subtracting (9) from (8) we have

$$(10) \quad (\Delta u)(x_0) = g(x_0, u_2(x_0), (\text{grad } u_2)(x_0)) - g(x_0, u_1(x_0), (\text{grad } u_1)(x_0)).$$

Here arguments $(\text{grad } u_2)(x_0), (\text{grad } u_1)(x_0)$ are common in virtue of equalities $u_{x_i}(x_0) = 0 \quad i = \overline{1, n}$ (see (7)), therefore using the relations

$$0 > u(x_0) = m \equiv u_1(x_0) - u_2(x_0)$$

and their consequence $u_2(x_0) > u_1(x_0)$; from the monotonicity condition on $w(t)$ we get

$$f(x_0, u_2(x_0), (\text{grad } u_2)(x_0)) - f(x_0, u_1(x_0), (\text{grad } u_1)(x_0)) < 0.$$

So, in (10) we have

$$(11) \quad (\Delta u)(x_0) < 0,$$

that contradicts inequality of (7). Theorem 2 is proved.

Remark 1. For the proof of Theorem 1 it is enough to apply Theorem 2 for the case $\Omega := B$ with the nonlinear part g (appearing in **Problem 2**) defined by the formula

$$g(x, u, u_{x_1}, \dots, u_{x_n}) := f \left(|x|, u, \left(\sum_{i=1}^n u_{x_i}^2 \right)^{1/2} \right) \quad x \in \overline{B},$$

where f is the nonlinearity appearing in **Problem 1**.

Next we explain another result on the uniqueness of the solution to the **Problem 1** without assumption on strong decrease of $f(\alpha, t, \beta)$ in t . However we need that $f(\alpha, t_1, t_2)$ is nonincreasing both in t_1 and t_2 . Here in the proof we will use the radial symmetricity of the solutions.

Theorem 3. Let function f appearing in differential equation of **Problem 1** satisfy conditions:

(i) $w(t) := f(\alpha, t, \beta)$ is nonincreasing in $t \in [a, \infty)$ for every fixed $\alpha, \beta (\alpha \in [0, 1], \beta \in [0, \infty))$, and

(ii) $\tilde{w}(t) := f(\alpha, \beta, t)$ is nonincreasing in $t \in [0, \infty)$ for every fixed $\alpha, \beta (\alpha \in [0, 1], \beta \in [a, \infty))$.

Then **Problem 1** has no more, than one solution.

Proof. Suppose, that there exist two different solutions: $u_1, u_2 (u_1(x) = v_1(|x|), u_2(x) = v_2(|x|) x \in \overline{B})$ of **Problem 1** with the same boundary value $a \in \mathbb{R}$. We introduce the notation

$$v(\rho) := v_1(\rho) - v_2(\rho) \quad \rho \in [0, 1].$$

From the assumption, that $f > 0$ and u_1, u_2 are solutions of Problem 1 (especially they are superharmonic and radially symmetric in B) easily follows that

$$(12) \quad \begin{aligned} v &\in C^2([0, 1]) \cap C([0, 1]), \quad v(1) = 0, \quad v'(0) = 0, \\ \Delta u_i(x) + f(|x|, u_i(x), |\nabla u_i(x)|) &= \\ &= v_i''(\rho) + \frac{n-1}{\rho} v_i'(\rho) + f(\rho, v_i(\rho), -v_i'(\rho)) = 0 \quad x \in B, \quad \rho \in (0, 1) \quad i = 1, 2, \end{aligned}$$

and the multiplied by ρ^{n-1} version of the last equality of(12) holds:

$$(13) \quad (\rho^{n-1} v_i'(\rho))' + \rho^{n-1} f(\rho, v_i(\rho), -v_i'(\rho)) = 0 \quad \rho \in [0, 1], \quad i = 1, 2.$$

It can be supposed – without loss of the generality – that there exists a point $a_1 \in [0, 1]$ such, that $v(a_1) > 0$. Using the continuity of v on $[0, 1]$ it is trivial, that the interval $(0, 1)$ also contains a point a_1 such, that $v(a_1) > 0$. Let us fix such a point a_1 for the sequel. Our aim is to construct an interval $[\alpha, \beta] \subseteq [0, 1]$ such that

$$v(\rho) > 0 \quad \rho \in [\alpha, \beta], \quad v'(\rho) < 0 \quad \rho \in (\alpha, \beta], \quad v'(\alpha) = 0.$$

Let be

$$b := \sup\{\rho \in [0, 1] \mid v(\rho) > 0\}$$

It is clear, that

$$b \in (0, 1], \quad v(b) = 0,$$

and that

$$a_1 \in (0, b).$$

Further let

$$d := \inf\{\rho \in (a_1, b] \mid v(\rho) = 0\}.$$

It is clear, that

$$v(d) = 0.$$

Then let be

$$(14) \quad c := 0 \quad \text{if} \quad v(\rho) > 0 \quad \rho \in [0, a_1],$$

and

$$(15) \quad c := \sup\{\rho \in [0, a_1] \mid v(\rho) = 0\} \quad \text{otherwise.}$$

In the case of (2.43)

$$v(c) = 0$$

holds automatically. Further, denoting by M the maximum of the function v on $[c, d]$ ($M > 0$) let us introduce

$$e := \sup\{\rho \in [c, d] \mid v(\rho) = M\}.$$

We remark that for the case of (14) $e \in [c, d] \equiv [0, d]$ and

$$(16) \quad v'(e) \equiv v'(0) = 0$$

in virtue of (12) if $e = c = 0$; and $v'(e) = 0$ if $e \in (c, d) \equiv (0, d)$ using the fact that $v(e) = M$ i.e. e is a point of interior global maximum of the function v on the interval $[c, d]$. In the case of (15)

$$(17) \quad e \in (c, d), \quad v(e) = M, \quad v'(e) = 0$$

hold automatically because e is an interior point of global maximum of v on $[c, d]$.

The assumption $v'(\rho) \geq 0 \quad \rho \in [e, d]$ leads to contradiction in both cases (14) and (15), because if $d_1 < d$, and $d_1 \rightarrow d$ then

$$v(e) + \int_e^{d_1} v'(\rho) \, d\rho \rightarrow v(d) = 0$$

and

$$v(e) + \int_e^{d_1} v'(\rho) \, d\rho \equiv M + \int_e^{d_1} v'(\rho) \, d\rho \geq M > 0 \quad (\forall d_1 \in (e, d)).$$

Consequently there exists a point $\beta \in (e, d)$ such, that $v'(\beta) < 0$. Fixing such a point, it is easy to show – using (16) and continuity of v' on $[0, 1]$ – that there exists an interval $[\alpha, \beta] \subseteq [c, d]$ such, that

$$(18) \quad v'(\rho) < 0 \quad \rho \in (\alpha, \beta], \quad v'(\alpha) = 0, \quad v(\rho) > 0 \quad \rho \in [\alpha, \beta].$$

Namely, for the both of the cases (14) and (15) α may be chosen as

$$(19) \quad \alpha := \sup\{\rho \in [e, \beta] | v'(\rho) = 0\} \equiv \sup \mathcal{M}$$

because the set \mathcal{M} is non empty ($e \in \mathcal{M}$), and using the property $v' \in C[0, 1]$ (see (12))

$$(20) \quad \alpha \in [e, \beta), \quad v'(\alpha) = 0.$$

The next step of the proof is the using of the validity of differential equation of **Problem 1** for v_1, v_2 on the interval $I \equiv (\alpha, \beta)$ choiced above:

$$(\rho^{n-1}v'_1(\rho))' + \rho^{n-1}f(\rho, v_1(\rho), -v'_1(\rho)) = 0,$$

$$(\rho^{n-1}v'_2(\rho))' + \rho^{n-1}f(\rho, v_2(\rho), -v'_2(\rho)) = 0,$$

from which after subtracting we get

$$(\rho^{n-1}v'(\rho))' + \rho^{n-1}[f(\rho, v_1, -v'_1) - f(\rho, v_2, -v'_2)] = 0$$

i.e.

$$(\rho^{n-1}v'(\rho))' = \rho^{n-1}[f(\rho, v_2, -v'_2) - f(\rho, v_1, -v'_1)].$$

Subtracting and adding in the brackets of the right hand side the term

$$f(\rho, v_1(\rho), -v'_2(\rho))$$

we get

$$(21) \quad (\rho^{n-1}v'(\rho))' = \delta_1(\rho) + \delta_2(\rho) \equiv \delta(\rho) \quad \rho \in [\alpha, \beta],$$

where

$$\delta_1(\rho) := \rho^{n-1}[f(\rho, v_2(\rho), -v'_2(\rho)) - f(\rho, v_1(\rho), -v'_2(\rho))] \quad \rho \in [\alpha, \beta],$$

$$\delta_2(\rho) := \rho^{n-1}[f(\rho, v_1(\rho), -v'_2(\rho)) - f(\rho, v_1(\rho), -v'_1(\rho))] \quad \rho \in [\alpha, \beta].$$

Of course $\delta_i \in C[\alpha, \beta]$ $i = 1, 2$. Moreover, taking into account the choice of the interval $[\alpha, \beta]$ we have the relations

$$(22) \quad v(\rho) \equiv v_1(\rho) - v_2(\rho) > 0 \quad \rho \in [\alpha, \beta], \quad v'(\rho) \equiv v'_1(\rho) - v'_2(\rho) < 0 \quad \rho \in (\alpha, \beta].$$

They imply the inequalities

$$(23) \quad \delta_i(\rho) \geq 0 \quad \rho \in [\alpha, \beta]$$

in virtue of the monotonicity - assumptions (i), (ii) of the theorem. Summarising the precedings results, we get

$$(24) \quad \delta \in C[\alpha, \beta], \quad \delta(\rho) \geq 0 \quad \rho \in [\alpha, \beta].$$

Integrating equality (21) over the interval (α, β) we get after rearranging:

$$\beta^{n-1}v'(\beta) = \alpha^{n-1}v'(\alpha) + \int_{\alpha}^{\beta} \delta(\rho) d\rho,$$

from which using equality $v'(\alpha) = 0$ we get

$$\beta^{n-1}v'(\beta) = \int_{\alpha}^{\beta} \delta(\rho) d\rho \geq 0,$$

consequently $v'(\beta) \geq 0$ that contradicts the choice of β as a point, such, that $v'(\beta) < 0$. Theorem is proved.

Now, let us formulate a weakly generalized Problem 1, namely **Problem 3**:

$$(25) \quad u \in C^2(B_0^R) \cap C(\overline{B_0^R}),$$

$$(26) \quad \Delta u(x) + f(|x|, u(x), |\nabla u(x)|) = 0 \quad x \in B_0^R,$$

$$(27) \quad \exists v : [0, R] \rightarrow \mathbb{R}, \quad v(|x|) = u(x) \quad x \in \overline{B_0^R} \quad (|x| \in [0, R]),$$

$$(28) \quad u|_{\Gamma} = a \in \mathbb{R},$$

where $a \in \mathbb{R}$ is arbitrarily fixed; $R \in (0, \infty)$,

$$B_0^R := \{x \in \mathbb{R}^n \mid |x| < R\}, \quad \Gamma = \partial B_0^R \equiv \{x \in \mathbb{R}^n \mid |x| = R\},$$

and

$$(29) \quad f \in C(G_{a,R}; (0, \infty)),$$

where

$$G_{a,R} := [0, R] \times [a, \infty) \times [0, \infty).$$

Theorem 4. Let function f satisfy the monotonicity conditions: (i) $w(t) := f(\alpha, t, \beta)$ is nonincreasing in $t \in [a, \infty)$ for every fixed α, β ($\alpha \in [0, R], \beta \in [0, \infty)$),

(ii) $\tilde{w}(t) := f(\alpha, \beta, t)$ is nonincreasing in $t \in [0, \infty)$ for every fixed α, β ($\alpha \in [0, R], \beta \in [a, \infty)$).

Then **Problem 3** has no more than one solution.

Proof. The arguments used in the proof of Theorem 3 applied to $[0, R]$ instead of $[0, 1]$ show the validity of Theorem 4.

2. Comparison results

Theorem 5. Suppose that all of the assumptions included in the formulation of **Problem 3** are fulfilled, moreover assumptions (i), (ii) of Theorem 4 hold. Consider the problems

$$(30) \quad u_i \in C^2(B_0^R) \cap C(\overline{B_0^R}) \quad i = 1, 2,$$

$$(31) \quad \Delta u_i(x) + f(|x|, u_i(x), |\nabla u_i(x)|) = 0 \quad x \in B_0^R, \quad i = 1, 2,$$

$$(32) \quad \exists v_i : [0, R] \rightarrow \mathbb{R}, \quad v_i(|x|) = u_i(x) \quad x \in \overline{B_0^R} \quad (|x| \in [0, R]), \quad i = 1, 2,$$

$$(33) \quad u_i|_{\Gamma} = a_i \in \mathbb{R} \quad i = 1, 2,$$

where

$$a_1, a_2 \in \mathbb{R}, \quad a_1 > a_2 \geq a.$$

If $u_i \sim v_i \quad i = 1, 2$ are solutions of problems (30) - (33), then

$$(34) \quad v_1(\rho) \geq v_2(\rho) \quad \rho \in [0, R],$$

$$(35) \quad (0 \geq) v_1'(\rho) \geq v_2'(\rho) \quad \rho \in [0, R], \quad v_1'(0) = v_2'(0) = 0.$$

Proof. Let us begin with the proof of inequality (34). We introduce the notation

$$v(\rho) := v_1(\rho) - v_2(\rho) \quad \rho \in [0, R].$$

The arguments used for the derivation of the relations (12), (13) applied to B_0^R instead of B_0^1 give

$$(36) \quad v \in C^2([0, R)) \cap C([0, R]), \quad v(R) = a_1 - a_2 > 0, \quad v'(0) = 0,$$

and

$$(37) \quad (\rho^{n-1} v_i'(\rho))' + \rho^{n-1} f(\rho, v_i(\rho), -v_i'(\rho)) = 0 \quad \rho \in [0, R); \quad i = 1, 2.$$

If $v(\rho) > 0 \quad \rho \in [0, R]$ is also fulfilled, then $v_1(\rho) > v_2(\rho) \quad \rho \in [0, R]$ and (34) is proved. In the case, when there exists a point $b_1 \in [0, R)$ such that $v(b_1) = 0$ let

$$(38) \quad b := \sup\{\rho \in [0, R) | v(\rho) = 0\}.$$

It is clear that

$$b \in [0, R), \quad v(b) = 0.$$

If $b = 0$, then

$$(39) \quad v_1(\rho) > v_2(\rho) \quad \rho \in (0, R], \quad v_1(0) = v_2(0),$$

so (34) is fulfilled. If $b > 0$, then $b \in (0, R)$ and Theorem 4 applied to the ball B_0^b gives

$$(40) \quad v_1(\rho) = v_2(\rho) \quad \rho \in [0, b].$$

On the other hand $v(R) > 0$, and the definition of b implies the inequality

$$(41) \quad v_1(\rho) > v_2(\rho) \quad \rho \in (b, R].$$

Relations (40) combined with (41) give

$$v_1(\rho) \geq v_2(\rho) \quad \rho \in [0, R].$$

Next we prove the inequality (35). Suppose the contrary. Then using also the first one of the relations in (36) there exists a point $c_1 \in (0, R)$ such that $v'(c_1) < 0$. Introduce the notation

$$(42) \quad c := \sup\{c_1 \in (0, R] \mid v'(c_1) < 0\}.$$

It is clear that $c \in (0, R]$ and $v'(c) \leq 0$. Then we consider the three possible cases

$$(A) \quad v(\rho) > 0 \quad \rho \in [0, R],$$

$$(B) \quad v(0) = 0, \quad v(\rho) > 0 \quad \rho \in (0, R] \quad (b = 0),$$

$$(C) \quad v(\rho) \equiv 0 \quad \rho \in [0, b], \quad v(\rho) > 0 \quad \rho \in (b, R] \quad (b \in (0, R)).$$

In the cases (A),(B) let us choose a point $d \in (0, c)$ such, that $v'(d) < 0$. Then we define the set \mathcal{M} :

$$\mathcal{M} := \{\rho \in [0, d] \mid v'(\rho) = 0\}.$$

It is obvious, that $\mathcal{M} \neq \emptyset$ because $v'(0) = 0$ (see the last of the relations in (36)). Then let

$$e := \sup \mathcal{M}.$$

It is trivial that

$$e \in [0, d), \quad v'(e) = 0, \quad v'(\rho) < 0 \quad \rho \in (e, d].$$

Summarising, in the cases (A), (B) we have

$$v(\rho) > 0 \quad \rho \in (e, d], \quad v(e) \geq 0; \quad v'(\rho) < 0 \quad \rho \in (e, d], \quad v'(e) = 0,$$

consequently, the same arguments as in the proof of Theorems 3, 4, applied to the interval $(\alpha, \beta) := (e, d)$ lead to the inequality $v'(d) \geq 0$ that contradicts the choice of d for which $v'(d) < 0$.

For the case (C), first, remark that in virtue of the inequality (41)

$$(43) \quad v(\rho) > 0 \quad \rho \in (b, R],$$

moreover

$$(44) \quad v(b) = 0, \quad v(\rho) = 0 \quad \rho \in [0, b) \quad (v'_1(\rho) = v'_2(\rho) \quad \rho \in [0, b))$$

according to the definition of b and to Theorem 4 on the uniqueness in the ball B_0^b . Now (44) - using the property $v \in C^2[0, 1]$ - implies $v'(b) = 0$, consequently we have the same situation as in the case (B), but on the interval $[b, R]$ instead of interval $[0, R]$. The theorem is proven.

Remark 2. In fact, we proved a stronger result, than inequality (34) : namely, may occur three and only the following three cases:

$$(A) \quad v_1(\rho) > v_2(\rho) \quad \rho \in [0, R],$$

or

$$(B) \quad v_1(\rho) > v_2(\rho) \quad \rho \in (0, R], \quad v_1(0) = v_2(0),$$

or there exists a number $b \in (0, R)$ such that

$$(C) \quad v_1(\rho) = v_2(\rho) \quad \rho \in [0, b], \quad v_1(\rho) > v_2(\rho) \quad \rho \in (b, R].$$

On the other hand inequality (35)

$$(0 \geq) v'_1(\rho) \geq v'_2(\rho) \quad \rho \in [0, R] \quad (v'_1(0) = v'_2(0) = 0)$$

– in general – cannot be replaced by another, stronger one under assumptions of Theorem 5 (see e.g. the case, when f does not depend on argument u).

Theorem 6. All of the statements of Theorem 5 remain - except for inequality (35) - if in conditions of Theorem 5 assumptions (i), (ii) of Theorem 4 are replaced by condition:

$$w(t) := f(|x|, t, |\nabla u|) \sim f(\alpha, t, \beta)$$

is strongly decreasing in $t \in [a, \infty)$ for every fixed α, β ($\alpha \in [0, R], \beta \in [0, \infty)$).

This theorem is a corollary of a general comparison result, namely:

Theorem 7. Let u_1, u_2 be solutions of **Problem 2** satisfying conditions

$$u_i|_{\Gamma} = \varphi_i \in C(\Gamma) \quad i = 1, 2; \quad \varphi_1 \geq \varphi_2,$$

and suppose that function

$$w(t) := f(\underline{\alpha}, t, \underline{\beta}) \quad t \in \mathbb{R}$$

is strongly decreasing in $t \in \mathbb{R}$ for any $\underline{\alpha} \in \Omega$, $\underline{\beta} \in \mathbb{R}^n$ fixed. Then

$$u_1(x) \geq u_2(x) \quad x \in \Omega.$$

Moreover, if there exists a point $y \in \Gamma$ such that $\varphi_1(y) > \varphi_2(y)$, then may occur two, and only the following two cases:

$$(A) \quad u_1(x) > u_2(x) \quad \forall x \in \Omega,$$

or there exists a subset $\Omega_1 \neq \emptyset$ of Ω such that

$$0 < \mu(\Omega_1) \leq \mu(\Omega)$$

(μ is the n -dimensional Lebesgue measure) and

$$(B) \quad u_1(x) > u_2(x) \quad \forall x \in \Omega_1; \quad u_1(x) = u_2(x) \quad \forall x \in \Omega \setminus \Omega_1.$$

Proof. Let $u := u_1 - u_2$, and suppose that there exists a point $y \in \Omega$ such that $u(y) < 0$. Then there is a point $x_0 \in \Omega$ with the property:

$$u(x_0) = \min_{x \in \overline{\Omega}} u(x) \equiv m < 0,$$

and all that remains is to repeat the proof of Theorem 2 for to get a contradiction. Theorem is proven.

3. Concavity results.

Here we will present certain results on the concavity of the function $v : [0, 1] \rightarrow \mathbb{R}$, defined in the Introduction ((1.2)) by the relation $v(|x|) = u(x) \quad x \in \overline{B}$, where the function u is supposed to be a solution of **Problem 1**.

Theorem 8. Let $a \in \mathbb{R}$ in **Problem 1** be fixed, and suppose that

$$(i) \quad w(t) := f(\alpha, t, \beta)$$

is nonincreasing in $t \in [a, \infty)$ for every α, β fixed ($\alpha \in [0, 1]$, $\beta \in [0, \infty)$),

$$(ii) \quad \tilde{w}(t) := f(\alpha, \beta, t)$$

is nonincreasing in $t \in [0, \infty)$ for every α, β fixed ($\alpha \in [0, 1]$, $\beta \in [a, \infty)$).

If, in addition,

$$(iii) \quad f\left(t, a + \frac{K_a}{n} \frac{1-t^2}{2}, \frac{K_a}{n} t\right) \geq K_a \left(1 - \frac{1}{n}\right) \quad t \in [0, 1),$$

where

$$K_a := \sup_{G_a} f(= \max_{\rho \in [0,1]} f(\rho, a, 0))$$

then function v is concave (in non strong sense) on the interval $[0, 1)$.

In other words - if $\gamma = (\gamma_1, \gamma_2, \gamma_3)$ is a curve in \mathbb{R}^3 :

$$\gamma : \gamma_1 = t, \quad \gamma_2 = a + \frac{K_a(1-t^2)}{2n}, \quad \gamma_3 = \frac{K_a t}{n} \quad t \in [0, 1),$$

then condition (iii) means that

$$(iv) \quad f|_{\gamma} \geq K_a \left(1 - \frac{1}{n}\right).$$

Proof. Assumptions of the Theorem guarantee the uniqueness (see Theorem 3 in above) of the solution $u \sim v$ to the **Problem 1**. We know (see (12), (13)) that v has the following properties:

$$(45) \quad \begin{aligned} v &\in C^2[0, 1) \cap C[0, 1], \quad v(1) = a, \quad v'(0) = 0, \\ \Delta u(x) + f(|x|, u(x), |\nabla u(x)|) &= v''(\rho) + \frac{n-1}{\rho} v'(\rho) + f(\rho, v(\rho), -v'(\rho)) = 0 \\ x \in B, \quad \rho &\in (0, 1), \end{aligned}$$

and

$$(46) \quad (\rho^{n-1} v'(\rho))' + \rho^{n-1} f(\rho, v(\rho), -v'(\rho)) = 0 \quad \rho \in [0, 1).$$

Integrating equality (46) over the interval $[\delta, t]$ ($0 < \delta < t < 1$) we get

$$(47) \quad t^{n-1} v'(t) = \delta^{n-1} v'(\delta) - \int_{\delta}^t \rho^{n-1} f(\rho, v(\rho), -v'(\rho)) d\rho$$

from which passing to the limit as $\delta \rightarrow 0+0$ we obtain

$$(48) \quad t^{n-1} v'(t) = - \int_0^t \rho^{n-1} f(\rho, v(\rho), -v'(\rho)) d\rho \quad \forall t \in (0, 1).$$

Using the notation

$$\nu := -v' \quad (\nu(t) := -v'(t) \quad \forall t \in [0, 1])$$

the first and second of the relations of (45) give

$$v(t) - v(t_1) = \int_t^{t_1} \nu(s) ds \quad 0 \leq t < t_1 < 1, \quad v(t) - v(t_1) \rightarrow v(t) - a \text{ as } t_1 \rightarrow 1 - 0,$$

consequently there exists the improper integral

$$\int_t^1 \nu(s) ds := \lim_{t_1 \rightarrow 1-0} \int_t^{t_1} \nu(s) ds \quad t \in [0, 1),$$

and

$$(49) \quad v(t) = a + \int_t^1 \nu(s) ds \quad \forall t \in [0, 1],$$

From (48),(49) we obtain that function ν satisfies equality

$$(50) \quad \nu(t) = \int_0^t \left(\frac{\rho}{t}\right)^{n-1} f(\rho, a + \int_\rho^1 \nu(s) ds, \nu(\rho)) d\rho \quad t \in [0 + 0, 1)$$

which is understood at $t = 0 + 0$ in the limit sense. From the definition of K_a and equality (50) we get the inequality

$$(51) \quad (0 \leq) \nu(t) \leq \frac{K_a}{n} t \quad \forall t \in [0, 1).$$

To prove the theorem we have to show that

$$(52) \quad \nu'(t) \geq 0 \quad t \in [0, 1),$$

i.e.-using the last of the equalities in (45) for $\rho \in [0 + 0, 1)$ combined with (50) - the inequality

$$(53) \quad \begin{aligned} \nu'(t) &\equiv f(t, a + \int_t^1 \nu(s) ds, \nu(t)) - \\ &- \frac{n-1}{t} \int_0^t \left(\frac{\rho}{t}\right)^{n-1} f(\rho, a + \int_\rho^1 \nu(s) ds, \nu(\rho)) d\rho \geq 0 \quad \rho \in [0 + 0, 1). \end{aligned}$$

From (51) we obtain that

$$(54) \quad \begin{aligned} \nu'(t) &\geq f\left(t, a + \frac{K_a}{n} \frac{1-t^2}{2}, \frac{K_a}{n} t\right) - \frac{n-1}{t} \nu(t) \geq \\ &\geq f\left(t, a + \frac{K_a}{n} \frac{1-t^2}{2}, \frac{K_a}{n} t\right) - \frac{n-1}{t} \frac{K_a}{n} t \geq 0 \quad t \in [0, 1) \end{aligned}$$

in virtue of condition (iii). Theorem is proven.

Some concrete sufficient conditions for the special case of **Problem 1**, when

$$(55) \quad f(\rho, u, |\nabla u|) = e^{\lambda u + \mathcal{K}|\nabla u|} \quad \lambda, \mathcal{K} \in \mathbb{R}; \quad \lambda, \mathcal{K} \leq 0$$

are presented in the following

Theorem 9. Let $a \in \mathbb{R}$ be arbitrarily fixed in **Problem 1** with nonlinearity f of the form in (55). Then solution $u \sim v$ of **Problem 1** exists ([7]), is unique, and any of the following conditions (i) - (vi) guarantees the nonstrong concavity of solution v on $[0, 1]$; where we use the notation

$$d_n := \ln \left[\left(1 - \frac{1}{n} \right)^n \right] \quad n \in \mathbb{N}, \quad n \text{ is fixed} \quad n \geq 2 \quad (d_n < 0),$$

$$(i) \quad \lambda = \mathcal{K} = 0,$$

$$(ii) \quad \lambda = 0, \quad 0 > \mathcal{K} \geq d_n,$$

$$(iii) \quad \mathcal{K} = 0, \quad 0 > \frac{\lambda e^{\lambda a}}{2} \geq d_n,$$

$$(iv) \quad \mathcal{K} < \lambda < 0, \quad \mathcal{K} e^{\lambda a} \geq d_n,$$

$$(v) \quad \mathcal{K} = \lambda < 0, \quad \lambda e^{\lambda a} \geq d_n,$$

$$(vi) \quad \lambda < \mathcal{K} < 0, \quad \frac{e^{\lambda a} \cdot \lambda}{2} \left(1 + \frac{\mathcal{K}^2}{\lambda^2} \right) \geq d_n.$$

Proof. It is enough to prove that inequality (iii) of Theorem 8 is fulfilled in every of the cases (i) - (vi) of the present Theorem. Using that

$$f(t_1, t_2, t_3) \sim f(t_2, t_3) = e^{\lambda t_2 + \mathcal{K} t_3} \quad t_2 \in [a, \infty), \quad t_3 \in [0, \infty)$$

and relations

$$f(a, 0) = e^{\lambda a} \geq f(t_2, t_3) \quad t_2 \in [a, \infty), \quad t_3 \in [0, \infty)$$

we get that $K_a = e^{\lambda a}$. Substituting this value into inequality (iii) of Theorem 8, the desirable inequality gains the form

$$e^{\lambda \left[a + \frac{\epsilon \lambda a}{n} \frac{1-t^2}{2} \right] + \mathcal{K} \frac{\epsilon \lambda a}{n} t} \geq e^{\lambda a} \left(1 - \frac{1}{n} \right) \quad t \in [0, 1)$$

i.e.

$$e^{e^{\lambda a} [\lambda \frac{1-t^2}{2} + \mathcal{K}t] \frac{1}{n}} \geq (1 - \frac{1}{n}) \quad t \in [0, 1)$$

i.e.

$$e^{\lambda a [\lambda \frac{1-t^2}{2} + \mathcal{K}t]} \equiv g(t) \geq \ln[(1 - \frac{1}{n})^n] \equiv d_n \quad t \in [0, 1).$$

It is easy to prove in every of the cases (i) - (vi) that

$$\min_{t \in [0,1]} g(t) \geq d_n,$$

which completes the proof.

References.

- [1] GIDAS B., NI W. M., NIRENBERG L.; Symmetry and Related Properties via the Maximum Principle, Communications in Mathematical Physics, Volume **68**, Number 3 (1979), 209-243.
- [2] ATKINSON F. V., PELETIER L. A.; Ground States of $-\Delta u = f(u)$ and the Emden-Fowler Equation, Archive for Rational Mechanics and Analysis, Volume **93**, Number 2 (1986), 103-127.
- [3] BAXLEY J. V.; Some singular nonlinear boundary value problems, SIAM Journal on Mathematical Analysis, Volume **22**, Number 2 (1991), 463-479.
- [4] FINK A. M., GATICA J. A., HERNANDEZ G. E., WALTMAN P.; Approximation of solutions of singular second-order boundary value problems, SIAM Journal on Mathematical Analysis, Volume **22**, Number 2 (1991), 440-462.
- [5] SERRIN J.; A symmetry problem in potential theory, Archive for Rational Mechanics and Analysis, Volume **43** (1971), 304-318.
- [6] GAZZOLA F., SERRIN J., TANG M.; Existence of ground states and free boundary problems for quasilinear elliptic operators, Advances in Differential Equations, Volume 5, Numbers 1-3 (2000), 1-30.
- [7] BOGNÁR G.; On the radially symmetric solutions to a nonlinear PDE, Publ. Univ. of Miskolc, Series D. Natural Sciences., Vol. **36** No. 2., Mathematics (1996), 13-20.
- [8] GILBARG D., TRUDINGER N.S., Elliptic Partial Differential Equations of Second Order, Grundlehren der mathematischen Wissenschaften 224, A Series of Comprehensive Studies in Mathematics; Springer-Verlag, Berlin - Heidelberg - New York, 1977; pp. 1-401.
- [9] HEGEDŰS J.; On the solutions of a BVP for a class of nonlinear elliptic equations; Conference EQUADIFF 9, [Brno (Czech Republic), August 25-29, 1997; (lecture)].