

Construction of non-constant lower and upper functions for impulsive periodic problems

Irena Rachůnková* and Milan Tvrdý†

Summary. We present conditions ensuring the existence of piecewise linear lower and upper functions for the nonlinear impulsive periodic boundary value problem $u'' = f(t, u, u')$, $u(t_i+) = J_i(u(t_i))$, $u'(t_i+) = M_i(u'(t_i))$, $i = 1, 2, \dots, m$, $u(0) = u(T)$, $u'(0) = u'(T)$. This together with the existence principles which we proved in [5]–[7] allows us to prove new existence criteria, see Theorems 3.1 and 3.2.

Mathematics Subject Classification 2000. 34B37, 34B15, 34C25

Keywords. Second order nonlinear ordinary differential equation, impulse, periodic solution, lower function, upper function.

1. Introduction

This paper deals with the impulsive periodic boundary value problem

$$(1.1) \quad u'' = f(t, u, u'),$$

$$(1.2) \quad u(t_i+) = J_i(u(t_i)), \quad u'(t_i+) = M_i(u'(t_i)), \quad i = 1, 2, \dots, m,$$

$$(1.3) \quad u(0) = u(T), \quad u'(0) = u'(T),$$

where

$$(1.4) \quad \begin{cases} 0 < t_1 < \dots < t_m < T < \infty, \\ f \text{ satisfies the Carathéodory conditions on } [0, T] \times \mathbb{R}^2, \\ J_i \text{ and } M_i, i = 1, 2, \dots, m, \text{ are continuous functions on } \mathbb{R}. \end{cases}$$

There are several papers providing the existence results for such problems in terms of lower and upper functions, see e.g. [1]–[4], [8] and our papers [5]–[7]. However, up to now, only Proposition 1.3 in [6] gives conditions ensuring the existence of nonconstant (in particular, piecewise constant) lower and upper functions. The main

*Supported by the grant No. 201/01/1451 of the Grant Agency of the Czech Republic and by the Council of Czech Government J14/98:153100011

†Supported by the grant No. 201/01/1199 of the Grant Agency of the Czech Republic

This paper is in final form and no version of it will be submitted for publications elsewhere.

goal of this paper is to find conditions for f , J_i , M_i giving piecewise linear lower and upper functions for (1.1)–(1.3). This together with the existence principles which we proved in [5]–[7] allows us to present new existence criteria.

Throughout the paper we keep the following notation and conventions: For $J \subset \mathbb{R}$, $\mathbb{C}(J)$ is the set of real valued functions which are continuous on J , $\mathbb{C}^1(J)$ is the set of functions having continuous first derivatives on J , $\mathbb{L}(J)$ is the set of functions Lebesgue integrable on J and $\mathbb{L}_\infty(J)$ is the set of functions essentially bounded on J . For $u \in \mathbb{L}_\infty[0, T]$, we denote $\|u\|_\infty = \sup \operatorname{ess}_{t \in [0, T]} |u(t)|$. Further, $D = \{t_1, t_2, \dots, t_m\}$, $t_0=0$, $t_{m+1}=T$ and $\mathbb{C}_D^1[0, T]$ is the set of functions $u: [0, T] \mapsto \mathbb{R}$ of the form

$$u(t) = \begin{cases} u_{[0]}(t) & \text{if } t \in [0, t_1], \\ u_{[1]}(t) & \text{if } t \in (t_1, t_2], \\ \dots & \dots \\ u_{[m]}(t) & \text{if } t \in (t_m, T], \end{cases}$$

where $u_{[i]} \in \mathbb{C}^1[t_i, t_{i+1}]$ for $i = 0, 1, \dots, m$. Moreover, $\mathbb{AC}_D^1[0, T]$ stands for the set of functions $u \in \mathbb{C}_D^1[0, T]$ having first derivatives absolutely continuous on each subinterval (t_i, t_{i+1}) , $i = 0, 1, \dots, m$. For $u \in \mathbb{C}_D^1[0, T]$ and $i = 1, 2, \dots, m+1$ we define $u'(t_i) = u'(t_i-) = \lim_{t \rightarrow t_i-} u'(t)$ and $u'(0) = u'(0+) = \lim_{t \rightarrow 0+} u'(t)$.

1.1 Definition. A solution of the problem (1.1)–(1.3) is a function $u \in \mathbb{AC}_D^1[0, T]$ which satisfies the conditions (1.2) and (1.3) and for a.e. $t \in [0, T]$ fulfils the equation (1.1).

1.2 Definition. A function $\sigma_1 \in \mathbb{AC}_D^1[0, T]$ is called a lower function of the problem (1.1)–(1.3) if

$$(1.5) \quad \sigma_1''(t) \geq f(t, \sigma_1(t), \sigma_1'(t)) \quad \text{for a.e. } t \in [0, T],$$

$$(1.6) \quad \sigma_1(t_i+) = J_i(\sigma_1(t_i)), \quad \sigma_1'(t_i+) \geq M_i(\sigma_1'(t_i)), \quad i = 1, 2, \dots, m,$$

$$(1.7) \quad \sigma_1(0) = \sigma_1(T), \quad \sigma_1'(0) \geq \sigma_1'(T).$$

A function $\sigma_2 \in \mathbb{AC}_D^1[0, T]$ is an upper function of (1.1)–(1.3) if it satisfies

$$(1.8) \quad \sigma_2''(t) \leq f(t, \sigma_2(t), \sigma_2'(t)) \quad \text{for a.e. } t \in [0, T],$$

$$(1.9) \quad \sigma_2(t_i+) = J_i(\sigma_2(t_i)), \quad \sigma_2'(t_i+) \leq M_i(\sigma_2'(t_i)), \quad i = 1, 2, \dots, m,$$

$$(1.10) \quad \sigma_2(0) = \sigma_2(T), \quad \sigma_2'(0) \leq \sigma_2'(T).$$

2. Construction of nonconstant lower and upper functions

2.1 Theorem. Assume (1.4) and

$$(2.1) \quad \lim_{x \rightarrow \infty} (J_i(x) - x) = c_i \in \mathbb{R} \quad \text{for } i = 1, 2, \dots, m.$$

Denote

$$(2.2) \quad c = -\frac{1}{T} \sum_{i=1}^m c_i$$

and suppose that there are $A \in \mathbb{R}$, $\delta > 0$ and $\nu \in \{1, 2\}$ such that

$$(2.3) \quad (-1)^\nu f(t, x, y) \geq 0 \quad \text{for a.e. } t \in [0, T] \quad \text{and all } x \geq A, y \in [c - \delta, c + \delta]$$

and

$$(2.4) \quad (-1)^\nu (M_i(y) - y) \geq 0 \quad \text{for } y \in [c - \delta, c + \delta], \quad i = 1, 2, \dots, m.$$

Then for each $\tilde{A} \in [A, \infty)$ there exist $\tilde{k} \in (c - \delta, c + \delta)$ and $\sigma_\nu \in \mathbb{A}\mathbb{C}_D^1[0, T]$ such that $\sigma_\nu(t) \geq \tilde{A}$, $\sigma'_\nu(t) = \tilde{k}$ for $t \in [0, T]$ and, for $\nu = 1$ ($\nu = 2$), σ_ν is a lower (upper) function of (1.1)–(1.3).

Proof. Let $\nu = 2$ and $\tilde{A} \in [A, \infty)$.

• **STEP 1.** For $a, k \in \mathbb{R}$, $x \in \mathbb{R}$ and $i = 1, 2, \dots, m$, define

$$\varphi(t, a, k) = \begin{cases} a + kt & \text{if } t \in [0, t_1], \\ J_i(\varphi(t_i, a, k)) + k(t - t_i) & \text{if } t \in (t_i, t_{i+1}], \quad i = 1, 2, \dots, m, \end{cases}$$

and

$$\varepsilon_i(x) = J_i(x) - x - c_i.$$

By virtue of (2.1), there are constants $A_i > \tilde{A}$, $i = 1, 2, \dots, m$, such that

$$(2.5) \quad |\varepsilon_i(x)| < \frac{\delta T}{m} \quad \text{for all } x \geq A_i \quad \text{and } i = 1, 2, \dots, m.$$

Furthermore, we have

$$(2.6) \quad \varphi(t, a, k) = \begin{cases} a + kt & \text{if } t \in [0, t_1], \\ a + kt + \sum_{j=1}^i (c_j + \varepsilon_j(\varphi(t_j, a, k))) & \text{if } t \in (t_i, t_{i+1}], \quad i = 1, 2, \dots, m. \end{cases}$$

Put

$$\tilde{a} = \max_{j=1, \dots, m} A_j + 2 \left(\sum_{j=1}^m |c_j| + \delta T \right)$$

and suppose that $k \in [c - \delta, c + \delta]$. Then

$$\varphi(t_1, \tilde{a}, k) = \tilde{a} + k t_1 \geq A_1 + \left(1 - \frac{t_1}{T}\right) \sum_{j=1}^m |c_j| \geq A_1.$$

Furthermore, by (2.5), we have $|\varepsilon_1(\varphi(t_1, \tilde{a}, k))| \leq \frac{\delta T}{m}$. Consequently, in view of (2.6), we get

$$\begin{aligned} \varphi(t_1+, \tilde{a}, k) &= \tilde{a} + k t_1 + c_1 + \varepsilon_1(\varphi(t_1, \tilde{a}, k)) \\ &\geq A_2 + |c_1| + \delta T + \left(1 - \frac{t_1}{T}\right) \sum_{j=1}^m |c_j| - |c_1| - \frac{\delta T}{m} \geq A_2. \end{aligned}$$

Now, let $1 < i \leq m$ and let

$$(2.7) \quad \varphi(t_{j+}, \tilde{a}, k) \geq A_{j+1} \quad \text{and} \quad \varphi(t_j, \tilde{a}, k) \geq A_j$$

for each $j = 1, 2, \dots, i-1$. Then, by (2.5), we have

$$|\varepsilon_j(\varphi(t_j, \tilde{a}, k))| \leq \frac{\delta T}{m} \quad \text{for each } j = 1, 2, \dots, i-1.$$

Hence, using (2.6) and (2.5), we get

$$\varphi(t_i, \tilde{a}, k) \geq A_i + \sum_{j=1}^m \left(1 - \frac{t_i}{T}\right) |c_j| + \sum_{j=i}^m |c_j| + \left(1 - \frac{i-1}{m}\right) \delta T \geq A_i$$

and $|\varepsilon_i(\varphi(t_i, \tilde{a}, k))| \leq \frac{\delta T}{m}$. In view of (2.6), we have

$$\begin{aligned} \varphi(t_i+, \tilde{a}, k) &= \tilde{a} + k t_i + \sum_{j=1}^i (c_j + \varepsilon_j(\varphi(t_j, \tilde{a}, k))) \\ &\geq A_{i+1} + \sum_{j=1}^m \left(1 - \frac{t_i}{T}\right) |c_j| + \left(1 - \frac{i}{m}\right) \delta T \geq A_{i+1} \end{aligned}$$

which means that (2.7) is true for any $j \in \{1, 2, \dots, m\}$. Similarly we can show that $\varphi(T, \tilde{a}, k) \geq A_m$. Thus

$$(2.8) \quad \begin{cases} \varphi(t, \tilde{a}, k) \geq \tilde{A} \quad \text{and} \quad |\varepsilon_i(\varphi(t_i, \tilde{a}, k))| < \frac{\delta T}{m} \\ \text{for all } t \in [0, T], k \in [c - \delta, c + \delta] \quad \text{and} \quad i = 1, 2, \dots, m. \end{cases}$$

• STEP 2. We will prove that there is $\tilde{k} \in (c - \delta, c + \delta)$ such that

$$(2.9) \quad \varphi(0, \tilde{a}, \tilde{k}) = \varphi(T, \tilde{a}, \tilde{k}).$$

By virtue of (2.6) and (2.8), $\sum_{i=1}^m |\varepsilon_i(\varphi(t_i, \tilde{a}, k))| < \delta T$ and

$$\varphi(T, \tilde{a}, k) - \varphi(0, \tilde{a}, k) = T \left(k - c + \frac{1}{T} \sum_{i=1}^m \varepsilon_i(\varphi(t_i, \tilde{a}, k)) \right)$$

for each $k \in [c - \delta, c + \delta]$. In particular, $\varphi(T, \tilde{a}, c - \delta) - \varphi(0, \tilde{a}, c - \delta) < 0$ and $\varphi(T, \tilde{a}, c + \delta) - \varphi(0, \tilde{a}, c + \delta) > 0$. Since φ is continuous, the existence of $\tilde{k} \in (c - \delta, c + \delta)$ satisfying (2.9) follows.

• **STEP 3.** Define $\sigma_2(t) = \varphi(t, \tilde{a}, \tilde{k})$ for $t \in [0, T]$. Then $\sigma_2'(t) = \tilde{k}$ for $t \in [0, T]$ and $\sigma_2(t_i+) = J_i(\sigma_2(t_i))$ for $i = 1, 2, \dots, m$. By (2.4) we have $\tilde{k} \leq M_i(\tilde{k})$, i.e. σ_2 satisfies (1.9). Moreover, by (2.8) and (2.9), we have $\sigma_2(t) \geq \tilde{A}$ on $[0, T]$ and $\sigma_2(0) = \sigma_2(T)$ and so (1.10) is true. Finally, by (2.3), σ_2 fulfils (1.8), i.e. σ_2 is an upper function for (1.1)–(1.3).

The case $\nu = 1$ can be treated analogously. □

Theorem 2.1 gives piecewise linear lower and upper functions which are bounded below. Now we will show conditions guaranteeing the existence of lower or upper functions bounded above. This is the contents of the next theorem. Its proof is similar to that of Theorem 2.1.

2.2 Theorem. Assume (1.4). Further, let $d_i \in \mathbb{R}$, $i = 1, 2, \dots, m$, $d, B \in \mathbb{R}$, $\eta > 0$ and $\nu \in \{1, 2\}$ be such that

$$\lim_{x \rightarrow -\infty} (J_i(x) - x) = d_i \in \mathbb{R} \quad \text{for } i = 1, 2, \dots, m, \quad d = -\frac{1}{T} \sum_{i=1}^m d_i,$$

$$(-1)^\nu f(t, x, y) \geq 0 \quad \text{for a.e. } t \in [0, T] \quad \text{and all } x \leq B, \quad y \in [d - \eta, d + \eta],$$

$$(-1)^\nu (M_i(y) - y) \geq 0 \quad \text{for } y \in [d - \eta, d + \eta], \quad i = 1, 2, \dots, m,$$

Then for each $\tilde{B} \leq B$ there exist $\tilde{k} \in (d - \eta, d + \eta)$ and $\sigma_\nu \in \mathbb{A}\mathbb{C}_D^1[0, T]$ such that $\sigma_\nu(t) \leq \tilde{B}$, $\sigma_\nu'(t) = \tilde{k}$ for $t \in [0, T]$ and, for $\nu = 1$ ($\nu = 2$), σ_ν is a lower (upper) function of (1.1)–(1.3).

2.3 Remark. Let (1.4) hold. Assume that $c_i, d_i \in \mathbb{R}$ and $A \in (0, \infty)$ are such that

$$(2.10) \quad J_i(x) = \begin{cases} x + c_i & \text{for } x \geq A \\ x + d_i & \text{for } x \leq -A, \quad i = 1, 2, \dots, m, \end{cases}$$

$$\begin{cases} f(t, x, c) \geq 0 & \text{for a.e. } t \in [0, T] \quad \text{and all } x \geq A, \\ f(t, x, d) \leq 0 & \text{for a.e. } t \in [0, T] \quad \text{and all } x \leq -A, \end{cases}$$

and

$$(2.11) \quad M_i(c) \geq c, \quad M_i(d) \leq d, \quad i = 1, 2, \dots, m.$$

Let $\tilde{A} \geq A$. Then, according to the proof of Theorem 2.1, we can see that

$$\sigma_2(t) = \begin{cases} \tilde{A} + \sum_{j=1}^m |c_j| + ct & \text{for } t \in [0, t_1), \\ \tilde{A} + \sum_{j=1}^m |c_j| + ct + \sum_{j=1}^i c_j & \text{for } t \in (t_i, t_{i+1}], \quad i = 1, 2, \dots, m \end{cases}$$

and

$$\sigma_1(t) = \begin{cases} -\tilde{A} - \sum_{j=1}^m |d_j| + dt & \text{for } t \in [0, t_1), \\ -\tilde{A} - \sum_{j=1}^m |d_j| + dt + \sum_{j=1}^i d_j & \text{for } t \in (t_i, t_{i+1}], \quad i = 1, 2, \dots, m \end{cases}$$

are respectively upper and lower functions of (1.1)–(1.3) satisfying

$$\tilde{A} \leq \sigma_2(t) \leq \tilde{A} + 2 \sum_{j=1}^m |c_j| \quad \text{and} \quad -\tilde{A} - 2 \sum_{j=1}^m |d_j| \leq \sigma_1(t) \leq -\tilde{A} \quad \text{for } t \in [0, T].$$

If all inequalities in (2.10) and (2.11) are reversed, then σ_2 becomes a lower function and σ_1 an upper function.

3. New existence criteria

Our main results are Theorems 3.1 and 3.2 which provides new existence criteria for the problem (1.1)–(1.3).

3.1 Theorem. *Let the assumptions of Theorem 2.1 be satisfied for $\nu = 2$ and let the assumptions of Theorem 2.2 be satisfied for $\nu = 1$. Assume that J_i are increasing on \mathbb{R} and M_i are nondecreasing on \mathbb{R} for $i = 1, 2, \dots, m$. Finally, let for each compact interval $K \subset \mathbb{R}$ there exist $h_K \in \mathbb{L}[0, T]$ and $\omega_K \in \mathbb{C}([1, \infty))$ such that $h_K \geq 0$ on $[0, T]$, $\omega_K > 0$ on $[1, \infty)$, $\int_1^\infty ds/\omega_K(s) = \infty$ and $|f(t, x, y)| \leq \omega_K(|y|)(|y| + h_K(t))$ for a.e. $t \in [0, T]$ and all $x \in K$, $|y| > 1$. Then the problem (1.1)–(1.3) has a solution.*

Proof. By Theorem 2.1, for each $\tilde{A} \geq A$, there is an upper function σ_2 of (1.1)–(1.3) such that $\sigma_2 \geq \tilde{A}$ on $[0, T]$. By Theorem 2.2, for each $\tilde{B} \leq B$, there is a lower function σ_1 of (1.1)–(1.3) such that $\sigma_1 \leq \tilde{B}$ on $[0, T]$. Choose \tilde{A}, \tilde{B} in such a way that $\tilde{B} \leq \tilde{A}$. Hence $\sigma_1 \leq \sigma_2$ on $[0, T]$ and all the assumptions of [5, Theorem 3.1] are satisfied. Therefore (1.1)–(1.3) has a solution. \square

3.2 Theorem. *Let the assumptions of Theorem 2.1 be satisfied for $\nu = 1$ and let the assumptions of Theorem 2.2 be satisfied for $\nu = 2$. Assume that J_i are increasing on \mathbb{R} and M_i are nondecreasing on \mathbb{R} for $i = 1, 2, \dots, m$. Finally, let there exist $h \in \mathbb{L}[0, T]$ such that $|f(t, x, y)| \leq h(t)$ for a.e. $t \in [0, T]$ and all $x, y \in \mathbb{R}$. Then the problem (1.1)–(1.3) has a solution.*

Proof. By Theorems 2.1 and 2.2 there are a lower function σ_1 and an upper function σ_2 of (1.1)–(1.3). The existence of a solution to (1.1)–(1.3) follows by [7, Theorem 3.1]. \square

3.3 Example. Let $k \in \mathbb{N} \cup \{0\}$, $\gamma \in (0, \infty)$, $p_j \in \mathbb{L}_\infty[0, T]$, $j = 0, 1, \dots, 2k$, $p_{2k+1} \in \mathbb{L}[0, T]$, $p_{2k+1} \geq \gamma$ a.e. on $[0, T]$, $q_1, q_2 \in \mathbb{L}_\infty[0, T]$, $c_i \in (0, \infty)$, $\alpha_i \in [0, 1)$ and $\beta_i \in \mathbb{R}$, $i = 1, 2, \dots, m$. Consider the problem (1.1)–(1.3), where

$$f(t, x, y) = \sum_{j=0}^{2k+1} p_j(t) x^j + q_1(t) y + q_2(t) y^2, \quad J_i(x) = x + \frac{2c_i}{\pi} \arctan x, \quad M_i(y) = \alpha_i y + \beta_i$$

for a.e. $t \in [0, T]$, all $x, y \in \mathbb{R}$ and $i = 1, 2, \dots, m$. Let c be given by (2.2) and let $\beta_i \in (c(1 - \alpha_i), -c(1 - \alpha_i))$, $i = 1, 2, \dots, m$. Then the conditions of Theorem 2.1 are satisfied for $\nu = 2$ and the conditions of Theorem 2.2 are satisfied for $\nu = 1$ and $d_i = -c_i$, $i = 1, 2, \dots, m$. Since $c_i > 0$ and $\alpha_i \geq 0$, the functions J_i are increasing and M_i are nondecreasing on \mathbb{R} for $i = 1, 2, \dots, m$. Choose an arbitrary compact interval $K \subset \mathbb{R}$ and denote $\varkappa_1 = \max_{x \in K} (|x|^{2k+1})$ and $\varkappa_2 = \sum_{j=1}^{2k} (\|p_j\|_\infty \max_{x \in K} |x|^j)$. Then $|f(t, x, y)| \leq \omega(|y|) (|y| + h(t))$ for a.e. $t \in [0, T]$ and all $x \in K$, $|y| > 1$, where $\omega(s) = 1 + \|q_1\|_\infty + \|q_2\|_\infty s$ and $h(t) = \varkappa_1 |p_{2k+1}(t)| + \varkappa_2$. Thus, the existence of a solution to (1.1)–(1.3) follows by means of Theorem 3.1.

3.4 Example. Let $\gamma \in (0, \infty)$, $p \in \mathbb{L}_\infty[0, T]$, $q_1 \in \mathbb{L}[0, T]$, $q_2 \in \mathbb{L}_\infty[0, T]$, $q_1 \geq 0$, $q_2 \geq 0$ and $q_1 + q_2 \geq \gamma$ a.e. on $[0, T]$, $\varphi \in \mathbb{C}(\mathbb{R})$, $\lim_{|x| \rightarrow \infty} \varphi(x) = 0$. Consider the problem (1.1)–(1.3), where

$$f(t, x, y) = p(t) \varphi(x) + q_1(t) y + q_2(t) y^2 \operatorname{sgn} y \quad \text{for a.e. } t \in [0, T] \quad \text{and all } x, y \in \mathbb{R}$$

and $J_i(x)$ and $M_i(y)$, $i = 1, 2, \dots, m$, are given as in Example 3.3, but with $c_i \in (-\pi/2, 0)$ and $\alpha_i \in (1, \infty)$, $i = 1, 2, \dots, m$. Let c be given by (2.2) and let $\beta_i \in (c(1 - \alpha_i), c(\alpha_i - 1))$ for $i = 1, 2, \dots, m$. We have $c > 0$, $M_i(c) > c$, $M_i(-c) < -c$, $f(t, x, c) > 0$ and $f(t, x, -c) < 0$ for a.e. $t \in [0, T]$ and all $x \in \mathbb{R}$ with $|x|$ sufficiently large. Thus, the assumptions of Theorem 2.1 are satisfied for $\nu = 2$ and the assumptions of Theorem 2.2 are satisfied for $\nu = 1$ and $d_i = -c_i$, $i = 1, 2, \dots, m$. Furthermore, J_i are increasing and M_i are nondecreasing on \mathbb{R} for $i = 1, 2, \dots, m$. Since φ is bounded on \mathbb{R} , we can find $\omega \in \mathbb{C}([1, \infty))$ such that $\omega > 0$ on $[1, \infty)$, $\int_1^\infty ds/\omega(s) = \infty$ and $|f(t, x, y)| \leq \omega(|y|)|y|$ for a.e. $t \in [0, T]$ and all $x \in \mathbb{R}$, $|y| > 1$. Thus, by Theorem 3.1, the given problem has a solution.

3.5 Example. Let $\gamma \in (0, \infty)$, $p \in \mathbb{L}_\infty[0, T]$, $q \in \mathbb{L}[0, T]$, $q \geq \gamma$ a.e. on $[0, T]$, $\varphi \in \mathbb{C}(\mathbb{R})$, $\lim_{|x| \rightarrow \infty} \varphi(x) = 0$. Consider the problem (1.1)–(1.3), where

$$f(t, x, y) = p(t) \varphi(x) + \frac{q(t) y}{1 + y^2} \quad \text{for a.e. } t \in [0, T] \quad \text{and all } x, y \in \mathbb{R}$$

and $J_i(x)$ and $M_i(y)$, $i = 1, 2, \dots, m$, are given as in Example 3.4, but with $c_i \in (0, \infty)$. We can see that the assumptions of Theorem 2.1 are satisfied for $\nu = 1$ and the assumptions of Theorem 2.2 are satisfied for $\nu = 2$ and $d_i = -c_i$, $i = 1, 2, \dots, m$. As in the previous examples, J_i are increasing and M_i are nondecreasing on \mathbb{R} for $i = 1, 2, \dots, m$. Moreover, since the functions φ and $\frac{y}{y^2+1}$ are bounded on \mathbb{R} , we can find $h \in \mathbb{L}[0, T]$ such that $|f(t, x, y)| \leq h(t)$ for a.e. $t \in [0, T]$ and all $x, y \in \mathbb{R}$. Thus, by Theorem 3.2, our problem has a solution.

References

- [1] D. Bainov and P. Simeonov, *Impulsive Differential Equations: Periodic Solutions and Applications*. Longman Sci. Tech., Harlow, 1993.
- [2] DONG YUJUN. Periodic solutions for second order impulsive differential systems. *Nonlinear Anal.* **27** (1996), 811-820.
- [3] HU SHOUCHUAN AND V. LAKSMIKANTHAM. Periodic boundary value problems for second order impulsive differential systems. *Nonlinear Anal.* **13** (1989), 75-85.
- [4] E. LIZ AND J. J. NIETO. Periodic solutions of discontinuous impulsive differential systems. *J. Math. Anal. Appl.* **161** (1991), 388-394.
- [5] I. RACHŮNKOVÁ AND M. TVRDÝ. Nonmonotone impulse effects in second order periodic boundary value problems. *Abstr. Anal. Appl.* **2004**: 7, 577-590.
- [6] I. RACHŮNKOVÁ AND M. TVRDÝ. Non-ordered lower and upper functions in second order impulsive periodic problems. *Dyn. Contin. Discrete Impuls. Syst., Ser. A, Math. Anal.*, to appear.
- [7] I. RACHŮNKOVÁ AND M. TVRDÝ. Existence results for impulsive second order periodic problems. *Nonlinear Anal., Theory Methods Appl.* **59** (2004) 133-146.
- [8] ZHANG ZHITAO. Existence of solutions for second order impulsive differential equations. *Appl. Math., Ser. B (Engl. Ed.)* **12**, (1997), 307-320.

Irena Rachůnková, Department of Mathematics, Palacký University, 779 00 OLOMOUC, Tomkova 40, Czech Republic (e-mail: rachunko@inf.upol.cz)

Milan Tvrďý, Mathematical Institute, Academy of Sciences of the Czech Republic, 115 67 PRAHA 1, Žitná 25, Czech Republic (e-mail: tvrды@math.cas.cz)

(Received September 30, 2003)