

On the asymptotic behavior of solutions of nonlinear differential equations of integer and also of non-integer order

Milan Medved'

Department of Mathematical Analysis
and Numerical Mathematics,
Comenius University, 842 48 Bratislava, Slovakia
e-mail: medved@fmph.uniba.sk

Abstract

We present conditions under which all solutions of the fractional differential equation with the Caputo derivative

$${}^c D_a^\alpha x(t) = f(t, x(t)), \quad a > 1, \quad \alpha \in (1, 2), \quad (1)$$

are asymptotic to $at + b$ as $t \rightarrow \infty$ for some real numbers a, b .

AMS Classification: 34E10, 34A34

Keywords: Asymptotic behavior, Caputo derivative, fractional differential equation

This work was supported by the Slovak Research and Development Agency under the contract No. APVV-0134-10 and by the Slovak Grant agency, project No. 1/0507/11.

This paper is in final form, has been submitted to the Proceedings of 9th Colloquium on the Qualitative Theory of Differential Equations and no version of it will be submitted for publication for publication elsewhere.

1 Introduction

In the asymptotic theory of n -th order nonlinear ordinary differential equations

$$y^{(n)} = f(t, y, y', \dots, y^{(n-1)}) \quad (2)$$

the classical problem is to establish some conditions for the existence of a solution which approach to a polynomial of degree $1 \leq m \leq n-1$ as $t \rightarrow \infty$. The first paper concerning this problem was published by D. Caligo [3] in 1941. He proved that if

$$|A(t)| < \frac{k}{t^{2+\rho}} \quad (3)$$

for all large t , where k, ρ are given, then any solution $y(t)$ of the linear differential equation

$$y''(t) + A(t)y(t) = 0, \quad t > 0, \quad (4)$$

can be represented asymptotically as $y(t) = c_1 t + c_2 + o(1)$ when $t \rightarrow +\infty$, $c_1, c_2 \in \mathbb{R}$ (see [1]). The first paper on the nonlinear second order differential equations

$$y''(t) = f(t, y(t)) \quad (5)$$

was published by W. F. Trench [27] (1963). He proved a sufficient condition on the existence of a solution of the equation (5) which is asymptotic to $a + bt$ as $t \rightarrow +\infty$, for some real numbers a, b . Different conditions under which all solutions of the equation

$$y''(t) = f(t, y(t)) \quad (6)$$

is approaching to $a + bt$ as $t \rightarrow \infty$ for some real numbers a, b . The asymptotic behavior of solutions of this type of equation has been discussed by D. S. Cohen [5] (1967), J. Tong [26] (1982), T. Kusano and W. F. Trench [10] (1985) and [11] (1985) and others. This problem has been solved for the equation

$$y''(t) = f(t, y(t), y'(t)) \quad (7)$$

by F. M. Dannan [8] (1985), A. Constantin [6] (1993) and [7] (2005), Y. V. Rogovchenko [23] (1998), S. P. Rogovchenko [24] (2000), O. G. Mustafa, Y. V. Rogovchenko [19] (2002), O. Lipovan [12] (2003) and others. In the proofs of their results the key role plays the Bihari inequality (see [2]) which is a generalization of the Gronwall inequality. Some results on the existence of solutions of the n -th order differential equation

$$y^{(n)}(t) = f(t, y(t)) \quad (8)$$

approaching to a polynomial function of the degree m with $1 \leq m \leq n - 1$ are proved by Ch. G. Philos, I. K. Purnaras and P. Ch. Tsamatos [21] (2004). Their proofs are based on an application of the Schauder Fixed Point Theorem. The paper by R. P. Agarwal, S. D. Djebali, T. Moussaoui and O. G. Mustafa [1] (2007) surveys the literature concerning the topic in asymptotic integration theory of ordinary differential equations. Several conditions under which all solutions of the one dimensional p -Laplacian equation

$$(|y'|^{p-1}y')' = f(t, y, y'), \quad p > 1 \quad (9)$$

are asymptotic to $a + bt$ as $t \rightarrow \infty$ for some real numbers a, b are proved in [17] (2008) and some sufficient conditions for the existence of solutions of the equation

$$(\Phi(y^{(n)}))' = f(t, y), \quad n \geq 1, \quad (10)$$

where $\Phi: \mathbb{R} \rightarrow \mathbb{R}$ is an increasing homeomorphism with a locally Lipschitz inverse satisfying $\Phi(0) = 0$ are given in the paper [16] (2010).

The aim of this paper is to give some conditions under which all solutions of the fractional differential equation (1) are asymptotic to $a + bt$ as $t \rightarrow \infty$ for some real numbers a, b . The proof of this result is based on a desingularization method proposed by the author in the paper [14] (see also [15]) in the study of nonlinear integral inequalities with weakly singular kernels.

2 Fractional Differential equations with the Caputo's derivative

Consider the initial value problem

$${}^c D_a^\alpha x(t) = f(t, x(t)), \quad t \geq a > 1, \quad 1 < \alpha < 2, \quad (11)$$

$$x(a) = c_0, \quad x'(a) = c_2, \quad (12)$$

where

$${}^c D_a^\alpha x(t) := \frac{1}{\Gamma(2 - \alpha)} \int_a^t (t - s)^{\alpha-1} x''(s) ds \quad (13)$$

is the Caputo derivative of the order $\alpha \in (1, 2)$ of a C^2 -scalar valued function $x(t)$ defined on the interval $[a, \infty)$, $x''(t) = \frac{d^2 x(t)}{dt^2}$. This definition has been given by M. Caputo in the paper [4]. For the definition of the Caputo derivative of order $\alpha \in (n - 1, n)$, $n \geq 1$ see [20] and also the monographs [18], [25]. We assume that any solution $x(t)$ of this problem exists on the interval

$[a, \infty)$. One can show that the initial value problem (11), (12) is equivalent to the integral equation

$$x(t) = c_0 + c_1(t - a) + \frac{1}{\Gamma(\alpha)} \int_a^t (t - s)^{\alpha-1} f(s, x(s)) ds. \quad (14)$$

Since $\alpha > 1$ the function $x(t)$ is differentiable and therefore (14) yields

$$x'(t) = c_1 + \frac{1}{\Gamma(\alpha - 1)} \int_a^t (t - s)^{\alpha-2} f(s, x(s)) ds. \quad (15)$$

Lemma 1 (see [22], [13]) *Let β, γ and p be positive constants such that $p(\beta - 1) + 1 > 0$, $p(\beta - 1) + 1 > 0$. Then*

$$\int_0^t (t - s)^{p(\beta-1)} s^{p(\gamma-1)} ds = t^\Theta B, \quad t \geq 0 \quad (16)$$

where $B := B[p(\gamma - 1) + 1, p(\beta - 1) + 1]$, $B[\xi, \eta] = \int_a^1 s^{\xi-1} (1 - s)^{\eta-1} ds$, ($\xi > 0, \eta > 0$) and $\Theta = p(\beta + \gamma - 2) + 1$.

Theorem 1 *Suppose that $1 < \alpha < 2$, $p > 1$, $p(\alpha - 2) + 1 > 0$, $a > 1$, $q = \frac{p}{p-1}$ and the function $f(t, u)$ satisfies the following conditions:*

- (i) $f(t, u)$ is continuous in $D = \{(t, v) : t \in [0, \infty), v \in \mathbb{R}\}$;
- (ii) There are continuous nonnegative functions $h : R_+ := [0, \infty) \rightarrow R_+$, $g : R_+ \rightarrow R_+$, and $\gamma > 0$ with $p(\gamma - 1) + 1 > 0$ such that

$$|f(t, x)| \leq t^{\gamma-1} h(t) g\left(\frac{|x|}{t}\right), \quad t > 0, \quad (t, x) \in D, \quad (17)$$

where $\gamma = 3 - \alpha + \frac{1}{p}$, i.e. $\Theta := p(\alpha + \gamma - 3) + 1 = 0$ and

$$\int_a^\infty h(s)^q ds < \infty. \quad (18)$$

(iii)

$$\int_a^\infty \frac{\tau^{q-1} d\tau}{g(\tau)^q} = \infty. \quad (19)$$

Then every solution $x(t)$ of the equation (11) is asymptotic to $c + dt$ for $t \rightarrow \infty$, where $c, d \in \mathbb{R}$.

Proof. By applying the condition (17) we obtain from the (14) and (15)

$$|x(t)| \leq Ct + B_1 \int_a^t (t-s)^{\alpha-1} h(s) s^{\gamma-1} g\left(\frac{|x|}{s}\right) ds, \quad (20)$$

where $C = |c_1| + |c_2|$, $B_1 = \frac{1}{\Gamma(\alpha)}$, i. e.

$$\begin{aligned} |x(t)| &\leq Ct + B_1 \int_a^t (t-s)^{\alpha-1} h(s) s^{\gamma-1} g\left(\frac{|x|}{s}\right) ds \\ &\leq Ct + B(t-a) \int_a^t (t-s)^{\alpha-2} h(s) s^{\gamma-1} g\left(\frac{|x|}{s}\right) ds \end{aligned} \quad (21)$$

This yields the inequality

$$\frac{|x(t)|}{t} \leq C + B \int_a^t (t-s)^{\alpha-2} h(s) s^{\gamma-1} g\left(\frac{|x|}{s}\right) ds. \quad (22)$$

If we denote by $z(t)$ the right-hand side of the inequality (22) we obtain the inequalities:

$$\frac{|x(t)|}{t} \leq z(t), \quad (23)$$

$$|x'(t)| \leq z(t). \quad (24)$$

Since the function g is nondecreasing, the inequality (23) yields

$$g\left(\frac{|x(t)|}{t}\right) \leq g(z(t)) \quad (25)$$

and from (22) we obtain

$$z(t) \leq 1 + C + B_1 \int_a^t (t-s)^{\beta-1} h(s) s^{\gamma-1} k(s) g(z(s)) ds, \quad (26)$$

$0 < \beta = \alpha - 1 < 1$. By applying the Hölder inequality and Lemma 1 we obtain

$$\begin{aligned} &\int_a^t (t-s)^{\beta-1} s^{\gamma-1} h(s) g(z(s)) ds \\ &\leq \left(\int_a^t (t-s)^{p(\beta-1)} s^{p(\gamma-1)} ds \right)^{\frac{1}{p}} \left(\int_a^t h(s)^q g(z(s))^q ds \right)^{\frac{1}{q}} \\ &\leq \left(\int_0^t (t-s)^{p(\beta-1)} s^{p(\gamma-1)} ds \right)^{\frac{1}{p}} \left(\int_a^t h(s)^q g(z(s))^q ds \right)^{\frac{1}{q}} \\ &\leq BB_1 t^\Theta \left(\int_a^t h(s)^q g(z(s))^q ds \right)^{\frac{1}{q}} \leq BB_1 t^\Theta \left(\int_a^t h(s)^q g(z(s))^q ds \right)^{\frac{1}{q}}, \end{aligned}$$

where $B = B[p(\gamma-1)+1, p(\beta-1)+1]$, $\Theta = p(\beta+\gamma-2)+1 = p(\alpha+\gamma-3)+1 = 0$, i.e.

$$\int_a^t (t-s)^{\beta-1} s^{\gamma-1} h(s) g(z(s)) ds \leq BB_1 \left(\int_a^t h(s)^q g(z(s))^q ds \right)^{\frac{1}{q}}. \quad (27)$$

Using this inequality and the elementary inequality $(a+b)^q \leq 2^{q-1}(a^q + b^q)$, $a, b \geq 0$ we obtain from (26)

$$z(t)^q \leq 2^{q-1}[(1+C)^q + (BB_1) \int_a^t k(s)^q g(z(s))^q ds]. \quad (28)$$

If we denote $u(t) = z(t)^q$, i.e. $z(t) = u(t)^{\frac{1}{q}}$, $P_1 = 2^{q-1}[(1+c)^q]$, $Q_1 = 2^{q-1}(B_1)^q$ then

$$u(t) \leq P_1 + Q_1 \int_a^t h(s)^q g(u(t)^{\frac{1}{q}})^q ds, \quad t \geq a. \quad (29)$$

Denote

$$\omega(v) = g(v^{\frac{1}{q}})^q, \quad \Omega(u) = \int_{u_0}^u \frac{d\sigma}{\omega(\sigma)}, \quad u_0 = u(a). \quad (30)$$

Since $\Omega(u) = q \int_{v_0}^v \frac{\tau^{q-1} d\tau}{g(\tau)^q}$, where $v_0 = (u_0)^{\frac{1}{q}}$, $v = u^{\frac{1}{q}}$ the condition (iii) of Theorem 1 implies that $\lim_{u \rightarrow \infty} \Omega(u) = \infty$, i.e. $\Omega([u_0, \infty)) = [0, \infty)$ then by the Bihari lemma

$$u(t) \leq K_0 := \Omega^{-1}[\Omega(P_1) + Q_1 \int_a^\infty h(s)^q ds] < \infty. \quad (31)$$

Since $u(t) = z(t)^q$ we obtain that $z(t) \leq K := K_0^{\frac{1}{q}}$ and from (24), (25) we have

$$\frac{|x(t)|}{t} \leq K_1, \quad |x'(t)| \leq K_1, \quad t \geq a. \quad (32)$$

From the condition (ii) of Theorem 1 we have

$$\begin{aligned} \int_a^t (t-s)^{\beta-1} |f(s, x(s))| ds &\leq \int_a^t (t-s)^{\beta-1} s^{\gamma-1} h(s) g(z(s)) ds \\ &\leq z(t) \leq K_1, \quad t \geq a, \end{aligned} \quad (33)$$

therefore $\int_a^\infty (t-s)^{\beta-1} |f(s, x(s))| ds$ exists. Therefore from the equality (15) it follows that $\lim_{t \rightarrow \infty} x'(t) = d$ exists and by the l'Hospital rule we conclude that

$$\lim_{t \rightarrow \infty} \frac{u(t)}{t} = \lim_{t \rightarrow \infty} x'(t) = d, \quad (34)$$

so the proof is now complete.

3 Example

Let $p > 1$, $\alpha = 2 - \frac{1}{2p}$, $\gamma = 3 - \alpha - \frac{1}{p}$, i.e. $p(\alpha + \gamma - 3) + 1 = 0$, $q = \frac{p}{p-1}$, i.e. $\frac{1}{p} + \frac{1}{q} = 1$, $g(u) = u^{\frac{q-1}{q}} [\ln(2+u)]^{\frac{1}{q}}$, $u \geq 0$, $f(t, u) = t^{\gamma-1} h(t) g(\frac{u}{t})$, where $h: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a continuous function with $\int_a^\infty h(s)^q ds < \infty$. Obviously $1 < \alpha < 2$, $p(\alpha - 2) + 1 = \frac{1}{2}$, $p(\gamma - 1) + 1 = 1 - \frac{1}{2p} > 0$ and

$$\int_a^\infty \frac{\tau^{q-1}}{g(\tau)^q} d\tau = \int_a^\infty \frac{1}{\ln(2+\tau)} d\tau = \infty. \quad (35)$$

Therefore the function $f(t, u)$ satisfy the conditions (i)–(iii) of Theorem 1.

References

- [1] R. P. Agarwal, S. Djebali, T. Moussaoui, O. G. Mustafa, On the asymptotic integration of nonlinear differential equations, *Journal of Comput. and Applied Math.*, 202 (2007), 352–376.
- [2] I. Bihari, Researches of the boundedness and stabilitz of solutions of non-linear differential equations, *Acta Math. Acad. Sci. Hung.*, Vol. 8 (1957), 261–278.
- [3] D. Caligo, Comportamento asintotico degli integrali dell'equazione $y''(x) + A(x)y(x) = 0$, nell'ipotesi $\lim_{x \rightarrow +\infty} A(x) = 0$, *Boll. Un. Mat. Ital.*, Vol. 6(1941), 286–295.
- [4] M. Caputo, Linear models of dissipation whose Q is almost frequency independent II, *Geophys. J. Royal Astronom. Soc.*, Vol. 13 (1967), 529–535.
- [5] D. S. Cohen, The asymptotic behavior of a class of nonlinear differential equations, *Proc. Amer. Math. Soc.* Vol 18, (1967), 607–609.
- [6] A. Constantin, On the asymptotic behavior of second order nonlinear differential equations, *Rend. Math. Appl.*, Vol. 13, No. 7 (1993), 627–634.
- [7] A. Constantin, On the existence of positive solutions of second order differential equations, *Annali di Matematica*, Vol. 184 (2005), 131–138.
- [8] F. M. Dannan, Integral inequalities of Gronwall-Bellman-Bihari type and asymptotic behavior of certain second order nonlinear differential equations, *J. Math. Anal. Appl.*, Vol. 108 (1985), 151–164.

- [9] K. M. Furati, N.-E. Tatar, Power-type estimates for nonlinear fractional differential equations, *Nonlinear Anal.*, TMA 62(2001) 1025–1036.
- [10] T. Kusano, W. F. Trench, Global existence of second order differential equations with integrable coefficients, *J. London Math. Soc.*, Vol. 31 (1985), 478–486.
- [11] T. Kusano, W. F. Trench, Existence of global solutions with prescribed asymptotic behavior for nonlinear ordinary differential equations, *Mat. Pura Appl. Vol.* 142(1985), 381–392.
- [12] O. Lipovan, On the asymptotic behaviour of the solutions to a class of second order nonlinear differential equations, *Glasg. Math. J.* 45 no. 1, (2003), 179–187.
- [13] Q.-H. Ma, J. Pečarič, J.-M. Zhang, Integral inequalities of systems and the estimate for solutions of certain nonlinear two-dimensional fractional differential systems, *Computers and Mathematics with Applications*, Vol. 61 (2011), 3258–3267.
- [14] M. Medved', A new approach to an analysis of Henry type integral inequalities and their Bihari type versions, *J. Math. Anal. Appl.*, Vol. 214 (1997), 349–366.
- [15] M. Medved', Integral inequalities and global solutions of semilinear evolution equations, *J. Math. Anal. Appl.*, Vol. 37, No. 4 (2002), 871–882.
- [16] M. Medved', T. Moussaoui, Asymptotic integration of nonlinear Φ -Laplacian differential equations, *Nonlinear Analysis, TMA*, Vol. 72 (2010), 1–8.
- [17] M. Medved', E. Pekárková, Long time behavior of second order differential equations with p -Laplacian, *Electronic Journal of Differential Equations*, Vol. 2008(108) (2008), 1–12.
- [18] K. S. Miller, B. Ross, *An Introduction to the Fractional Calculus and Differential Equations*, John Wiley, New York 1993.
- [19] O. G. Mustafa, Y. V. Rogovchenko, Global existence of solutions with prescribed asymptotic behavior for second-order nonlinear differential equations, *Nonl. Anal. TMA*, Vol. 51 (2002), 339–368.
- [20] I. Podlubny, *Fractional Differential Equations*, Academic Press, San Diego 1999.

- [21] Ch. G. Philos, I. K. Purnaras, P. Ch. Tsamatos, Large time asymptotic to polynomials solutions for nonlinear differential equations, *Nonl. Anal. TMA*, Vol. 59 (2004), 1157–1179.
- [22] A. P. Prudnikov, Zu. A. Brychkov, O. I. Marichev, Integral and Series, in: *Elementary Functions*, Vol. 1, Nauka, Moscow, 1981 (in Russian).
- [23] Y. V. Rogovchenko, On asymptotics behavior of solutions for a class of second order nonlinear differential equations, *Collect. Math.*, Vol. 49, No.1 (1998), 113–120.
- [24] S. P. Rogovchenko, Yu. V. Rogovchenko, Asymptotics of solutions for a class of second order nonlinear differential equations, *Portugaliae Math.*, Vol. 57, No. 1 (2000), 17–32.
- [25] S. G. Samko, A. A. Kilbas, O. I. Marichev, *Fractional Integrals and Derivatives: Theory and Applications*, Gordon and Breach Science Publishers, New York 1993.
- [26] J. Tong, The asymptotic behavior of a class of nonlinear differential equations of second order, *Proc. Amer. Math. Soc.*, Vol. 84 (1982), 235–236.
- [27] W. F. Trench, On the asymptotic behavior of solutions of second order linear differential equations, *Proc. Amer. Math. Soc.*, Vol. 54 (1963), 12–14.

(Received July 31, 2011)