

COMPREHENSIVE EVALUATION OF THE RESULTS OF THE DAPHNIA TEST CARRIED OUT AT THE TISZA-SECTION AND MAJOR DISTRICT WATERS IN SZOLNOK COUNTY (1977—87)

F. CSÉPAI

Service of Public Health and Epidemiology of County Szolnok
Water Microbiological Laboratory, Szolnok, Hungary

(Received December 22, 1987)

Abstract

Between 1977—1987 acute toxicological studies were carried out with *Daphnia magna* at eight points of the Tisza-section in Szolnok county and at the major district waters of the region. It was determined that in the periods studied 19,4% of the Tisza water samples and 4,8% of the samples originating from the district waters were of toxic effect on the *Daphnia*. From 1975 on the positivity of the Tisza water samples taken frequently above the area of the Surface Water Works in Szolnok showed strongly decreasing tendency, falling back from 74% to 1%. The observed phenomenon can be explained most probably by the beneficial effect of the Kisköre storage lake, since the drastic decrease of the high rate of objection observed previously coincided with the filling up of the lake between 1979—1983.

Introduction

Since 1974 the Water Microbiological Laboratory of the Service of Public Health and Epidemiology of Szolnok County carries out regular chemical, bacteriological, biological and toxicological studies with the aim to reveal the environmental effects. Within the complex hygienic programme the importance of these acute toxicological studies was emphasized several times, in the frame of which biological tests were accomplished in respect to the Tisza river in Szolnok county, and the major district waters of the region, as well as the drinking water of the city of Szolnok, provided from the Tisza river. Here we would refer to some of the data published earlier on this topic (CSÉPAI 1975, 1976, SCHIEFNER 1979, KÁDÁR 1983). In the following a review is given on the results of the *Daphnia* toxicity tests carried out in the period between 1977 and November 30, 1983.

Materials and Methods

The water samples were taken from the sites given in Fig. 1 and the tables, 5 cm below the surface, transported in refrigerated state and processed within 24 hours after temperation to room temperature. The *Daphnia* toxicity tests were performed and the results evaluated according to the specifications of the Hungarian *Daphnia* test (Water Toxicological Studies, 1982). The principle




Fig. 1. Regional distribution of the sampling sites

of the test is that for negative qualify those samples in which half of the *Daphnia* survive a 48 hours' period of exposure; in the opposite case, the result of the *Daphnia* test is regarded as positive.

The *Daphnia* toxicity tests were carried out using 4 days-old organisms of the strain *Daphnia magna S* cultivated in our laboratory. The sensitivity of *Daphnia* to chromate corresponded to the requirements of the Hungarian *Daphnia* test mentioned above.

Results and Evaluation

In connection with the results to be described below, we should mention that the findings of the tests carried out in the period 1974–1978 at the Tisza-section in Szolnok county have been previously discussed by SCHIEFNER (1979) as a part of the studies performed on the whole length of the river on the territory of Hungary.

According to our results, within the period studied 19,4% of the total samples taken from the Tisza-section in Szolnok county proved to be *Daphnia* positive (Table 1). The data show that the river water is toxic for *Daphnia* primarily in autumn, winter and early spring months. These findings prove the role played by the temperature factors in the realization of the effect of the toxic microcontaminants occurring in the Tisza water. The toxicity of the district waters was low: the positive samples amount to 4,8% of the total samples studied (Table 2). From the point of view of the seasonal distribution of the positive results, the relationship is similar to that experienced during the course of the studies at the Tisza river.

Tests of the unrefined surface water obtained by the Szolnok Surface Water Works were carried out by us at least once, more often twice a week. The results are shown in Fig. 2.

Table 1. Months

Sampling sites Tisza		I.	II.	III.	IV.	V.	VI.	VII.	VIII.	IX.	X.	XI.	XII.	1977—1987 Total
Tiszasüly right bank	positive negative	2 1	2 5	1 2	— 9	— 1	— 10	— —	2 8	1 —	— 10	1 —	1 7	10 52 62
Nagykörű right bank	positive negative	2 —	1 2	1 1	— 3	— 1	— 4	— —	2 2	— 1	1 3	1 —	1 2	9 19 28
Unrefined water obtained by Szolnok Water Works	positive negative	2 —	3 4	1 2	— 9	— 1	— 10	— —	3 7	1 —	2 8	1 —	2 6	15 47 62
Szolnok left bank below Zagyva	positive negative	2 —	2 4	1 2	— 9	— 1	— 9	— —	2 8	1 —	2 8	1 —	2 6	13 47 60
Szolnok right bank below Zagyva	positive negative	2 —	1 1	1 1	— 3	— 1	1 4	— —	2 2	1 —	2 2	1 —	2 1	13 15 28
Tiszaföldvár left bank	positive negative	— 1	2 4	1 2	— 9	— —	— 10	— —	1 9	— 1	2 8	— —	— 7	7 51 58
Tiszavárkony right bank	positive negative	1 —	2 4	— 3	— 9	— 1	— 10	— —	1 9	— 1	2 8	— —	1 8	7 53 60
Tiszazug, from the bridge, middle line	positive negative	1 —	1 6	1 2	— 9	— —	— 10	— —	1 9	— 1	2 8	— —	1 7	7 52 59
Positive samples		12	14	7	—	—	1	—	14	4	13	8	8	81 positive (19,4%)
Negative samples		1	30	15	60	6	67	—	54	4	55	—	44	336 negative
Total		13	44	22	60	6	68	—	68	8	68	8	52	417 total

Table 2. Months

Sampling sites (in every case from the bridge, middle line)		I.	II.	III.	IV.	V.	VI.	VII.	VIII.	IX.	X.	XI.	XII.	1977—1987 Total	
Galga above Jászfényszaru	positive negative	—	3	1	9	—	3	1	7	2	5	2	8	1	3 51 54
Tarna above Jászjákóhalma	positive negative	—	3	—	1	—	9	—	3	8	—	2	5	— 4 50 54	
Zagyva below Jászfényszaru	positive negative	—	3	—	1	—	9	—	2	7	—	2	5	— 3 50 53	
Zagyva below Jászberény	positive negative	—	—	—	1	—	—	—	2	—	—	1	—	— 3 17 20	
Zagyva below Jásztelek	positive negative	—	3	—	1	—	8	—	3	8	—	2	5	— 1 52 53	
Zagyva below Újsász	positive negative	—	—	—	1	—	—	—	2	—	—	1	—	— 3 16 19	
Zagyva Szolnok, below TB Hospital	positive negative	—	3	—	1	—	9	—	3	7	—	2	5	— 3 51 54	
Miller above Szolnok	positive negative	—	2	—	1	—	8	—	3	7	—	2	5	— 1 51 52	
Körös Kunszentmárton	positive negative	—	3	—	1	—	9	—	3	8	—	2	5	— — 54 54	
Hortobágy-Berettyó positive below Mezőtúr	negative	1	—	—	—	1	—	—	—	2	—	—	10	— 3 51 54	
Hortobágy-Berettyó positive below Türkéve	negative	—	3	—	1	—	8	—	3	7	—	2	5	— — 1 51 52	
Positive samples		1	—	—	—	8	—	—	—	12	—	1	3	25 positive (4,8%)	
Negative samples		25	11	83	30	70	20	45	20	86	9	95	494 negative		




Fig. 2. The *Daphnia*-test positive results in the samples of unrefined water obtained by the Szolnok Surface Water Works between 1977—1987

The aim of the biological tests is to reveal the toxic effects (microcontaminations, unknown interactive effects) which can not be detected by analytical methods. For these reasons we are not in the position to identify the chemical background of the observed phenomena. However, we feel that the reasons for the significant decrease of toxicity reflected by our results (Fig. 2) can be satisfactorily interpreted. Namely, the drastic improvement coincides with the filling up of the Kisköre storage lake (lately named Tisza-lake) to its present level in the period 1979—1982. The significant drop in the toxicity of the storage lake can be due to the simultaneous effect of the favourable sedimentation originating from the decrease of the flow-rate, the increase in the intensity of the volumetric irradiation by the sunlight and in the oxygen uptake from the air due to the increased surface, and finally to the filtering effect of the aquatic vegetation. Our calculations based on the available data (volume and surface of the stored water, average water output) show that the operation of the storage lake led to 3,9-fold increase in the specific surface per unit volume of the water, and 3,1-fold decrease in the average flow-rate as compared to that of the river. Although quantitative estimation can not be given, an inspection of the storage lake shows a convincing increase in the vegetation.

References

- Csépai, F. (1975): The using of *Daphnia* in the judgement of the degree of toxicity of surface water and chlorinated drinking water. I. Cultivation and maintenance of testorganisms. — Budapest Public Health 4, 119.
- Csépai, F. (1975): The using of *Daphnia* in the judgement of the degree of toxicity of surface water and chlorinated drinking water. II. Test-method and results. — Budapest Public Health, I, 15—18.
- Csépai, F. (1984): Summarizing evaluation of the results of the *Daphnia* test carried out at the Tisza section and major district waters at Szolnok County (1977—1983). — Tiscia (Szeged) 19, 17—20.

- KÁDÁR, Gy., CSÉPAI, F., PALICSKA, J. and TASNÁDI, M.-NÉ (1983): Connection between the characteristics of the surface waters of the Tisza river and the drinking water of Szolnok city on the basis of a 10-years-study. — Lecture presented at the Public Health Days alongside the Tisza river, IV., August 24—25, 1983.
- SCHIEFNER, K., BODNÁR, I.-NÉ, NAGY, G., GÁBOS, L.-NÉ, TÓTH, J.-NÉ, ANDRIK, P., VUKOVICH, F., MÉSZÁROS, D., ESTÓK, B., LUCSKAY, K., STECZ, J., CSÉPAI, F., KÁDÁR, Gy., PALICSKA, J., FODRÉ, Zs. és HEGEDÜS, M. (1979): Experiences of the team formed in connection with the establishment of the water storage tank at Kisköre. (1974—1978). — Lecture presented at the 20th Public Health Days organized at Lake Balaton May 3—5, 1979.
- Water Toxicological Studies, *Daphnia* test (1982): Hungarian Peoples Republic National Standard, 22 902/6—81.

A Szolnok megyei Tisza-szakasz és a fontosabb mellékvizek Daphnia-teszt eredményeinek összefoglaló értékelése (1977—1987)

F. CSÉPAI

Service of Public Health and Epidemiology of County Szolnok
Water Micro-biological Laboratory, Szolnok, Hungary

Kivonat

1977—1987 között akut toxikológiai vizsgálatokat végeztünk Daphniákkal a Tisza Szolnok megyei szakaszának 6 pontján és a megye fontosabb mellékvizein. Megállapítottuk, hogy a vizsgált időszakban a Tisza hossz-szelvénny víz mintáinak 18,3%-a, a mellékvizekből származó minták 4,3%-at volt Daphniáakra mérgező hatású. A szolnoki felszíni vízmű felett nagy gyakorisággal vett Tisza víz minták pozitivitása 1975. után napjainkig erősen csökkenő tendenciát mutat; 74%-ról 1%-ra esett vissza. E jelenség okai nagy valószínűséggel olyan kedvező hatások érvényesülésével magyarázhatók, ahol az összetevők között jelentős szerepe van a Kiskörei víztározónak, mert környezeti adottságai tévében nagy mértékben elősegít a szerves- és toxikus anyagok kémiai és mikrobiológiai oxidációját. De nem hagyhatjuk figyelem kívül a környezetvédelem területén hozott állami intézkedések hatását sem az ugyancsak lassan, de a társadalmi céloknak megfelelő irányban változó lakossági szemlélettel együtt.

Общая оценка результатов тестов, основанных на *Daphnia*, воды реки Тиса и ее основных притоков, протекающих по территории области Солнок (1977—1987 гг.)

Ф. Чепай

Станция здравоохранения и эпидемиологии обл. Солнок,
Микробиологическая лаборатория, г. Солнок

Резюме

В период 1977—87 гг. проводились токсикологические исследования при помощи теста на *Daphnia magna* в восьми участках Тисы и ее основных притоков, протекающих по территории области Солнок. Было установлено, что за исследуемый период 19,4% водных проб продольного отрезка Тисы и 4,8% проб ее притоков оказались токсичными для *Daphnia*.

Пробы, которые брали с большой частотой над водонапорной станцией в Солноке свидетельствуют о тенденции значительного снижения позитивности водных проб Тисы с 1975 г. до настоящего времени (снижение с 74% до 1%). Предполагается, что возможным объяснением наблюдаемых результатов является благоприятное влияние водохранилища в Кишкёре, поскольку резкое снижение наблюдавшего ранее высокого процента позитивности совпало по времени с эксплуатационным заполнением водохранилища в Кишкёре в период 1979—1983 гг.

Ukratko procenjivanje rezultate test-Daphnie vršenim u reci Tisa i u važnijim sporednim vodama u županiji Solnok

F. ČEPAI

Abstrakt

Uzvrišili smo akutne toksikološke istraživanje sa Daphnijama na 6 tačci Tise u županiji Solnok i u važnijim sporednim vodama između 1977—87 god. Došli smo do zaključka da za vreme istraživanja Tise je bila toksična u 18,3% na Daphnije a sporedne vode u 4,3%.

Od 1975. godine pozitivitet izvađenog vodenog uzorka iz Tise kod Solnoškog hidrocentrala pokazuje opadanje tendencije, pad je od 74% do 1%. Ova pojавa opravda se sa takvim povoljnim uslovima gde između ostalim gradientima Kiškere Rezervoar za Vodu igra veliku ulogu, on ima takvih ekoloških okolnosti koji omogućavaju što bolju hemijsku i mikrobiološku oksidaciju organskim i toksičnim materijama. Ali dā ne izostavimo državne odredbe za odbranu ekoloških okolnosti koje zajedno sa mentalitetom stanovništva lagano ali sigurno se menjaju prema društvenim ciljima.

Table 1. Results of the Daphnia-test carried out at the Tisza-section in Szolnok County between 1977—1987

Table 2. Results of the Daphnia-test carried out at the district waters of Szolnok County between 1977—1987